



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 06:54 PM UTC

PDB ID : 3HPT / pdb_00003hpt
Title : Crystal structure of human FxA in complex with (S)-2-cyano-1-(2-methylbenzofuran-5-yl)-3-(2-oxo-1-(2-oxo-2-(pyrrolidin-1-yl)ethyl)azepan-3-yl)guanidine
Authors : Klei, H.E.; Ghosh, K.; Rushith, A.; Kish, K.
Deposited on : 2009-06-04
Resolution : 2.19 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.19 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	94	7% 74% 14% • 11%
1	C	94	13% 88% • 7%
2	B	238	% 85% 13% •
2	D	238	2% 89% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	21	-	-	X	-
9	DMS	B	251	-	-	X	-

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 5683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Coagulation factor X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	84	Total	C	N	O	S	0	1	0
			624	375	108	128	13			
1	C	87	Total	C	N	O	S	0	0	1
			604	364	98	129	13			

- Molecule 2 is a protein called Coagulation factor X.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	238	Total	C	N	O	S	0	2	0
			1861	1173	324	350	14			
2	D	237	Total	C	N	O	S	0	5	0
			1878	1185	325	354	14			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 1-cyano-2-(2-methyl-1-benzofuran-5-yl)-3-[(3S)-2-oxo-1-(2-oxo-2-pyrrolidin-1-ylethyl)azepan-3-yl]guanidine (CCD ID: YET) (formula: $C_{23}H_{28}N_6O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			32	23	6	3		
4	D	1	Total	C	N	O	0	0
			32	23	6	3		


- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		
5	D	1	Total	Ca	0	0
			1	1		

- Molecule 6 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Na	0	0
			1	1		
6	D	1	Total	Na	0	0
			1	1		

- Molecule 7 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
7	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 8 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	B	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		
8	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	B	1	Total	C	O	S	0	0
			4	2	1	1		
9	D	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	71	Total	O	0	0
			71	71		
10	B	195	Total	O	0	0
			195	195		
10	C	63	Total	O	0	0
			63	63		
10	D	225	Total	O	0	0
			225	225		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Coagulation factor X





- Molecule 1: Coagulation factor X



- Molecule 2: Coagulation factor X



- Molecule 2: Coagulation factor X



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.98Å 78.66Å 73.67Å 90.00° 102.49° 90.00°	Depositor
Resolution (Å)	50.00 – 2.19 50.00 – 2.19	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-2.19) 99.7 (50.00-2.19)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 2.18Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.176 , 0.239 (Not available) , 0.237	Depositor DCC
R_{free} test set	1781 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.1	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5683	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, DMS, MES, GOL, NA, YET, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.50	0/635	0.77	0/853
1	C	0.52	0/612	0.83	0/826
2	B	1.54	2/1906 (0.1%)	0.88	2/2572 (0.1%)
2	D	1.16	2/1930 (0.1%)	0.90	5/2599 (0.2%)
All	All	1.21	4/5083 (0.1%)	0.87	7/6850 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	145	HIS	C-N	44.99	1.92	1.33
2	D	218	GLY	C-N	44.26	1.97	1.33
2	B	218	GLY	C-N	42.83	1.90	1.33
2	D	160	VAL	CA-CB	5.76	1.58	1.54

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	218	GLY	O-C-N	-16.48	106.08	123.05
2	B	145	HIS	O-C-N	-13.12	108.50	123.33
2	B	218	GLY	O-C-N	-10.94	108.47	122.70
2	D	218	GLY	CA-C-N	-8.18	110.68	122.94
2	D	218	GLY	C-N-CA	-8.18	110.68	122.94
2	D	197	GLY	CA-C-N	5.56	125.32	119.76
2	D	197	GLY	C-N-CA	5.56	125.32	119.76

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	624	0	551	16	0
1	C	604	0	503	3	0
2	B	1861	0	1790	20	0
2	D	1878	0	1813	16	0
3	A	12	0	16	4	0
3	B	6	0	8	0	0
3	D	12	0	16	0	0
4	B	32	0	27	2	0
4	D	32	0	27	2	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
6	B	1	0	0	0	0
6	D	1	0	0	0	0
7	B	12	0	12	1	0
7	D	12	0	12	0	0
8	B	16	0	12	0	0
8	D	16	0	12	0	0
9	B	4	0	6	5	0
9	D	4	0	6	2	0
10	A	71	0	0	0	0
10	B	195	0	0	6	0
10	C	63	0	0	1	0
10	D	225	0	0	3	0
All	All	5683	0	4811	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:218:GLY:C	2:B:220:CYS:N	1.90	1.27
2:B:145:HIS:C	2:B:147:GLU:N	1.92	1.26
2:D:218:GLY:C	2:D:220:CYS:N	1.97	1.21
9:B:251:DMS:H12	10:B:326:HOH:O	1.64	0.94

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:158:LEU:HD23	9:B:251:DMS:H11	1.57	0.87
1:C:141:HIS:HD2	10:C:184:HOH:O	1.69	0.75
1:A:174:LYS:HG3	3:A:21:GOL:H31	1.68	0.73
2:D:36:GLU:H	2:D:36:GLU:CD	1.98	0.72
1:A:174:LYS:HB3	3:A:21:GOL:H2	1.74	0.70
1:A:145:ASN:HD22	2:D:63:ARG:NH2	1.95	0.65
1:A:145:ASN:HD22	2:D:63:ARG:HH22	1.43	0.64
2:B:72:ASN:HA	2:B:153:THR:O	1.98	0.64
1:A:174:LYS:HG3	3:A:21:GOL:C3	2.32	0.59
4:B:1:YET:HN29	4:B:1:YET:C2	2.15	0.59
2:D:147:GLU:OE2	2:D:222:ARG:NH1	2.35	0.57
9:D:250:DMS:H12	10:D:346:HOH:O	2.05	0.56
1:A:174:LYS:NZ	2:B:206:THR:OG1	2.39	0.55
1:A:153[A]:ARG:HG2	10:B:321:HOH:O	2.09	0.53
1:A:103:ASP:HA	1:A:108:TYR:HA	1.91	0.53
4:D:2:YET:C2	4:D:2:YET:HN29	2.22	0.52
2:B:171:SER:O	2:B:224:LYS:NZ	2.39	0.51
2:B:158:LEU:HD21	2:B:188:GLU:HB3	1.93	0.51
1:A:98:GLN:HB2	1:A:113:LEU:HD11	1.93	0.51
2:B:233:ALA:O	7:B:9:MES:H31	2.12	0.50
2:B:236:LYS:HG2	2:B:240:ARG:NH1	2.27	0.49
1:A:174:LYS:CB	3:A:21:GOL:H2	2.42	0.49
9:B:251:DMS:H13	10:B:346:HOH:O	2.14	0.48
2:B:157:MET:HE2	9:B:251:DMS:H22	1.96	0.48
1:A:93:SER:N	1:A:94:PRO:HD3	2.29	0.47
2:D:218:GLY:C	2:D:220:CYS:CA	2.82	0.47
2:B:183:ALA:HB3	2:B:228:TYR:CE1	2.51	0.46
2:B:71:ARG:O	2:B:154:ARG:HA	2.14	0.46
2:B:90:LYS:HE2	10:B:355:HOH:O	2.15	0.46
1:A:93:SER:N	1:A:94:PRO:CD	2.80	0.45
9:B:251:DMS:C1	10:B:326:HOH:O	2.42	0.45
2:B:16:ILE:O	2:B:144:THR:HA	2.17	0.45
2:B:124:PRO:HB2	2:B:128:ALA:HB2	1.99	0.45
2:D:27:CYS:H	2:D:71[B]:ARG:HH22	1.64	0.44
2:B:145:HIS:HB3	2:B:149:GLY:HA3	2.00	0.43
2:D:20:GLN:HG3	9:D:250:DMS:H23	2.00	0.43
4:B:1:YET:C2	4:B:1:YET:N29	2.78	0.43
1:C:141:HIS:HE1	1:C:143:GLU:OE1	2.02	0.43
1:A:155:TYR:CZ	1:A:171:PRO:HB2	2.53	0.43
2:B:240:ARG:NH2	10:B:423:HOH:O	2.52	0.43
2:D:25:GLY:N	2:D:117:ASN:OD1	2.37	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:155:TYR:CZ	1:C:171:PRO:HB2	2.54	0.42
2:D:23:LYS:O	2:D:71[B]:ARG:NH1	2.52	0.42
2:D:72:ASN:HA	2:D:153:THR:O	2.17	0.42
2:B:143:ARG:HA	2:B:151:GLN:HA	2.01	0.42
2:D:124:PRO:HB2	2:D:128:ALA:HB2	2.01	0.42
1:A:128:LEU:HB3	1:A:142:GLU:OE1	2.18	0.42
1:A:132:ASP:OD1	2:D:90:LYS:NZ	2.45	0.42
2:D:166[A]:ASN:ND2	10:D:388:HOH:O	2.54	0.41
4:D:2:YET:C2	4:D:2:YET:N29	2.83	0.41
2:B:36:GLU:H	2:B:36:GLU:CD	2.29	0.40
2:B:24:ASP:OD1	2:B:117:ASN:ND2	2.45	0.40
2:D:245:ARG:HB3	10:D:544:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	80/94 (85%)	78 (98%)	2 (2%)	0	100	100
1	C	81/94 (86%)	77 (95%)	4 (5%)	0	100	100
2	B	238/238 (100%)	231 (97%)	7 (3%)	0	100	100
2	D	236/238 (99%)	229 (97%)	7 (3%)	0	100	100
All	All	635/664 (96%)	615 (97%)	20 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	70/80 (88%)	69 (99%)	1 (1%)	59	75
1	C	64/80 (80%)	64 (100%)	0	100	100
2	B	195/202 (96%)	189 (97%)	6 (3%)	35	48
2	D	199/202 (98%)	195 (98%)	4 (2%)	48	64
All	All	528/564 (94%)	517 (98%)	11 (2%)	47	63

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	ASP
2	B	88	VAL
2	B	97	GLU
2	B	133	GLN
2	B	158	LEU
2	B	210	THR
2	B	247	LEU
2	D	24	ASP
2	D	76	GLU
2	D	151	GLN
2	D	158	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	96	GLN
1	A	120	ASN
1	A	145	ASN
1	A	175	GLN
2	B	30	GLN
2	B	166	ASN
1	C	141	HIS
1	C	175	GLN
2	D	38	ASN
2	D	151	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 23 ligands modelled in this entry, 4 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	GOL	D	5	-	5,5,5	0.32	0	5,5,5	0.46	0
3	GOL	D	6	-	5,5,5	0.36	0	5,5,5	0.66	0
8	ACT	D	12	-	3,3,3	0.81	0	3,3,3	0.90	0
3	GOL	B	8	-	5,5,5	0.37	0	5,5,5	0.24	0
7	MES	D	10	-	12,12,12	2.07	1 (8%)	15,16,16	2.33	5 (33%)
8	ACT	B	11	-	3,3,3	0.86	0	3,3,3	1.18	0
3	GOL	A	21	-	5,5,5	0.86	0	5,5,5	2.17	2 (40%)
4	YET	B	1	-	34,35,35	1.39	2 (5%)	43,48,48	2.69	10 (23%)
7	MES	B	9	-	12,12,12	2.07	1 (8%)	15,16,16	2.52	4 (26%)
8	ACT	B	250	-	3,3,3	0.84	0	3,3,3	0.80	0
8	ACT	D	251	-	3,3,3	0.83	0	3,3,3	0.74	0
8	ACT	B	14	-	3,3,3	0.80	0	3,3,3	0.92	0
9	DMS	B	251	-	3,3,3	2.77	1 (33%)	3,3,3	0.62	0
8	ACT	B	252	-	3,3,3	0.81	0	3,3,3	0.87	0
8	ACT	D	13	-	3,3,3	0.83	0	3,3,3	0.87	0
9	DMS	D	250	-	3,3,3	2.74	1 (33%)	3,3,3	0.59	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	YET	D	2	-	34,35,35	1.51	2 (5%)	43,48,48	2.66	11 (25%)
8	ACT	D	15	-	3,3,3	0.83	0	3,3,3	0.87	0
3	GOL	A	7	-	5,5,5	0.35	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	5	-	-	2/4/4/4	-
3	GOL	D	6	-	-	0/4/4/4	-
7	MES	B	9	-	-	1/6/14/14	0/1/1/1
7	MES	D	10	-	-	1/6/14/14	0/1/1/1
3	GOL	A	21	-	-	2/4/4/4	-
4	YET	D	2	-	-	0/16/41/41	0/4/4/4
4	YET	B	1	-	-	0/16/41/41	0/4/4/4
3	GOL	A	7	-	-	0/4/4/4	-
3	GOL	B	8	-	-	0/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	2	YET	C5-C9	7.06	1.41	1.33
7	B	9	MES	C8-S	-6.86	1.68	1.77
7	D	10	MES	C8-S	-6.86	1.68	1.77
4	B	1	YET	C5-C9	6.33	1.40	1.33
9	B	251	DMS	O-S	4.70	1.81	1.50
9	D	250	DMS	O-S	4.62	1.80	1.50
4	D	2	YET	C12-N25	-3.50	1.33	1.38
4	B	1	YET	C12-N25	-2.67	1.34	1.38

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1	YET	C22-C9-C5	-11.11	124.30	134.08
4	D	2	YET	C22-C9-C5	-10.44	124.89	134.08
7	B	9	MES	C5-N4-C3	7.64	125.31	108.84
4	D	2	YET	C6-C5-C9	-7.15	101.00	107.14
4	B	1	YET	O32-C9-C22	6.87	120.70	114.69
4	B	1	YET	C6-C5-C9	-6.80	101.29	107.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	D	10	MES	C5-N4-C3	6.43	122.69	108.84
4	D	2	YET	O32-C9-C22	6.22	120.14	114.69
4	B	1	YET	O32-C9-C5	5.05	115.00	111.20
4	D	2	YET	O32-C9-C5	5.01	114.97	111.20
4	D	2	YET	C8-C6-C5	4.20	109.96	105.64
4	B	1	YET	C8-C6-C5	3.82	109.57	105.64
7	D	10	MES	C7-N4-C3	3.36	120.19	111.24
3	A	21	GOL	O1-C1-C2	3.05	124.09	110.38
4	B	1	YET	C6-C4-C7	-2.99	118.61	121.43
4	D	2	YET	C21-N29-C12	2.94	127.84	122.13
3	A	21	GOL	O3-C3-C2	2.89	123.39	110.38
7	B	9	MES	O1S-S-C8	2.89	111.09	106.73
4	B	1	YET	C18-N26-C10	2.89	125.78	122.42
7	B	9	MES	O3S-S-C8	2.86	111.59	106.00
7	D	10	MES	O3S-S-C8	2.79	111.46	106.00
7	B	9	MES	C7-N4-C3	2.65	118.30	111.24
4	D	2	YET	C6-C4-C7	-2.63	118.95	121.43
4	D	2	YET	C17-C21-C10	-2.51	108.45	111.65
4	B	1	YET	C14-C18-N26	-2.47	107.69	113.91
4	B	1	YET	C8-O32-C9	-2.47	104.48	105.72
4	B	1	YET	C19-N27-C20	2.35	114.48	111.37
4	D	2	YET	C14-C18-N26	-2.24	108.27	113.91
7	D	10	MES	C7-N4-C5	2.20	117.11	111.24
4	D	2	YET	C18-N26-C10	2.17	124.95	122.42
7	D	10	MES	O1S-S-C8	2.15	109.97	106.73
4	D	2	YET	C8-O32-C9	-2.05	104.69	105.72

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	21	GOL	O1-C1-C2-C3
3	D	5	GOL	O1-C1-C2-C3
7	D	10	MES	C8-C7-N4-C5
3	A	21	GOL	O1-C1-C2-O2
3	D	5	GOL	O1-C1-C2-O2
7	B	9	MES	C8-C7-N4-C5


There are no ring outliers.

6 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	21	GOL	4	0
4	B	1	YET	2	0
7	B	9	MES	1	0
9	B	251	DMS	5	0
9	D	250	DMS	2	0
4	D	2	YET	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	2
2	B	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	145:HIS	C	147:GLU	N	2.42
1	D	218:GLY	C	220:CYS	N	1.97
1	B	145:HIS	C	147:GLU	N	1.92
1	B	218:GLY	C	220:CYS	N	1.90

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	84/94 (89%)	0.41	7 (8%) 17 15	21, 38, 63, 68	1 (1%)
1	C	87/94 (92%)	0.68	12 (13%) 6 5	21, 40, 81, 83	0
2	B	238/238 (100%)	-0.11	2 (0%) 82 80	16, 29, 48, 56	2 (0%)
2	D	237/238 (99%)	-0.11	4 (1%) 69 66	15, 28, 48, 59	5 (2%)
All	All	646/664 (97%)	0.06	25 (3%) 43 40	15, 31, 59, 83	8 (1%)

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	90	CYS	5.2
1	C	89	GLN	4.1
1	C	90	CYS	3.9
1	C	108	TYR	3.5
1	C	101	CYS	3.4
1	C	104	GLY	3.2
1	C	91	GLU	3.1
1	C	111	THR	3.0
2	D	152	SER	2.7
1	C	95	CYS	2.7
2	D	244	THR	2.6
2	D	150	ARG	2.6
1	A	108	TYR	2.5
1	A	109	THR	2.4
1	C	99	GLY	2.4
2	D	142	GLY	2.4
1	A	102	LYS	2.3
1	A	101	CYS	2.3
1	A	103	ASP	2.2
1	C	109	THR	2.2
2	B	74	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	93	SER	2.1
1	C	113	LEU	2.1
1	A	124	PHE	2.0
2	B	148	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	ACT	D	251	4/4	0.68	0.16	54,54,54,54	0
8	ACT	D	13	4/4	0.72	0.19	51,51,52,52	0
3	GOL	A	21	6/6	0.76	0.24	29,34,37,38	0
8	ACT	B	250	4/4	0.78	0.18	48,49,49,49	0
8	ACT	D	12	4/4	0.78	0.15	53,53,53,53	0
9	DMS	D	250	4/4	0.79	0.22	86,87,87,87	0
8	ACT	B	11	4/4	0.81	0.14	43,43,43,43	0
8	ACT	D	15	4/4	0.83	0.17	52,52,53,53	0
5	CA	B	249	1/1	0.85	0.10	86,86,86,86	0
8	ACT	B	252	4/4	0.87	0.16	61,61,61,61	0
8	ACT	B	14	4/4	0.87	0.13	56,56,56,56	0
3	GOL	D	5	6/6	0.87	0.11	48,49,49,49	0
9	DMS	B	251	4/4	0.88	0.16	67,68,68,68	0
3	GOL	B	8	6/6	0.89	0.11	40,43,44,45	0
6	NA	D	4	1/1	0.90	0.06	49,49,49,49	0
7	MES	B	9	12/12	0.93	0.09	34,37,43,43	0
6	NA	B	3	1/1	0.94	0.08	35,35,35,35	0
4	YET	B	1	32/32	0.95	0.06	24,29,31,31	0
7	MES	D	10	12/12	0.95	0.08	32,34,40,40	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	D	6	6/6	0.95	0.08	22,26,27,28	0
4	YET	D	2	32/32	0.96	0.06	24,26,29,33	0
5	CA	D	249	1/1	0.96	0.05	71,71,71,71	0
3	GOL	A	7	6/6	0.97	0.07	27,28,30,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around YET B 1:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.