



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 02:24 PM UTC

PDB ID : 4HPQ / pdb_00004hpq
Title : Crystal Structure of the Atg17-Atg31-Atg29 Complex
Authors : Stanley, R.E.; Ragusa, M.J.; Hurley, J.H.
Deposited on : 2012-10-24
Resolution : 3.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	69	22% 65% 28% 6%
1	D	69	25% 62% 28% 9%
2	B	159	14% 39% 25% 9% 28%
2	E	159	14% 33% 30% 9% 28%
3	C	413	21% 35% 44% 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	413	21% 35% 45% 15% ••

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9220 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Atg29.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	69	Total	C	N	O	S	0	0	0
			429	266	82	80	1			
1	D	69	Total	C	N	O	S	0	0	0
			429	266	82	80	1			

- Molecule 2 is a protein called Atg31.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	115	Total	C	N	O	S	0	0	0
			932	587	156	185	4			
2	E	115	Total	C	N	O	S	0	0	0
			932	587	156	185	4			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	87	MET	LEU	engineered mutation	UNP C5DEB9
B	110	MET	LEU	engineered mutation	UNP C5DEB9
B	146	ALA	-	expression tag	UNP C5DEB9
B	147	GLY	-	expression tag	UNP C5DEB9
B	148	GLN	-	expression tag	UNP C5DEB9
B	149	PHE	-	expression tag	UNP C5DEB9
B	150	TYR	-	expression tag	UNP C5DEB9
B	151	LEU	-	expression tag	UNP C5DEB9
B	152	ASN	-	expression tag	UNP C5DEB9
B	153	ALA	-	expression tag	UNP C5DEB9
B	154	HIS	-	expression tag	UNP C5DEB9
B	155	HIS	-	expression tag	UNP C5DEB9
B	156	HIS	-	expression tag	UNP C5DEB9
B	157	HIS	-	expression tag	UNP C5DEB9
B	158	HIS	-	expression tag	UNP C5DEB9
B	159	HIS	-	expression tag	UNP C5DEB9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	87	MET	LEU	engineered mutation	UNP C5DEB9
E	110	MET	LEU	engineered mutation	UNP C5DEB9
E	146	ALA	-	expression tag	UNP C5DEB9
E	147	GLY	-	expression tag	UNP C5DEB9
E	148	GLN	-	expression tag	UNP C5DEB9
E	149	PHE	-	expression tag	UNP C5DEB9
E	150	TYR	-	expression tag	UNP C5DEB9
E	151	LEU	-	expression tag	UNP C5DEB9
E	152	ASN	-	expression tag	UNP C5DEB9
E	153	ALA	-	expression tag	UNP C5DEB9
E	154	HIS	-	expression tag	UNP C5DEB9
E	155	HIS	-	expression tag	UNP C5DEB9
E	156	HIS	-	expression tag	UNP C5DEB9
E	157	HIS	-	expression tag	UNP C5DEB9
E	158	HIS	-	expression tag	UNP C5DEB9
E	159	HIS	-	expression tag	UNP C5DEB9


- Molecule 3 is a protein called Atg17.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	396	Total	C	N	O	S	0	0	0
			3249	2036	565	635	13			
3	F	396	Total	C	N	O	S	0	0	0
			3249	2036	565	635	13			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Atg29




• Molecule 1: Atg29




• Molecule 2: Atg31




• Molecule 2: Atg31







• Molecule 3: Atg17



• Molecule 3: Atg17





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	144.37Å 64.20Å 184.21Å 90.00° 110.79° 90.00°	Depositor
Resolution (Å)	46.56 – 3.06 46.56 – 3.06	Depositor EDS
% Data completeness (in resolution range)	75.4 (46.56-3.06) 75.8 (46.56-3.06)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.37 (at 3.06Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.303 , 0.336 0.302 , 0.334	Depositor DCC
R_{free} test set	3071 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	64.1	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.003 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	9220	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/288	0.94	0/391
1	D	0.77	0/288	1.00	1/391 (0.3%)
2	B	0.71	0/944	0.99	2/1276 (0.2%)
2	E	0.96	0/944	1.10	3/1276 (0.2%)
3	C	0.73	0/3292	0.97	5/4426 (0.1%)
3	F	0.93	0/3292	1.03	4/4426 (0.1%)
All	All	0.83	0/9048	1.01	15/12186 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	E	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	150	ILE	CB-CA-C	-6.28	103.84	111.88
2	B	112	ARG	N-CA-CB	-6.24	107.06	114.17
3	C	397	ILE	CB-CA-C	-6.03	104.25	111.97
3	F	219	LYS	N-CA-C	-5.80	104.87	111.07
3	F	397	ILE	CB-CA-C	-5.62	104.78	111.97

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	111	SER	Peptide
2	E	111	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	429	0	278	38	0
1	D	429	0	280	42	0
2	B	932	0	911	62	0
2	E	932	0	911	65	0
3	C	3249	0	3253	248	0
3	F	3249	0	3253	238	0
All	All	9220	0	8886	595	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 595 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:109:ILE:HG21	3:C:121:LEU:HD22	1.17	1.14
3:F:109:ILE:HG21	3:F:121:LEU:HD22	1.18	1.13
3:C:67:LEU:HD11	3:C:157:MET:HE2	1.51	0.91
3:F:67:LEU:HD11	3:F:157:MET:HE2	1.50	0.91
1:D:1:MET:HB3	2:E:44:MET:SD	2.11	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	38/69 (55%)	31 (82%)	6 (16%)	1 (3%)	4	17
1	D	38/69 (55%)	30 (79%)	7 (18%)	1 (3%)	4	17
2	B	109/159 (69%)	94 (86%)	10 (9%)	5 (5%)	2	9
2	E	109/159 (69%)	94 (86%)	9 (8%)	6 (6%)	1	7
3	C	392/413 (95%)	339 (86%)	49 (12%)	4 (1%)	12	37
3	F	392/413 (95%)	340 (87%)	50 (13%)	2 (0%)	24	52
All	All	1078/1282 (84%)	928 (86%)	131 (12%)	19 (2%)	6	24

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	111	ASN
3	F	111	ASN
2	B	77	ASP
2	B	113	PHE
3	C	113	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	23/36 (64%)	17 (74%)	6 (26%)	0	1
1	D	23/36 (64%)	17 (74%)	6 (26%)	0	1
2	B	107/145 (74%)	86 (80%)	21 (20%)	1	5
2	E	107/145 (74%)	84 (78%)	23 (22%)	1	4
3	C	367/383 (96%)	266 (72%)	101 (28%)	0	1
3	F	367/383 (96%)	262 (71%)	105 (29%)	0	1
All	All	994/1128 (88%)	732 (74%)	262 (26%)	0	1

5 of 262 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	F	306	ILE
3	F	327	LEU
3	F	403	LEU
3	C	306	ILE
3	C	292	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	91	GLN
3	F	149	ASN
2	E	118	GLN
3	F	82	GLN
3	F	216	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1
1	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	40:VAL	C	51:UNK	N	15.44
1	D	40:VAL	C	51:UNK	N	14.58

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	40/69 (57%)	1.95	15 (37%) 1 0	59, 94, 148, 158	0
1	D	40/69 (57%)	2.19	17 (42%) 0 0	53, 69, 117, 161	0
2	B	115/159 (72%)	1.24	23 (20%) 3 1	45, 78, 111, 126	0
2	E	115/159 (72%)	1.02	23 (20%) 3 1	23, 50, 94, 115	0
3	C	396/413 (95%)	1.29	86 (21%) 2 1	31, 72, 148, 189	0
3	F	396/413 (95%)	1.16	88 (22%) 2 1	22, 52, 126, 165	0
All	All	1102/1282 (85%)	1.27	252 (22%) 2 1	22, 67, 135, 189	0

The worst 5 of 252 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	233	ALA	8.9
3	F	110	THR	8.3
1	D	16	ALA	7.1
1	D	15	ARG	6.8
3	C	237	PHE	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.