



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 08:13 PM UTC

PDB ID : 6HP7 / pdb_00006hp7
Title : ARBITRIUM PEPTIDE RECEPTOR FROM SPBETA PHAGE in complex with 43 mer DNA
Authors : Marina, A.; Gallego del Sol, F.
Deposited on : 2018-09-19
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	0% 87% 10% ..
1	B	386	2% 85% 11% .
2	C	43	51% 40% 9%
3	D	43	2% 49% 42% 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8420 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SPBc2 prophage-derived uncharacterized protein YopK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	386	Total	C	N	O	S	0	1	0
			3189	2035	528	605	21			
1	B	386	Total	C	N	O	S	0	0	0
			3183	2031	528	603	21			

- Molecule 2 is a DNA chain called DNA (43-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	43	Total	C	N	O	P	0	0	0
			884	425	157	259	43			

- Molecule 3 is a DNA chain called DNA (43-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	43	Total	C	N	O	P	0	0	0
			879	423	156	257	43			

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	C	1	Total	O	P	0	0
			5	4	1		
4	D	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	116	Total	O	0	0
			116	116		
5	B	119	Total	O	0	0
			119	119		
5	C	13	Total	O	0	0
			13	13		
5	D	7	Total	O	0	0
			7	7		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: SPBc2 prophage-derived uncharacterized protein YopK




- Molecule 1: SPBc2 prophage-derived uncharacterized protein YopK



- Molecule 2: DNA (43-MER)



- Molecule 3: DNA (43-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	159.16Å 159.16Å 245.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	133.61 – 2.20 133.61 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (133.61-2.20) 100.0 (133.61-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.216 , 0.246 (Not available) , 0.234	Depositor DCC
R_{free} test set	4007 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	52.5	Xtriage
Anisotropy	0.406	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 30.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8420	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.93	1/3249 (0.0%)	1.01	4/4369 (0.1%)
1	B	0.92	0/3240	1.02	4/4357 (0.1%)
2	C	0.47	0/991	1.20	10/1528 (0.7%)
3	D	0.42	0/985	1.26	11/1517 (0.7%)
All	All	0.84	1/8465 (0.0%)	1.07	29/11771 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	19	MET	SD-CE	-6.41	1.63	1.79

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	16	DT	C4'-C3'-O3'	11.99	127.98	110.00
3	D	11	DA	C4'-C3'-O3'	11.52	127.27	110.00
2	C	1	DT	C2'-C3'-O3'	11.07	128.10	111.50
3	D	31	DA	C4'-C3'-O3'	10.27	125.41	110.00
2	C	18	DT	C2'-C3'-O3'	8.66	124.48	111.50
3	D	17	DT	C4'-C3'-O3'	8.22	122.34	110.00
1	B	82	ARG	NE-CZ-NH2	-7.14	112.77	119.20
3	D	11	DA	C2'-C3'-O3'	-6.76	101.35	111.50
3	D	2	DC	C4'-C3'-O3'	6.53	119.80	110.00
2	C	15	DT	C4'-C3'-O3'	-6.52	100.22	110.00
1	A	5	ARG	NE-CZ-NH2	6.44	125.00	119.20
3	D	19	DG	C4'-C3'-O3'	6.44	119.65	110.00
1	A	5	ARG	NE-CZ-NH1	-6.43	115.07	121.50
2	C	28	DA	C4'-C3'-O3'	6.43	119.64	110.00
3	D	9	DT	C4'-C3'-O3'	6.28	119.42	110.00
3	D	39	DA	C4'-C3'-O3'	-6.21	100.68	110.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	GLN	N-CA-C	6.09	119.81	112.38
2	C	13	DT	C2'-C3'-O3'	6.01	120.52	111.50
2	C	36	DG	C4'-C3'-O3'	5.98	118.97	110.00
3	D	39	DA	C2'-C3'-O3'	5.56	119.84	111.50
1	B	306	ILE	N-CA-CB	5.55	118.08	110.54
2	C	12	DA	C2'-C3'-O3'	5.37	119.56	111.50
1	A	206	ASN	N-CA-C	5.34	117.10	111.28
1	B	82	ARG	NE-CZ-NH1	5.29	126.79	121.50
3	D	41	DC	C2'-C3'-O3'	5.26	119.39	111.50
2	C	33	DT	C2'-C3'-O3'	-5.24	103.64	111.50
3	D	30	DC	C2'-C3'-O3'	5.21	119.32	111.50
2	C	29	DT	C2'-C3'-O3'	5.17	119.26	111.50
1	B	156	LEU	CA-CB-CG	5.08	134.07	116.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3189	0	3185	27	0
1	B	3183	0	3179	24	0
2	C	884	0	490	13	0
3	D	879	0	489	11	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
4	C	5	0	0	0	0
4	D	5	0	0	0	0
5	A	116	0	0	0	0
5	B	119	0	0	1	0
5	C	13	0	0	0	0
5	D	7	0	0	0	0
All	All	8420	0	7343	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:136:VAL:HG22	1:B:136:VAL:HG22	1.29	1.13
1:A:19:MET:HE2	1:A:22:TRP:HE3	1.27	0.95
2:C:25:DC:N3	3:D:19:DG:N2	2.14	0.95
1:A:19:MET:HE2	1:A:22:TRP:CE3	2.09	0.87
1:B:30:ASN:C	1:B:30:ASN:HD22	1.84	0.86
1:A:19:MET:CE	1:A:22:TRP:CE3	2.63	0.82
2:C:19:DT:H2'	2:C:20:DA:C8	2.21	0.76
2:C:13:DT:H2''	2:C:14:DG:H5''	1.69	0.73
1:B:267:ASN:O	5:B:501:HOH:O	2.10	0.68
3:D:19:DG:H2'	3:D:20:DG:C8	2.29	0.68
1:A:19:MET:CE	1:A:22:TRP:CZ3	2.79	0.66
1:B:189:ARG:HH11	1:B:190:ASN:HD21	1.44	0.66
2:C:18:DT:H2''	2:C:19:DT:O5'	1.98	0.63
1:A:372:GLN:NE2	1:A:378:GLN:OE1	2.32	0.61
1:A:162:LEU:HA	1:A:170:MET:CE	2.31	0.60
1:A:162:LEU:HA	1:A:170:MET:HE1	1.83	0.59
1:A:167:PHE:CZ	1:A:206:ASN:ND2	2.71	0.59
1:A:162:LEU:N	1:A:170:MET:HE2	2.21	0.56
1:B:30:ASN:ND2	1:B:32:ASN:H	2.04	0.56
1:A:162:LEU:CA	1:A:170:MET:CE	2.84	0.55
1:B:241:LEU:HD13	1:B:248:LEU:HB3	1.88	0.55
1:A:158:LEU:HD13	1:A:173:LEU:HD23	1.89	0.54
1:B:1:MET:HE1	1:B:9:LYS:HE3	1.88	0.54
1:B:158:LEU:HD13	1:B:173:LEU:HD23	1.89	0.53
1:A:19:MET:HE1	1:A:22:TRP:CZ3	2.43	0.53
3:D:11:DA:H4'	3:D:12:DA:OP1	2.08	0.53
2:C:6:DC:H2''	2:C:7:DA:C8	2.43	0.53
1:B:172:GLN:NE2	1:B:176:GLN:HE21	2.08	0.52
1:A:19:MET:HE1	1:A:22:TRP:HZ3	1.75	0.52
3:D:7:DA:H2''	3:D:8:DC:H5''	1.91	0.51
1:B:30:ASN:C	1:B:30:ASN:ND2	2.59	0.51
3:D:17:DT:H2''	3:D:18:DA:C8	2.47	0.50
2:C:13:DT:C2'	2:C:14:DG:H5''	2.38	0.50
1:A:262:GLN:NE2	1:A:262:GLN:H	2.10	0.50
1:A:153:SER:HA	1:A:156:LEU:HD13	1.94	0.49
1:B:369:LEU:HD13	1:B:373:LYS:HE2	1.94	0.49
2:C:18:DT:H2''	2:C:19:DT:C5'	2.43	0.49
1:A:131:THR:OG1	1:A:134[A]:GLU:HG3	2.14	0.48
1:B:12:LEU:HD21	1:B:19:MET:HG2	1.94	0.48
1:B:172:GLN:NE2	1:B:176:GLN:NE2	2.62	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:GLN:NE2	1:B:262:GLN:H	2.12	0.48
1:A:162:LEU:N	1:A:170:MET:CE	2.77	0.47
1:B:153:SER:HA	1:B:156:LEU:HD13	1.97	0.47
1:B:260:SER:HB2	1:B:266:TYR:HB2	1.98	0.46
1:A:312:SER:O	1:A:316:GLU:HG2	2.16	0.45
1:B:32:ASN:ND2	2:C:36:DG:O6	2.47	0.45
1:B:266:TYR:HA	1:B:269:ILE:HD12	1.97	0.45
1:A:43:LYS:NZ	3:D:34:DT:OP2	2.50	0.45
1:A:52:VAL:HG22	1:A:67:MET:HE3	2.00	0.44
2:C:1:DT:H2'	2:C:2:DT:C6	2.53	0.44
1:B:189:ARG:HH11	1:B:190:ASN:ND2	2.12	0.44
3:D:42:DA:H2''	3:D:43:DA:OP2	2.18	0.44
2:C:9:DT:H2'	2:C:10:DA:C8	2.53	0.43
1:A:52:VAL:HA	1:A:67:MET:CE	2.49	0.43
1:A:172:GLN:OE1	1:A:176:GLN:NE2	2.52	0.43
1:B:71:CYS:HA	1:B:74:LEU:HD22	2.01	0.43
2:C:23:DA:N1	3:D:21:DT:O4	2.52	0.43
1:A:242:LEU:HD12	1:A:277:LEU:HD13	1.99	0.42
1:B:333:MET:HG3	1:B:337:LEU:HD22	2.01	0.42
1:B:16:ASN:C	1:B:16:ASN:HD22	2.26	0.42
2:C:30:DA:C2	3:D:15:DA:C2	3.08	0.41
1:A:262:GLN:H	1:A:262:GLN:HE21	1.67	0.41
1:A:5:ARG:HD3	1:A:37:PHE:O	2.20	0.41
1:B:312:SER:O	1:B:315:LYS:HG3	2.21	0.41
1:A:16:ASN:HD22	1:A:16:ASN:C	2.29	0.41
3:D:9:DT:H2'	3:D:10:DT:H71	2.03	0.40
3:D:3:DC:H2''	3:D:4:DA:C8	2.57	0.40
2:C:18:DT:H2''	2:C:19:DT:O4'	2.21	0.40
1:B:242:LEU:HD13	1:B:277:LEU:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	385/386 (100%)	380 (99%)	5 (1%)	0	100	100
1	B	384/386 (100%)	378 (98%)	6 (2%)	0	100	100
All	All	769/772 (100%)	758 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	361/360 (100%)	330 (91%)	31 (9%)	10	10
1	B	360/360 (100%)	322 (89%)	38 (11%)	6	6
All	All	721/720 (100%)	652 (90%)	69 (10%)	8	8

All (69) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	GLU
1	A	16	ASN
1	A	19	MET
1	A	21	LYS
1	A	74	LEU
1	A	76	VAL
1	A	77	LYS
1	A	105	ILE
1	A	136	VAL
1	A	141	LYS
1	A	156	LEU
1	A	157	LEU
1	A	158	LEU
1	A	168	SER
1	A	172	GLN
1	A	198	VAL
1	A	206	ASN
1	A	218	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	262	GLN
1	A	298	LEU
1	A	312	SER
1	A	320	LYS
1	A	323	LEU
1	A	324	LEU
1	A	338	LYS
1	A	341	LEU
1	A	342	GLU
1	A	363	LEU
1	A	368	LEU
1	A	380	LEU
1	A	386	LEU
1	B	1	MET
1	B	13	GLU
1	B	16	ASN
1	B	21	LYS
1	B	30	ASN
1	B	34	LEU
1	B	72	LEU
1	B	74	LEU
1	B	136	VAL
1	B	156	LEU
1	B	157	LEU
1	B	158	LEU
1	B	168	SER
1	B	189	ARG
1	B	198	VAL
1	B	211	GLU
1	B	242	LEU
1	B	255	LYS
1	B	262	GLN
1	B	293	ASP
1	B	296	MET
1	B	298	LEU
1	B	306	ILE
1	B	310	GLU
1	B	312	SER
1	B	315	LYS
1	B	316	GLU
1	B	323	LEU
1	B	331	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	337	LEU
1	B	342	GLU
1	B	356	LYS
1	B	366	LEU
1	B	368	LEU
1	B	369	LEU
1	B	378	GLN
1	B	380	LEU
1	B	386	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	ASN
1	A	32	ASN
1	A	39	ASN
1	A	206	ASN
1	A	250	GLN
1	A	262	GLN
1	A	265	ASN
1	A	359	ASN
1	B	16	ASN
1	B	30	ASN
1	B	39	ASN
1	B	143	ASN
1	B	172	GLN
1	B	190	ASN
1	B	262	GLN
1	B	265	ASN
1	B	299	GLN
1	B	301	GLN
1	B	311	ASN
1	B	372	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PO4	B	401	-	4,4,4	0.82	0	6,6,6	0.60	0
4	PO4	A	402	-	4,4,4	0.75	0	6,6,6	0.72	0
4	PO4	B	402	-	4,4,4	0.78	0	6,6,6	0.39	0
4	PO4	A	401	-	4,4,4	0.89	0	6,6,6	0.53	0
4	PO4	D	101	-	4,4,4	0.93	0	6,6,6	0.50	0
4	PO4	C	101	-	4,4,4	0.90	0	6,6,6	0.46	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	386/386 (100%)	0.11	2 (0%) 87 85	28, 49, 73, 93	1 (0%)
1	B	386/386 (100%)	0.21	9 (2%) 61 58	37, 51, 81, 103	0
2	C	43/43 (100%)	0.29	0 100 100	39, 61, 86, 120	0
3	D	43/43 (100%)	0.35	1 (2%) 61 58	41, 61, 83, 152	0
All	All	858/858 (100%)	0.18	12 (1%) 73 71	28, 51, 80, 152	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	276	PHE	4.9
1	A	276	PHE	4.6
3	D	1	DG	3.7
1	B	1	MET	3.4
1	B	325	VAL	3.0
1	B	265	ASN	2.5
1	B	323	LEU	2.5
1	B	295	ILE	2.3
1	B	320	LYS	2.1
1	B	306	ILE	2.1
1	A	325	VAL	2.1
1	B	331	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PO4	C	101	5/5	0.77	0.16	119,122,130,137	0
4	PO4	B	402	5/5	0.78	0.14	103,105,110,123	0
4	PO4	D	101	5/5	0.81	0.16	124,126,129,135	0
4	PO4	A	402	5/5	0.86	0.13	93,100,103,116	0
4	PO4	B	401	5/5	0.88	0.15	89,111,117,119	0
4	PO4	A	401	5/5	0.88	0.12	108,109,117,124	0

6.5 Other polymers [i](#)

There are no such residues in this entry.