



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:42 AM UTC

PDB ID : 6HQD / pdb_00006hqd
Title : Cytochrome P450-153 from Pseudomonas sp. 19-rlim
Authors : Fiorentini, F.; Mattevi, A.
Deposited on : 2018-09-24
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	
1	B	420	
1	C	420	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SRT	A	502	-	X	-	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 10426 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	414	Total	C	N	O	S	0	0	0
			3307	2099	576	612	20			
1	B	412	Total	C	N	O	S	0	0	0
			3300	2093	577	610	20			
1	C	415	Total	C	N	O	S	0	0	0
			3322	2107	579	616	20			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is S,R MESO-TARTARIC ACID (CCD ID: SRT) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	4	2		
4	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	4	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		
5	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	83	Total	O	0	0
			83	83		
6	B	123	Total	O	0	0
			123	123		
6	C	115	Total	O	0	0
			115	115		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cytochrome P450



• Molecule 1: Cytochrome P450



• Molecule 1: Cytochrome P450



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.31Å 96.64Å 87.78Å 90.00° 119.18° 90.00°	Depositor
Resolution (Å)	44.00 – 1.80 44.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (44.00-1.80) 99.2 (44.00-1.80)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.67 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.170 , 0.200 0.181 , 0.211	Depositor DCC
R_{free} test set	5690 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.026 for -h-l,k,h 0.026 for l,k,-h-l 0.027 for h,-k,-h-l 0.029 for -h-l,-k,l 0.028 for l,-k,h	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10426	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, GOL, PGE, SRT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	2/3388 (0.1%)	1.04	3/4603 (0.1%)
1	B	1.02	2/3381 (0.1%)	1.03	5/4592 (0.1%)
1	C	1.01	0/3404	1.04	4/4624 (0.1%)
All	All	1.01	4/10173 (0.0%)	1.04	12/13819 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	2
1	C	0	8
All	All	0	17

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	85	SER	N-CA	5.31	1.53	1.46
1	B	35	HIS	CE1-NE2	5.27	1.37	1.32
1	A	27	GLU	N-CA	5.16	1.52	1.46
1	B	21	ILE	N-CA	5.06	1.51	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	71	ASP	CA-CB-CG	6.55	119.15	112.60
1	A	71	ASP	CA-CB-CG	6.30	118.91	112.60
1	C	105	PHE	CA-CB-CG	5.97	119.77	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	109	ASP	CB-CA-C	5.87	117.91	109.11
1	C	71	ASP	CA-CB-CG	5.80	118.40	112.60
1	B	341	ASP	CA-CB-CG	5.50	118.09	112.60
1	B	109	ASP	CA-CB-CG	5.32	117.92	112.60
1	A	105	PHE	CA-CB-CG	5.29	119.09	113.80
1	C	346	ARG	CB-CA-C	-5.28	107.45	111.20
1	B	237	HIS	CA-CB-CG	-5.17	108.63	113.80
1	A	230	ASP	CA-CB-CG	5.11	117.71	112.60
1	C	230	ASP	CA-CB-CG	5.08	117.67	112.60

There are no chirality outliers.

All (17) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	179	ARG	Sidechain
1	A	222	ARG	Sidechain
1	A	265	ARG	Sidechain
1	A	320	ARG	Sidechain
1	A	339	ARG	Sidechain
1	A	390	ARG	Sidechain
1	A	43	ARG	Sidechain
1	B	310	ARG	Sidechain
1	B	369	ARG	Sidechain
1	C	142	ARG	Sidechain
1	C	217	ARG	Sidechain
1	C	289	ARG	Sidechain
1	C	310	ARG	Sidechain
1	C	339	ARG	Sidechain
1	C	346	ARG	Sidechain
1	C	358	ARG	Sidechain
1	C	374	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3307	0	3227	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3300	0	3225	17	0
1	C	3322	0	3244	8	0
2	A	43	0	30	2	0
2	B	43	0	30	3	0
2	C	43	0	30	2	0
3	A	10	0	4	0	0
4	A	6	0	7	1	0
4	B	7	0	9	3	0
4	C	6	0	7	1	0
5	B	6	0	7	3	0
5	C	12	0	16	1	0
6	A	83	0	0	1	0
6	B	123	0	0	8	0
6	C	115	0	0	2	0
All	All	10426	0	9836	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:243:ASP:HB3	6:B:718:HOH:O	1.62	0.99
1:B:228:LYS:HE3	1:C:277:LEU:HD22	1.54	0.90
1:B:309:MET:SD	6:B:631:HOH:O	2.46	0.72
5:B:502:GOL:H11	6:B:634:HOH:O	1.91	0.71
2:B:501:HEM:HBC2	2:B:501:HEM:HMC2	1.72	0.70
1:C:358:ARG:NH2	1:C:377:GLU:OE1	2.27	0.66
1:B:116:HIS:CD2	1:B:241:MET:HE1	2.32	0.65
1:B:226:PRO:HB3	1:C:348:MET:CE	2.30	0.61
1:B:12:ILE:HD12	6:B:611:HOH:O	2.01	0.61
2:A:501:HEM:HMC2	2:A:501:HEM:HBC2	1.86	0.57
1:C:14:ARG:CB	6:C:710:HOH:O	2.51	0.57
1:B:228:LYS:HE3	1:C:277:LEU:CD2	2.32	0.56
4:B:503:PGE:C6	6:B:686:HOH:O	2.55	0.55
2:B:501:HEM:HBC2	2:B:501:HEM:CMC	2.34	0.55
4:B:503:PGE:H62	6:B:686:HOH:O	2.09	0.53
1:B:406:ASN:N	5:B:502:GOL:O1	2.27	0.53
1:C:339:ARG:HA	1:C:347:PRO:HB2	1.90	0.53
1:A:130:LEU:CD2	6:B:612:HOH:O	2.57	0.52
1:B:405:SER:HA	5:B:502:GOL:H12	1.92	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:C:504:PGE:H1	6:C:700:HOH:O	2.10	0.50
1:B:286:ARG:HH11	1:B:286:ARG:HG2	1.76	0.50
1:A:126:PRO:O	1:A:130:LEU:HD23	2.12	0.49
1:B:263:THR:HB	2:B:501:HEM:C3B	2.52	0.45
1:A:351:ILE:HD12	1:A:354:ARG:NH2	2.31	0.45
1:B:186:SER:CB	1:B:261:ASN:HD22	2.29	0.45
4:A:503:PGE:H6	6:A:613:HOH:O	2.17	0.44
1:B:121:SER:HB2	1:B:122:PRO:HD3	1.98	0.44
4:B:503:PGE:H6	6:B:686:HOH:O	2.16	0.44
1:A:263:THR:HB	2:A:501:HEM:C3B	2.53	0.44
1:B:164:THR:HG21	1:B:261:ASN:O	2.18	0.44
1:A:341:ASP:HA	1:A:344:ILE:O	2.19	0.42
1:B:339:ARG:HA	1:B:347:PRO:HB2	2.02	0.42
2:C:501:HEM:HMB2	2:C:501:HEM:HBB2	2.01	0.41
1:B:133:GLU:HG3	1:B:378:LEU:HD11	2.01	0.41
1:A:339:ARG:HA	1:A:347:PRO:HB2	2.03	0.41
1:A:8:ASP:OD2	1:A:11:GLU:HB2	2.21	0.41
1:B:123:ILE:HG23	1:B:124:VAL:HG13	2.02	0.40
1:C:42:ARG:HE	5:C:503:GOL:H32	1.86	0.40
1:C:81:HIS:ND1	1:C:101:TYR:OH	2.39	0.40
2:C:501:HEM:HBC2	2:C:501:HEM:HHD	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	412/420 (98%)	406 (98%)	6 (2%)	0	100	100
1	B	410/420 (98%)	402 (98%)	8 (2%)	0	100	100
1	C	413/420 (98%)	405 (98%)	8 (2%)	0	100	100
All	All	1235/1260 (98%)	1213 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	360/375 (96%)	352 (98%)	8 (2%)	45	34
1	B	360/375 (96%)	352 (98%)	8 (2%)	45	34
1	C	363/375 (97%)	353 (97%)	10 (3%)	38	26
All	All	1083/1125 (96%)	1057 (98%)	26 (2%)	43	31

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	SER
1	A	136	ILE
1	A	138	GLU
1	A	183	THR
1	A	230	ASP
1	A	240	GLU
1	A	309	MET
1	A	416	VAL
1	B	25	LYS
1	B	51	LYS
1	B	53	SER
1	B	85	SER
1	B	87	VAL
1	B	177	ASP
1	B	240	GLU
1	B	317	VAL
1	C	75	LYS
1	C	118	MET
1	C	177	ASP
1	C	230	ASP
1	C	240	GLU
1	C	261	ASN
1	C	287	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	299	VAL
1	C	317	VAL
1	C	416	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	94	ASN
1	A	127	GLN
1	A	287	ASN
1	A	303	GLN
1	A	373	ASN
1	B	74	HIS
1	B	223	GLN
1	B	237	HIS
1	B	261	ASN
1	B	303	GLN
1	C	104	ASN
1	C	237	HIS
1	C	280	GLN
1	C	419	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	B	501	6,1	50,50,50	1.51	10 (20%)	67,82,82	1.75	15 (22%)
2	HEM	C	501	6,1	50,50,50	1.62	9 (18%)	67,82,82	1.63	15 (22%)
5	GOL	C	503	-	5,5,5	0.81	0	5,5,5	0.93	0
4	PGE	B	503	-	6,6,9	0.31	0	5,5,8	0.81	0
3	SRT	A	502	-	9,9,9	0.99	0	12,12,12	3.33	4 (33%)
5	GOL	C	502	-	5,5,5	0.53	0	5,5,5	1.03	0
4	PGE	C	504	-	5,5,9	0.80	0	4,4,8	0.39	0
2	HEM	A	501	6,1	50,50,50	1.59	7 (14%)	67,82,82	1.88	18 (26%)
5	GOL	B	502	-	5,5,5	0.95	0	5,5,5	0.93	0
4	PGE	A	503	-	5,5,9	0.85	0	4,4,8	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	501	6,1	-	1/14/54/54	-
2	HEM	C	501	6,1	-	3/14/54/54	-
5	GOL	C	503	-	-	2/4/4/4	-
4	PGE	B	503	-	-	1/4/4/7	-
3	SRT	A	502	-	-	10/12/12/12	-
5	GOL	C	502	-	-	4/4/4/4	-
4	PGE	C	504	-	-	1/3/3/7	-
2	HEM	A	501	6,1	-	2/14/54/54	-
5	GOL	B	502	-	-	2/4/4/4	-
4	PGE	A	503	-	-	0/3/3/7	-

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	HEM	FE-NB	4.87	2.09	1.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	HEM	FE-NB	4.46	2.08	1.94
2	B	501	HEM	FE-NB	4.21	2.07	1.94
2	A	501	HEM	C1D-C2D	4.15	1.52	1.44
2	C	501	HEM	FE-NC	3.98	2.08	1.95
2	A	501	HEM	FE-NC	3.53	2.06	1.95
2	B	501	HEM	FE-NC	3.49	2.06	1.95
2	A	501	HEM	C4D-C3D	3.31	1.50	1.45
2	C	501	HEM	C4B-NB	-3.19	1.32	1.38
2	C	501	HEM	C1B-NB	-2.87	1.35	1.40
2	C	501	HEM	C4D-ND	-2.86	1.35	1.40
2	B	501	HEM	C1D-C2D	2.80	1.50	1.44
2	A	501	HEM	C1C-C2C	-2.56	1.40	1.45
2	A	501	HEM	C1B-NB	-2.53	1.35	1.40
2	B	501	HEM	C3B-C4B	2.47	1.49	1.44
2	B	501	HEM	O1D-CGD	2.41	1.30	1.22
2	C	501	HEM	C4D-C3D	2.38	1.49	1.45
2	C	501	HEM	C1C-NC	-2.36	1.35	1.39
2	B	501	HEM	C4B-NB	-2.29	1.34	1.38
2	B	501	HEM	C1B-NB	-2.28	1.36	1.40
2	B	501	HEM	FE-NA	2.26	2.02	1.95
2	B	501	HEM	CBD-CGD	2.18	1.55	1.50
2	A	501	HEM	O1A-CGA	2.16	1.29	1.22
2	C	501	HEM	O1A-CGA	2.16	1.29	1.22
2	C	501	HEM	FE-NA	2.04	2.01	1.95
2	B	501	HEM	CHC-C4B	2.03	1.43	1.39

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	SRT	O3-C3-C2	-7.83	94.24	110.17
3	A	502	SRT	C2-C3-C4	6.33	123.86	109.82
2	A	501	HEM	CHC-C4B-NB	5.52	130.37	124.42
2	B	501	HEM	CHC-C4B-NB	5.38	130.21	124.42
2	A	501	HEM	C1B-NB-C4B	5.02	111.15	105.21
2	A	501	HEM	CHD-C4C-NC	4.67	129.54	124.45
2	C	501	HEM	C1B-NB-C4B	4.36	110.37	105.21
2	C	501	HEM	CHC-C4B-NB	4.27	129.02	124.42
2	B	501	HEM	C1B-NB-C4B	4.06	110.01	105.21
2	C	501	HEM	CAC-C3C-C4C	4.04	134.46	124.82
2	B	501	HEM	CHD-C4C-NC	3.71	128.50	124.45
2	A	501	HEM	CAD-C3D-C4D	3.42	130.65	124.70
2	C	501	HEM	CAC-C3C-C2C	-3.30	117.69	128.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	HEM	C3B-C4B-NB	-3.27	107.12	109.47
2	A	501	HEM	C3B-C4B-NB	-3.18	107.19	109.47
2	B	501	HEM	O2A-CGA-CBA	3.12	123.87	114.00
2	A	501	HEM	C3B-C2B-C1B	3.12	108.75	106.41
2	C	501	HEM	CHD-C1D-ND	3.00	127.65	124.42
3	A	502	SRT	O3-C3-C4	-2.97	104.34	110.69
2	B	501	HEM	CHA-C4D-C3D	-2.95	119.78	125.23
2	B	501	HEM	C3D-C4D-ND	2.93	113.39	110.17
2	A	501	HEM	C3D-C4D-ND	2.84	113.28	110.17
2	C	501	HEM	CHA-C4D-ND	2.82	127.85	124.37
2	A	501	HEM	CHB-C1B-NB	2.80	127.83	124.37
2	A	501	HEM	CMD-C2D-C1D	2.79	129.39	125.03
2	C	501	HEM	C3B-C4B-NB	-2.77	107.48	109.47
2	B	501	HEM	O1A-CGA-CBA	-2.76	114.35	123.09
2	A	501	HEM	O2A-CGA-CBA	2.71	122.55	114.00
2	C	501	HEM	CMB-C2B-C1B	2.67	129.20	125.03
2	B	501	HEM	CHD-C1D-C2D	-2.57	120.98	125.03
2	A	501	HEM	CMA-C3A-C4A	-2.56	121.52	125.42
3	A	502	SRT	O41-C4-C3	2.50	120.26	113.31
2	C	501	HEM	CHB-C1B-NB	2.45	127.40	124.37
2	A	501	HEM	O2D-CGD-CBD	2.41	121.61	114.00
2	C	501	HEM	C1A-CHA-C4D	-2.38	120.65	126.25
2	B	501	HEM	C2D-C1D-ND	2.37	112.65	109.90
2	A	501	HEM	CMA-C3A-C2A	2.37	130.65	125.62
2	C	501	HEM	CMC-C2C-C1C	2.34	128.84	124.73
2	C	501	HEM	CHD-C1D-C2D	-2.32	121.36	125.03
2	A	501	HEM	CHA-C4D-C3D	-2.32	120.94	125.23
2	A	501	HEM	C4C-NC-C1C	2.29	109.56	105.82
2	C	501	HEM	CHA-C4D-C3D	-2.27	121.05	125.23
2	A	501	HEM	C2B-C1B-NB	-2.25	107.25	109.84
2	C	501	HEM	C4C-NC-C1C	2.18	109.38	105.82
2	B	501	HEM	CHA-C1A-NA	2.17	127.79	123.86
2	B	501	HEM	O2D-CGD-O1D	-2.12	117.88	123.33
2	B	501	HEM	C4C-NC-C1C	2.11	109.26	105.82
2	C	501	HEM	O2A-CGA-CBA	2.09	120.60	114.00
2	B	501	HEM	O2D-CGD-CBD	2.08	120.57	114.00
2	A	501	HEM	C4A-CHB-C1B	-2.08	121.37	126.25
2	A	501	HEM	C4D-C3D-C2D	-2.04	103.93	106.89
2	B	501	HEM	CAA-C2A-C1A	2.01	128.86	124.94

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502	SRT	C1-C2-C3-O3
5	C	502	GOL	O2-C2-C3-O3
5	C	503	GOL	C1-C2-C3-O3
3	A	502	SRT	O2-C2-C3-O3
3	A	502	SRT	C1-C2-C3-C4
3	A	502	SRT	O1-C1-C2-C3
3	A	502	SRT	O11-C1-C2-C3
3	A	502	SRT	O2-C2-C3-C4
3	A	502	SRT	O1-C1-C2-O2
3	A	502	SRT	O11-C1-C2-O2
5	C	503	GOL	O2-C2-C3-O3
5	B	502	GOL	C1-C2-C3-O3
5	C	502	GOL	O1-C1-C2-C3
5	C	502	GOL	C1-C2-C3-O3
5	C	502	GOL	O1-C1-C2-O2
4	B	503	PGE	O3-C5-C6-O4
2	C	501	HEM	C4C-C3C-CAC-CBC
5	B	502	GOL	O2-C2-C3-O3
4	C	504	PGE	O1-C1-C2-O2
3	A	502	SRT	C2-C3-C4-O41
2	C	501	HEM	CAD-CBD-CGD-O2D
2	B	501	HEM	CAD-CBD-CGD-O2D
2	A	501	HEM	CAD-CBD-CGD-O2D
3	A	502	SRT	O3-C3-C4-O4
2	C	501	HEM	CAD-CBD-CGD-O1D
2	A	501	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


8 monomers are involved in 16 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	HEM	3	0
2	C	501	HEM	2	0
5	C	503	GOL	1	0
4	B	503	PGE	3	0
4	C	504	PGE	1	0
2	A	501	HEM	2	0
5	B	502	GOL	3	0
4	A	503	PGE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	414/420 (98%)	-0.02	4 (0%) 79 80	17, 26, 42, 59	0
1	B	412/420 (98%)	-0.20	6 (1%) 72 72	15, 23, 41, 61	0
1	C	415/420 (98%)	-0.26	2 (0%) 87 88	15, 22, 37, 55	0
All	All	1241/1260 (98%)	-0.16	12 (0%) 79 80	15, 24, 40, 61	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	6	PRO	4.4
1	B	9	ASP	4.3
1	B	10	ALA	4.0
1	C	365	PHE	3.2
1	A	420	ALA	2.7
1	B	224	ALA	2.6
1	A	176	PHE	2.4
1	B	243	ASP	2.3
1	B	420	ALA	2.3
1	B	12	ILE	2.2
1	A	7	ILE	2.1
1	A	195	TYR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	C	503	6/6	0.76	0.14	33,38,38,39	0
4	PGE	A	503	6/10	0.82	0.15	35,37,38,38	0
4	PGE	C	504	6/10	0.83	0.15	28,36,41,44	0
4	PGE	B	503	7/10	0.84	0.13	36,39,43,47	0
5	GOL	C	502	6/6	0.88	0.10	29,36,41,41	0
3	SRT	A	502	10/10	0.88	0.09	31,38,41,41	0
5	GOL	B	502	6/6	0.90	0.15	22,24,34,38	0
2	HEM	C	501	43/43	0.98	0.05	14,17,19,31	0
2	HEM	B	501	43/43	0.98	0.05	15,17,20,31	0
2	HEM	A	501	43/43	0.99	0.05	15,17,21,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around HEM C 501:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.