



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 03:04 AM UTC

PDB ID : 8HQQ / pdb_00008hqq
EMDB ID : EMD-34949
Title : Capsid of DT57C bacteriophage in the empty state
Authors : Ayala, R.; Moiseenko, A.V.; Kulikov, E.E.; Golomidova, A.K.; Orekhov, P.S.;
Street, M.A.; Sokolova, O.S.; Letarov, A.V.; Wolf, M.
Deposited on : 2022-12-13
Resolution : 3.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	12797 (3.10 - 4.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	458	 51% 14% 35%
1	B	458	 55% 10% 35%
1	C	458	 53% 11% 35%
1	D	458	 51% 13% 35%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	458	 50% 14% 35%
1	F	458	 51% 14% 35%
1	G	458	 53% 11% 35%
1	H	458	 51% 13% 35%
1	I	458	 51% 14% 35%
1	J	458	 52% 12% 35%
1	K	458	 52% 13% 35%
1	L	458	 52% 12% 35%
1	M	458	 54% 9% 37%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 29730 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major head protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	B	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	C	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	D	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	E	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	F	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	G	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	H	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	I	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	J	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	K	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	L	296	Total 2291	C 1448	N 389	O 449	S 5	0	0
1	M	288	Total 2238	C 1417	N 379	O 437	S 5	0	0

35%

- Molecule 1: Major head protein

35%

- Molecule 1: Major head protein



35%







• Molecule 1: Major head protein

Chain F: 51% 14% 35%




• Molecule 1: Major head protein


Chain G: 53% 11% 35%



• Molecule 1: Major head protein


Chain H: 51% 13% 35%






• Molecule 1: Major head protein

Chain I: 51% 14% 35%



• Molecule 1: Major head protein

Chain J: 52% 12% 35%



• Molecule 1: Major head protein

Chain K: 52% 13% 35%

D432				
N441				
L442				
Q443				
N448				
A457				
SER				

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	27507	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	67	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	4000	Depositor
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	19.346	Depositor
Minimum map value	-5.025	Depositor
Average map value	0.003	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	5.4	Depositor
Map size (Å)	1248.2999, 1248.2999, 1248.2999	wwPDB
Map dimensions	900, 900, 900	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.387, 1.387, 1.387	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/2326	0.37	0/3144
1	B	0.26	0/2326	0.34	0/3144
1	C	0.27	0/2326	0.36	0/3144
1	D	0.25	0/2326	0.34	0/3144
1	E	0.26	0/2326	0.35	0/3144
1	F	0.27	0/2326	0.37	0/3144
1	G	0.26	0/2326	0.35	0/3144
1	H	0.26	0/2326	0.37	0/3144
1	I	0.26	0/2326	0.37	0/3144
1	J	0.25	0/2326	0.34	0/3144
1	K	0.26	0/2326	0.36	0/3144
1	L	0.25	0/2326	0.36	0/3144
1	M	0.15	0/2272	0.30	0/3068
All	All	0.25	0/30184	0.35	0/40796

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2291	0	2307	49	0
1	B	2291	0	2307	36	0
1	C	2291	0	2307	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2291	0	2307	44	0
1	E	2291	0	2307	56	0
1	F	2291	0	2307	54	0
1	G	2291	0	2307	40	0
1	H	2291	0	2307	49	0
1	I	2291	0	2307	43	0
1	J	2291	0	2307	39	0
1	K	2291	0	2307	45	0
1	L	2291	0	2307	45	0
1	M	2238	0	2257	26	0
All	All	29730	0	29941	471	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 471 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:VAL:HA	1:A:195:LEU:HD22	1.67	0.76
1:D:274:ARG:NH2	1:E:236:THR:O	2.20	0.75
1:K:426:GLN:OE1	1:K:431:ARG:NH2	2.22	0.72
1:K:331:ARG:NH1	1:K:338:GLY:O	2.24	0.71
1:C:331:ARG:NH1	1:C:338:GLY:O	2.24	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/458 (64%)	276 (94%)	18 (6%)	0	100	100
1	B	294/458 (64%)	277 (94%)	17 (6%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	294/458 (64%)	276 (94%)	18 (6%)	0	100	100
1	D	294/458 (64%)	276 (94%)	18 (6%)	0	100	100
1	E	294/458 (64%)	279 (95%)	15 (5%)	0	100	100
1	F	294/458 (64%)	276 (94%)	18 (6%)	0	100	100
1	G	294/458 (64%)	278 (95%)	16 (5%)	0	100	100
1	H	294/458 (64%)	275 (94%)	19 (6%)	0	100	100
1	I	294/458 (64%)	277 (94%)	17 (6%)	0	100	100
1	J	294/458 (64%)	277 (94%)	17 (6%)	0	100	100
1	K	294/458 (64%)	275 (94%)	19 (6%)	0	100	100
1	L	294/458 (64%)	275 (94%)	19 (6%)	0	100	100
1	M	284/458 (62%)	267 (94%)	16 (6%)	1 (0%)	30	61
All	All	3812/5954 (64%)	3584 (94%)	227 (6%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	316	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/382 (64%)	243 (99%)	2 (1%)	73	77
1	B	245/382 (64%)	245 (100%)	0	100	100
1	C	245/382 (64%)	245 (100%)	0	100	100
1	D	245/382 (64%)	244 (100%)	1 (0%)	84	81
1	E	245/382 (64%)	245 (100%)	0	100	100
1	F	245/382 (64%)	244 (100%)	1 (0%)	84	81
1	G	245/382 (64%)	245 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	245/382 (64%)	243 (99%)	2 (1%)	73	77
1	I	245/382 (64%)	245 (100%)	0	100	100
1	J	245/382 (64%)	243 (99%)	2 (1%)	73	77
1	K	245/382 (64%)	244 (100%)	1 (0%)	84	81
1	L	245/382 (64%)	245 (100%)	0	100	100
1	M	240/382 (63%)	240 (100%)	0	100	100
All	All	3180/4966 (64%)	3171 (100%)	9 (0%)	84	83

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	350	MET
1	K	334	LEU
1	F	382	ILE
1	H	350	MET
1	H	436	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	162	ASN
1	M	438	GLN
1	G	416	GLN
1	M	416	GLN
1	L	162	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-34949. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 450




Y Index: 450




Z Index: 450


6.2.2 Raw map



X Index: 450



Y Index: 450




Z Index: 450


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 733




Y Index: 167




Z Index: 167


6.3.2 Raw map



X Index: 733



Y Index: 733




Z Index: 167


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

This section was not generated.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 14845 nm³; this corresponds to an approximate mass of 13410 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.278 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.60	-	-
Author-provided FSC curve	3.62	4.02	3.67
Unmasked-calculated*	4.48	6.27	4.53

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.48 differs from the reported value 3.6 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-34949 and PDB model 8HQQ. Per-residue inclusion information can be found in section [3](#) on page [5](#).

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (5.4).

9.4 Atom inclusion [i](#)



At the recommended contour level, 97% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (5.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8770	 0.5070
A	 0.8850	 0.5110
B	 0.8860	 0.5110
C	 0.8830	 0.5100
D	 0.8810	 0.5060
E	 0.8880	 0.5110
F	 0.8800	 0.5090
G	 0.8780	 0.5090
H	 0.8830	 0.5110
I	 0.8850	 0.5100
J	 0.8790	 0.5080
K	 0.8700	 0.5040
L	 0.8700	 0.5030
M	 0.8270	 0.4850

