



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 02:34 PM UTC

PDB ID : 6HR2 / pdb_00006hr2
Title : Crystal structure of PROTAC 2 in complex with the bromodomain of human SMARCA4 and pVHL:ElonginC:ElonginB
Authors : Roy, M.; Bader, G.; Diers, E.; Trainor, N.; Farnaby, W.; Ciulli, A.
Deposited on : 2018-09-26
Resolution : 1.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	120	
1	E	120	
2	B	149	
2	F	149	
3	C	97	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	97	
4	D	104	
4	H	104	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 15746 atoms, of which 7698 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription activator BRG1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	117	Total	C	H	N	O	S	1005	0	0
			1972	615	1005	167	182	3			
1	E	120	Total	C	H	N	O	S	1045	2	0
			2046	637	1045	172	188	4			

- Molecule 2 is a protein called von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	149	Total	C	H	N	O	S	1264	7	0
			2525	799	1264	234	226	2			
2	F	149	Total	C	H	N	O	S	1241	4	0
			2482	786	1241	230	223	2			

- Molecule 3 is a protein called Elongin-C.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	87	Total	C	H	N	O	S	698	1	0
			1393	448	698	111	129	7			
3	G	87	Total	C	H	N	O	S	697	0	0
			1392	448	697	111	129	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	16	MET	-	initiating methionine	UNP Q15369
G	16	MET	-	initiating methionine	UNP Q15369

- Molecule 4 is a protein called Elongin-B.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	D	104	Total	C	H	N	O	S	821	0	0
			1644	520	821	138	160	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	H	103	Total	C	H	N	O	S	811	0	0
			1624	514	811	136	158	5			

- Molecule 5 is (2 {S},4 {R})- {N}-[[2-[2-[4-[[4-[3-azanyl-6-(2-hydroxyphenyl)pyridazin-4-yl]piperazin-1-yl]methyl]phenyl]ethoxy]-4-(4-methyl-1,3-thiazol-5-yl)phenyl]methyl]-1-[(2 {S})-2-[(1-fluoranylcyclopropyl)carbonylamino]-3,3-dimethyl-butanoyl]-4-oxidanyl-pyrrolidine-2-carboxamide (CCD ID: FWZ) (formula: C₄₉H₅₈FN₉O₆S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
5	B	1	Total	C	F	H	N	O	S	58	0
			124	49	1	58	9	6	1		
5	F	1	Total	C	F	H	N	O	S	58	0
			124	49	1	58	9	6	1		

- Molecule 6 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	O	S	0	0
			4	2	1	1		
6	E	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	F	1	Total	C	O	0	0
			4	2	2		
7	H	1	Total	C	O	0	0
			4	2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	28	Total 28	O 28	0	0
8	B	78	Total 78	O 78	0	0
8	C	37	Total 37	O 37	0	0
8	D	71	Total 71	O 71	0	0
8	E	47	Total 47	O 47	0	0
8	F	71	Total 71	O 71	0	0
8	G	32	Total 32	O 32	0	0
8	H	40	Total 40	O 40	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Transcription activator BRG1




- Molecule 1: Transcription activator BRG1




- Molecule 2: von Hippel-Lindau disease tumor suppressor





- Molecule 2: von Hippel-Lindau disease tumor suppressor




- Molecule 3: Elongin-C




- Molecule 3: Elongin-C



● Molecule 4: Elongin-B



● Molecule 4: Elongin-B



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.33Å 61.68Å 81.60Å 69.43° 83.38° 86.42°	Depositor
Resolution (Å)	39.80 – 1.76 39.80 – 1.76	Depositor EDS
% Data completeness (in resolution range)	63.2 (39.80-1.76) 63.3 (39.80-1.76)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.09 (at 1.76Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.216 , 0.236 0.211 , 0.229	Depositor DCC
R_{free} test set	2903 reflections (4.27%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.017	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15746	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, FWZ, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.75	0/982	1.03	0/1317
1	E	0.76	0/1024	1.04	0/1372
2	B	0.72	0/1309	0.97	2/1785 (0.1%)
2	F	0.74	0/1285	1.00	2/1751 (0.1%)
3	C	0.80	0/718	1.05	2/967 (0.2%)
3	G	0.79	0/709	1.03	1/955 (0.1%)
4	D	0.76	0/839	0.99	1/1132 (0.1%)
4	H	0.72	0/829	1.01	1/1121 (0.1%)
All	All	0.75	0/7695	1.01	9/10400 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	76	PHE	CA-CB-CG	5.39	119.19	113.80
4	D	52	ASP	CA-CB-CG	5.32	117.92	112.60
4	H	40	ASP	CA-CB-CG	5.31	117.91	112.60
3	G	77	PHE	CA-CB-CG	5.31	119.11	113.80
3	C	77	PHE	CA-CB-CG	5.30	119.10	113.80
3	C	85	ASN	CA-CB-CG	5.16	117.76	112.60
2	B	76	PHE	CA-CB-CG	5.15	118.95	113.80
2	F	83	VAL	N-CA-C	-5.05	101.07	108.54
2	B	83	VAL	N-CA-C	-5.03	101.09	108.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	967	1005	1005	0	0
1	E	1001	1045	1033	0	0
2	B	1261	1264	1242	4	0
2	F	1241	1241	1225	3	0
3	C	695	698	691	0	0
3	G	695	697	697	0	0
4	D	823	821	824	0	0
4	H	813	811	811	0	0
5	B	66	58	0	0	0
5	F	66	58	0	0	0
6	B	4	0	6	0	0
6	E	4	0	6	0	0
7	F	4	0	6	0	0
7	H	4	0	6	0	0
8	A	28	0	0	0	0
8	B	78	0	0	0	0
8	C	37	0	0	0	0
8	D	71	0	0	0	0
8	E	47	0	0	0	0
8	F	71	0	0	0	0
8	G	32	0	0	0	0
8	H	40	0	0	0	0
All	All	8048	7698	7552	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:139[A]:SER:HB2	2:F:139:SER:OG	2.12	0.49
2:B:115:HIS:O	2:B:138[A]:PRO:HD2	2.15	0.47
2:B:140[A]:LEU:HB2	2:F:140:LEU:HB2	1.98	0.45
2:B:133:THR:HG21	2:F:145:GLN:HG2	2.00	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/120 (96%)	114 (99%)	1 (1%)	0	100	100
1	E	120/120 (100%)	118 (98%)	2 (2%)	0	100	100
2	B	154/149 (103%)	149 (97%)	5 (3%)	0	100	100
2	F	151/149 (101%)	146 (97%)	5 (3%)	0	100	100
3	C	84/97 (87%)	84 (100%)	0	0	100	100
3	G	83/97 (86%)	82 (99%)	1 (1%)	0	100	100
4	D	102/104 (98%)	99 (97%)	3 (3%)	0	100	100
4	H	101/104 (97%)	96 (95%)	5 (5%)	0	100	100
All	All	910/940 (97%)	888 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/116 (97%)	111 (98%)	2 (2%)	51	33
1	E	118/116 (102%)	113 (96%)	5 (4%)	26	8
2	B	146/139 (105%)	140 (96%)	6 (4%)	27	8
2	F	142/139 (102%)	138 (97%)	4 (3%)	38	18

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	80/86 (93%)	80 (100%)	0	100	100
3	G	79/86 (92%)	79 (100%)	0	100	100
4	D	92/92 (100%)	91 (99%)	1 (1%)	65	52
4	H	91/92 (99%)	90 (99%)	1 (1%)	65	52
All	All	861/866 (99%)	842 (98%)	19 (2%)	50	26

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1488	LEU
1	A	1568	ASP
2	B	140[A]	LEU
2	B	140[B]	LEU
2	B	141[A]	ASN
2	B	141[B]	ASN
2	B	182	ARG
2	B	196	LYS
4	D	37	ARG
1	E	1451	LEU
1	E	1488	LEU
1	E	1490	SER
1	E	1522	LEU
1	E	1568	ASP
2	F	113[A]	ARG
2	F	113[B]	ARG
2	F	140	LEU
2	F	196	LYS
4	H	46	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1457	ASN
1	A	1537	GLN
1	A	1554	GLN
2	B	191	HIS
3	C	61	ASN
1	E	1457	ASN
1	E	1537	GLN
1	E	1554	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	F	208	HIS
3	G	27	HIS
3	G	61	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	EDO	F	302	-	3,3,3	0.49	0	2,2,2	0.41	0
7	EDO	H	201	-	3,3,3	0.51	0	2,2,2	0.30	0
6	DMS	E	1601	-	3,3,3	0.30	0	3,3,3	0.36	0
5	FWZ	F	301	-	71,73,73	0.41	1 (1%)	97,107,107	0.82	1 (1%)
6	DMS	B	302	-	3,3,3	0.37	0	3,3,3	0.11	0
5	FWZ	B	301	-	71,73,73	0.29	0	97,107,107	0.76	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	H	201	-	-	0/1/1/1	-
7	EDO	F	302	-	-	0/1/1/1	-
5	FWZ	B	301	-	-	3/53/81/81	0/8/8/8
5	FWZ	F	301	-	-	2/53/81/81	0/8/8/8

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	301	FWZ	C47-N44	2.07	1.45	1.41

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	301	FWZ	C36-C32-C31	4.90	126.22	116.20
5	B	301	FWZ	C36-C32-C31	3.97	124.31	116.20

There are no chirality outliers.


All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	F	301	FWZ	C64-C40-N41-C46
5	F	301	FWZ	C64-C40-N41-C42
5	B	301	FWZ	C64-C40-N41-C46
5	B	301	FWZ	C64-C40-N41-C42
5	B	301	FWZ	C48-C47-N44-C45

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	117/120 (97%)	1.47	33 (28%)	1 1	14, 30, 64, 117	0
1	E	120/120 (100%)	0.92	17 (14%)	6 7	8, 24, 60, 111	1 (0%)
2	B	149/149 (100%)	1.07	29 (19%)	3 3	8, 20, 62, 84	5 (3%)
2	F	149/149 (100%)	1.37	33 (22%)	2 2	8, 21, 89, 128	2 (1%)
3	C	87/97 (89%)	0.91	10 (11%)	9 11	14, 23, 42, 80	0
3	G	87/97 (89%)	1.28	19 (21%)	2 2	17, 25, 45, 84	0
4	D	104/104 (100%)	0.70	12 (11%)	9 11	11, 20, 54, 105	0
4	H	103/104 (99%)	1.08	16 (15%)	5 5	15, 25, 51, 124	0
All	All	916/940 (97%)	1.11	169 (18%)	3 3	8, 23, 67, 128	8 (0%)

All (169) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	181	VAL	11.6
2	B	140[A]	LEU	10.1
4	D	102	VAL	10.1
2	F	184	LEU	9.1
2	F	178	LEU	8.9
2	F	180	ILE	8.4
4	H	99	LEU	8.0
2	B	139[A]	SER	8.0
2	F	175	TYR	8.0
4	H	103	MET	7.9
4	H	102	VAL	7.8
2	B	178	LEU	7.6
2	B	180	ILE	7.4
4	D	103	MET	7.2
2	B	181	VAL	6.8
2	F	185	TYR	6.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	100	PRO	6.5
1	A	1564	ILE	6.2
4	D	99	LEU	6.2
2	F	170	VAL	6.1
2	B	175	TYR	6.1
4	H	100	PRO	6.0
2	F	140	LEU	5.9
2	F	188	LEU	5.8
3	G	46	LEU	5.4
2	F	174	ASN	5.2
2	B	184	LEU	4.9
1	E	1564	ILE	4.8
2	F	169	LEU	4.8
2	B	174	ASN	4.7
2	F	62	VAL	4.7
2	B	172	PRO	4.7
2	F	182	ARG	4.6
2	F	206	ILE	4.6
2	B	177	ARG	4.5
1	E	1451	LEU	4.5
2	F	183	SER	4.5
2	F	179	ASP	4.4
2	F	187	ASP	4.4
1	A	1453	PRO	4.3
2	F	172	PRO	4.3
1	E	1453	PRO	4.2
4	D	101	ASP	4.2
1	A	1452	SER	4.2
4	H	97	PRO	4.1
2	B	209	GLN	4.1
3	G	45	MET	4.1
2	B	182	ARG	3.9
2	B	206	ILE	3.8
2	B	185	TYR	3.8
2	F	177	ARG	3.8
4	D	104	LYS	3.8
2	F	176	ARG	3.8
4	D	97	PRO	3.8
1	E	1450	LYS	3.8
3	C	46	LEU	3.8
2	B	183	SER	3.8
3	G	47	SER	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1455	PRO	3.7
1	A	1472	TYR	3.7
3	C	47	SER	3.7
2	F	139	SER	3.6
2	B	61	PRO	3.6
2	B	176	ARG	3.6
2	F	207	ALA	3.5
4	H	1	MET	3.4
1	A	1492	LYS	3.4
1	E	1467	ASP	3.4
4	D	36	LYS	3.3
2	F	113[A]	ARG	3.3
1	E	1458	LEU	3.3
1	E	1568	ASP	3.3
2	B	193	ASN	3.3
3	C	88	THR	3.2
3	G	88	THR	3.2
3	C	85	ASN	3.2
1	A	1527	LYS	3.2
3	G	17	MET	3.1
1	A	1470	ILE	3.1
2	F	171	LYS	3.1
2	B	188	LEU	3.1
2	B	170	VAL	3.1
1	A	1481	LEU	3.1
3	C	63	ARG	3.0
1	A	1522	LEU	3.0
1	A	1566	LYS	3.0
1	E	1455	PRO	3.0
4	H	96	PRO	3.0
1	E	1449	GLU	3.0
4	D	96	PRO	3.0
1	A	1469	VAL	2.9
1	E	1522	LEU	2.9
1	E	1452	SER	2.9
3	G	34	GLU	2.9
1	A	1568	ASP	2.9
3	C	45	MET	2.9
4	D	1	MET	2.9
4	D	83	ASP	2.9
1	A	1556	VAL	2.8
3	G	44	ALA	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	1563	LYS	2.8
2	F	141	ASN	2.8
4	D	98	GLU	2.8
4	H	84	THR	2.8
3	G	42	ILE	2.8
1	A	1456	PRO	2.7
1	A	1562	GLN	2.7
1	A	1514	ILE	2.7
1	A	1473	LYS	2.7
1	E	1566	LYS	2.7
2	F	186	GLU	2.7
1	A	1475	SER	2.7
1	A	1558	THR	2.6
1	A	1477	SER	2.6
3	C	60	VAL	2.6
3	G	60	VAL	2.6
1	A	1557	PHE	2.6
4	H	85	PHE	2.6
1	A	1454	ASN	2.6
1	A	1467	ASP	2.5
3	G	16	MET	2.5
4	H	101	ASP	2.5
2	B	208	HIS	2.5
4	H	25	PHE	2.5
2	B	169	LEU	2.5
2	B	179	ASP	2.5
1	E	1508	LYS	2.5
4	H	98	GLU	2.5
2	B	105	THR	2.4
3	G	99	ILE	2.4
4	H	64	SER	2.4
2	B	141[A]	ASN	2.4
1	A	1502	ARG	2.4
3	G	58	ASN	2.4
2	F	193	ASN	2.3
1	A	1567	GLU	2.3
1	A	1500	LEU	2.3
2	B	207	ALA	2.3
2	F	142	VAL	2.3
1	E	1454	ASN	2.3
1	A	1471	LYS	2.3
2	B	173	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	205	ARG	2.3
4	H	36	LYS	2.3
1	A	1516	ASN	2.2
2	B	171	LYS	2.2
3	G	62	PHE	2.2
3	G	90	ILE	2.2
3	C	43	LYS	2.2
2	F	168	SER	2.2
2	F	144	GLY	2.2
1	A	1560	VAL	2.1
1	A	1517	HIS	2.1
2	F	208	HIS	2.1
1	A	1563	LYS	2.1
3	C	16	MET	2.1
3	G	64	GLU	2.1
2	B	165	VAL	2.1
1	E	1567	GLU	2.1
3	C	64[A]	GLU	2.1
3	G	61	ASN	2.1
2	F	173	GLU	2.1
4	H	49	GLN	2.1
1	E	1476	SER	2.1
3	G	43	LYS	2.1
3	G	18	TYR	2.1
3	G	89	GLU	2.0
1	A	1476	SER	2.0
4	H	46	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	DMS	B	302	4/4	0.84	0.24	81,82,82,83	0
6	DMS	E	1601	4/4	0.84	0.23	72,73,74,74	0
7	EDO	F	302	4/4	0.84	0.16	42,44,47,48	0
7	EDO	H	201	4/4	0.85	0.16	65,66,66,66	0
5	FWZ	B	301	66/66	0.95	0.07	29,34,42,45	58
5	FWZ	F	301	66/66	0.96	0.07	22,31,37,38	58

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.