



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:43 PM UTC

PDB ID : 1HSR / pdb_00001hsr
Title : BINDING MODE OF BENZHYDROXAMIC ACID TO ARTHROMYCES
RAMOSUS PEROXIDASE
Authors : Fukuyama, K.; Itakura, H.
Deposited on : 1997-07-01
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	344	
2	B	2	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2789 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2465	1537	421	492	15			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: C₃₄H₃₂FeN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 5 is BENZHYDROXAMIC ACID (CCD ID: BHO) (formula: $C_7H_7NO_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			10	7	1	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	241	Total 241	O 241	0	0

Note EDS was not executed.

Chain A: 78% 15% ...



Chain B: 100%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.50Å 74.50Å 118.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.60	Depositor
% Data completeness (in resolution range)	(Not available) (7.00-1.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.187 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2789	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, BHO, NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	3/2521 (0.1%)	1.62	46/3436 (1.3%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	86	ILE	CA-CB	6.30	1.62	1.54
1	A	318	ILE	CA-CB	6.27	1.62	1.54
1	A	29	VAL	CA-CB	5.62	1.62	1.54

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	202	ASP	CA-CB-CG	11.08	123.68	112.60
1	A	10	VAL	N-CA-C	-10.77	91.75	108.23
1	A	309	ILE	N-CA-C	-10.74	96.92	107.76
1	A	210	THR	N-CA-CB	-9.72	94.08	110.41
1	A	221	THR	N-CA-CB	-8.53	98.18	110.81
1	A	264	SER	N-CA-C	7.74	122.91	113.38
1	A	295	ASP	CA-CB-CG	7.60	120.20	112.60
1	A	83	HIS	CA-CB-CG	-7.51	106.29	113.80
1	A	20	ASN	CA-CB-CG	7.33	119.93	112.60
1	A	210	THR	OG1-CB-CG2	7.30	123.89	109.30
1	A	237	PHE	CA-CB-CG	7.18	120.98	113.80
1	A	34	GLN	OE1-CD-NE2	-6.52	116.08	122.60
1	A	85	ASN	OD1-CG-ND2	-6.42	116.18	122.60
1	A	70	GLN	OE1-CD-NE2	-6.38	116.22	122.60
1	A	83	HIS	CB-CG-CD2	-6.23	123.10	131.20
1	A	304	ASN	OD1-CG-ND2	-6.17	116.44	122.60
1	A	110	HIS	ND1-CG-CD2	6.12	112.22	106.10
1	A	119	GLN	OE1-CD-NE2	-6.00	116.60	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	110	HIS	CB-CG-CD2	-5.97	123.44	131.20
1	A	158	ASN	OD1-CG-ND2	-5.96	116.64	122.60
1	A	110	HIS	CA-CB-CG	-5.90	107.90	113.80
1	A	110	HIS	CG-CD2-NE2	-5.89	101.31	107.20
1	A	11	THR	N-CA-C	-5.82	99.70	109.07
1	A	155	GLY	CA-C-N	5.77	125.39	119.56
1	A	155	GLY	C-N-CA	5.77	125.39	119.56
1	A	193	ASN	OD1-CG-ND2	-5.77	116.83	122.60
1	A	339	SER	N-CA-C	-5.67	99.04	108.34
1	A	39	GLN	OE1-CD-NE2	-5.63	116.97	122.60
1	A	306	ALA	CA-C-N	5.59	125.52	119.76
1	A	306	ALA	C-N-CA	5.59	125.52	119.76
1	A	187	ALA	N-CA-C	5.59	117.76	108.99
1	A	318	ILE	CB-CG1-CD1	-5.46	102.34	113.80
1	A	223	GLN	N-CA-C	-5.44	98.90	108.69
1	A	83	HIS	CG-CD2-NE2	-5.35	101.85	107.20
1	A	210	THR	CB-CA-C	5.35	120.60	109.95
1	A	148	SER	CA-C-N	5.34	123.56	119.66
1	A	148	SER	C-N-CA	5.34	123.56	119.66
1	A	311	GLY	N-CA-C	5.32	121.24	113.48
1	A	288	ASN	OD1-CG-ND2	-5.24	117.36	122.60
1	A	83	HIS	ND1-CG-CD2	5.19	111.29	106.10
1	A	199	SER	CA-C-N	5.17	125.07	119.85
1	A	199	SER	C-N-CA	5.17	125.07	119.85
1	A	97	THR	N-CA-C	5.11	117.24	111.11
1	A	128	ASN	OD1-CG-ND2	-5.05	117.55	122.60
1	A	237	PHE	CA-C-N	5.04	125.04	119.90
1	A	237	PHE	C-N-CA	5.04	125.04	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2465	0	2373	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	28	0	25	0	0
3	A	2	0	0	0	0
4	A	43	0	30	0	0
5	A	10	0	7	0	0
6	A	241	0	0	4	0
All	All	2789	0	2435	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:MET:SD	6:A:843:HOH:O	2.40	0.79
1:A:86:ILE:HG12	1:A:147:PRO:HB3	1.81	0.62
1:A:34:GLN:HE21	1:A:34:GLN:HA	1.66	0.60
1:A:32:ASP:OD2	1:A:110:HIS:HE1	1.87	0.58
1:A:217:LEU:HD13	1:A:250:ALA:HB1	1.87	0.57
1:A:210:THR:HG22	6:A:530:HOH:O	2.07	0.55
1:A:138:LEU:HD12	1:A:292:ASP:HA	1.91	0.53
1:A:180:LEU:HD13	1:A:276:ALA:HB1	1.93	0.51
1:A:243:MET:HE3	1:A:245:SER:OG	2.12	0.49
1:A:184:HIS:HE1	1:A:243:MET:CE	2.26	0.48
1:A:221:THR:HG22	1:A:222:THR:OG1	2.14	0.48
1:A:26:TRP:HA	1:A:29:VAL:HG13	1.96	0.48
1:A:201:LEU:HD21	6:A:843:HOH:O	2.15	0.46
1:A:86:ILE:HG13	1:A:87:GLU:N	2.32	0.45
1:A:85:ASN:HD22	1:A:85:ASN:H	1.67	0.42
1:A:29:VAL:HG22	1:A:120:PHE:CE2	2.54	0.42
1:A:221:THR:HG21	6:A:538:HOH:O	2.19	0.42
1:A:219:LYS:HG3	1:A:317:ASP:O	2.21	0.41
1:A:184:HIS:HE1	1:A:243:MET:HE2	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/344 (97%)	326 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/273 (99%)	245 (90%)	26 (10%)	8	1

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	SER
1	A	13	PRO
1	A	29	VAL
1	A	34	GLN
1	A	50	ILE
1	A	65	LEU
1	A	85	ASN
1	A	86	ILE
1	A	135	LEU
1	A	138	LEU
1	A	180	LEU
1	A	181	LEU
1	A	186	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	192	LEU
1	A	210	THR
1	A	217	LEU
1	A	218	LEU
1	A	219	LYS
1	A	221	THR
1	A	228	LEU
1	A	234	LEU
1	A	237	PHE
1	A	248	LEU
1	A	283	LEU
1	A	318	ILE
1	A	332	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	85	ASN
1	A	110	HIS
1	A	128	ASN
1	A	193	ASN
1	A	206	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.66	0	17,19,21	1.46	6 (35%)
2	NAG	B	2	2	14,14,15	0.49	0	17,19,21	1.26	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C6-C5-C4	-2.39	107.14	113.02
2	B	1	NAG	O5-C1-C2	-2.22	107.85	111.29
2	B	2	NAG	C6-C5-C4	2.20	118.42	113.02
2	B	2	NAG	O5-C5-C4	-2.10	105.71	110.83
2	B	1	NAG	C8-C7-N2	2.06	119.53	116.12
2	B	1	NAG	C3-C4-C5	2.05	113.95	110.23
2	B	1	NAG	C1-O5-C5	2.05	114.93	112.19
2	B	1	NAG	C1-C2-N2	2.03	113.63	110.43

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	BHO	A	800	-	10,10,10	1.82	2 (20%)	12,12,12	1.03	1 (8%)
4	HEM	A	345	6,1	50,50,50	1.14	6 (12%)	67,82,82	0.92	4 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BHO	A	800	-	-	0/6/6/6	0/1/1/1
4	HEM	A	345	6,1	-	4/14/54/54	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	800	BHO	C1-C	-4.02	1.41	1.50
4	A	345	HEM	CBC-CAC	3.27	1.46	1.30
4	A	345	HEM	CAB-C3B	-2.75	1.40	1.47
4	A	345	HEM	CBB-CAB	2.74	1.43	1.30
4	A	345	HEM	CAC-C3C	-2.51	1.40	1.47
5	A	800	BHO	C-N	2.27	1.35	1.32
4	A	345	HEM	O2A-CGA	-2.16	1.23	1.30
4	A	345	HEM	O2D-CGD	-2.15	1.23	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	345	HEM	C3B-C4B-NB	2.89	111.55	109.47
5	A	800	BHO	O2-N-C	-2.75	112.84	119.73
4	A	345	HEM	O2D-CGD-CBD	2.12	120.70	114.00
4	A	345	HEM	C1D-C2D-C3D	-2.11	104.76	106.98
4	A	345	HEM	C4C-C3C-C2C	-2.09	105.00	106.81

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	345	HEM	C2B-C3B-CAB-CBB
4	A	345	HEM	C4B-C3B-CAB-CBB
4	A	345	HEM	CAA-CBA-CGA-O1A
4	A	345	HEM	CAA-CBA-CGA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.