



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 18, 2026 – 12:44 PM UTC

PDB ID : 2HSJ / pdb_00002hsj
Title : The structure of a putative platelet activating factor from Streptococcus pneumonia.
Authors : Cuff, M.E.; Mulligan, R.; Abdullah, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2006-07-21
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 4% 89% 8% ..
1	B	214	 4% 93% 6% .
1	C	214	 4% 93% 6% .
1	D	214	 4% 95% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	1404	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative platelet activating factor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	211	1772	1139	290	339	1	3	0	8	0
1	B	212	1765	1131	291	339	2	2	0	7	0
1	C	212	1758	1128	288	337	2	3	0	7	0
1	D	214	1782	1147	292	339	1	3	0	6	0


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	cloning artifact	UNP Q97PY9
A	-1	ASN	-	cloning artifact	UNP Q97PY9
A	0	ALA	-	cloning artifact	UNP Q97PY9
A	1	MSE	MET	modified residue	UNP Q97PY9
A	165	MSE	MET	modified residue	UNP Q97PY9
B	-2	SER	-	cloning artifact	UNP Q97PY9
B	-1	ASN	-	cloning artifact	UNP Q97PY9
B	0	ALA	-	cloning artifact	UNP Q97PY9
B	1	MSE	MET	modified residue	UNP Q97PY9
B	165	MSE	MET	modified residue	UNP Q97PY9
C	-2	SER	-	cloning artifact	UNP Q97PY9
C	-1	ASN	-	cloning artifact	UNP Q97PY9
C	0	ALA	-	cloning artifact	UNP Q97PY9
C	1	MSE	MET	modified residue	UNP Q97PY9
C	165	MSE	MET	modified residue	UNP Q97PY9
D	-2	SER	-	cloning artifact	UNP Q97PY9
D	-1	ASN	-	cloning artifact	UNP Q97PY9
D	0	ALA	-	cloning artifact	UNP Q97PY9
D	1	MSE	MET	modified residue	UNP Q97PY9
D	165	MSE	MET	modified residue	UNP Q97PY9

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	3	Total Mg 3 3	0	0
2	C	1	Total Mg 1 1	0	0
2	D	2	Total Mg 2 2	0	0

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	305	Total	O	0	0
			305	305		
4	B	318	Total	O	0	0
			318	318		
4	C	350	Total	O	0	0
			350	350		
4	D	293	Total	O	0	0
			293	293		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.03Å 120.06Å 84.74Å 90.00° 99.48° 90.00°	Depositor
Resolution (Å)	40.46 – 1.50 40.46 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (40.46-1.50) 99.4 (40.46-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.160 , 0.193 0.159 , 0.192	Depositor DCC
R_{free} test set	6428 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	12.8	Xtrriage
Anisotropy	0.133	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8404	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.68	0/1805	0.78	0/2447
1	B	0.70	0/1798	0.76	0/2437
1	C	0.72	0/1788	0.81	2/2423 (0.1%)
1	D	0.67	0/1815	0.76	0/2458
All	All	0.69	0/7206	0.78	2/9765 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	165	MSE	CA-C-N	-7.83	110.12	122.73
1	C	165	MSE	C-N-CA	-7.83	110.12	122.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1772	0	1767	20	0
1	B	1765	0	1747	11	0
1	C	1758	0	1756	13	0
1	D	1782	0	1776	8	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	3	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	12	0	16	1	0
3	B	18	0	24	5	0
3	C	6	0	8	0	0
3	D	18	0	24	1	0
4	A	305	0	0	7	0
4	B	318	0	0	3	0
4	C	350	0	0	8	0
4	D	293	0	0	2	0
All	All	8404	0	7118	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:34[A]:ILE:HD13	1:A:210:LEU:CD2	1.94	0.97
1:C:11[B]:LEU:HG	4:C:1757:HOH:O	1.75	0.87
1:B:33[A]:ASN:ND2	1:B:84:ASP:OD2	2.15	0.80
1:D:121:LEU:HB2	3:D:1407:GOL:H12	1.66	0.78
1:A:34[A]:ILE:HD13	1:A:210:LEU:HD22	1.65	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	216/214 (101%)	211 (98%)	5 (2%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	217/214 (101%)	213 (98%)	4 (2%)	0	100	100
1	C	217/214 (101%)	212 (98%)	5 (2%)	0	100	100
1	D	218/214 (102%)	213 (98%)	5 (2%)	0	100	100
All	All	868/856 (101%)	849 (98%)	19 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/186 (104%)	189 (97%)	5 (3%)	40	13
1	B	193/186 (104%)	190 (98%)	3 (2%)	55	28
1	C	193/186 (104%)	191 (99%)	2 (1%)	68	45
1	D	194/186 (104%)	190 (98%)	4 (2%)	47	19
All	All	774/744 (104%)	760 (98%)	14 (2%)	55	24

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	136	GLU
1	C	4	GLN
1	D	65	TYR
1	D	1[B]	MSE
1	D	4	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	200	GLN
1	C	4	GLN
1	C	166	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	139	GLN
1	A	151	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 7 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	1404	-	5,5,5	0.68	0	5,5,5	0.53	0
3	GOL	B	1403	-	5,5,5	0.46	0	5,5,5	0.44	0
3	GOL	B	1405	-	5,5,5	0.35	0	5,5,5	0.44	0
3	GOL	A	1402	-	5,5,5	0.21	0	5,5,5	0.78	0
3	GOL	C	1408	-	5,5,5	0.22	0	5,5,5	0.77	0
3	GOL	D	1401	-	5,5,5	0.36	0	5,5,5	0.65	0
3	GOL	A	1409	-	5,5,5	0.41	0	5,5,5	0.61	0
3	GOL	D	1406	-	5,5,5	0.34	0	5,5,5	0.68	0
3	GOL	D	1407	-	5,5,5	0.33	0	5,5,5	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	1404	-	-	2/4/4/4	-
3	GOL	B	1403	-	-	0/4/4/4	-
3	GOL	B	1405	-	-	0/4/4/4	-
3	GOL	A	1402	-	-	0/4/4/4	-
3	GOL	C	1408	-	-	0/4/4/4	-
3	GOL	D	1401	-	-	0/4/4/4	-
3	GOL	A	1409	-	-	3/4/4/4	-
3	GOL	D	1406	-	-	0/4/4/4	-
3	GOL	D	1407	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1409	GOL	C1-C2-C3-O3
3	A	1409	GOL	O2-C2-C3-O3
3	B	1404	GOL	O1-C1-C2-C3
3	D	1407	GOL	C1-C2-C3-O3
3	B	1404	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1404	GOL	5	0
3	A	1409	GOL	1	0
3	D	1407	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	209/214 (97%)	-0.05	8 (3%) 44 48	4, 12, 23, 37	7 (3%)
1	B	210/214 (98%)	-0.12	8 (3%) 44 48	5, 11, 21, 41	7 (3%)
1	C	210/214 (98%)	-0.21	9 (4%) 40 44	5, 10, 21, 46	6 (2%)
1	D	212/214 (99%)	0.25	8 (3%) 44 48	5, 13, 22, 44	5 (2%)
All	All	841/856 (98%)	-0.03	33 (3%) 43 47	4, 12, 22, 46	25 (2%)

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	0	ALA	5.2
1	B	9[A]	TRP	4.6
1	C	3	VAL	4.5
1	B	5	LEU	4.5
1	D	-2	SER	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	B	1405	6/6	0.70	0.18	34,41,43,44	0
3	GOL	D	1407	6/6	0.83	0.13	31,36,38,39	0
3	GOL	A	1409	6/6	0.85	0.12	26,28,30,35	0
3	GOL	B	1404	6/6	0.86	0.11	20,23,30,30	0
3	GOL	D	1406	6/6	0.88	0.12	21,25,27,29	0
3	GOL	C	1408	6/6	0.91	0.13	15,15,18,19	0
2	MG	B	1306	1/1	0.91	0.16	34,34,34,34	0
3	GOL	B	1403	6/6	0.91	0.13	14,16,17,17	0
2	MG	D	1307	1/1	0.96	0.11	32,32,32,32	0
3	GOL	D	1401	6/6	0.97	0.12	15,17,18,20	0
2	MG	B	1302	1/1	0.98	0.22	17,17,17,17	0
3	GOL	A	1402	6/6	0.98	0.12	13,15,16,16	0
2	MG	B	1305	1/1	0.98	0.08	14,14,14,14	0
2	MG	A	1301	1/1	0.98	0.10	12,12,12,12	0
2	MG	C	1303	1/1	0.98	0.21	19,19,19,19	0
2	MG	D	1304	1/1	0.99	0.11	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.