



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:50 PM UTC

PDB ID : 5HSR / pdb_00005hsr
Title : Fluorine substituted 5-methyl-6-(3',4'-difluoromethoxyphenylthio)thieno[2,3-d
pyrimidine-2,4-diamine
Authors : Cody, V.
Deposited on : 2016-01-26
Resolution : 1.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2002 (1.24-1.20)
Clashscore	190562	2061 (1.24-1.20)
Ramachandran outliers	187476	2009 (1.24-1.20)
Sidechain outliers	187428	2008 (1.24-1.20)
RSRZ outliers	180081	2000 (1.24-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	63Y	A	202	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 1822 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydrofolate reductase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	186	1544	990	258	287	9	0	6	0

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is 5-methyl-6-[(2,3,4-trifluorophenyl)sulfonyl]thieno[2,3-d]pyrimidine-2,4-diamine (CCD ID: 63Y) (formula: $C_{13}H_9F_3N_4S_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	S	0	0
			21	13	2	4	2		
3	A	1	Total	C	F	N	S	0	0
			21	13	2	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	188	Total	O	0	0
			188	188		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydrofolate reductase



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	84.61Å 84.61Å 78.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.72 – 1.21 26.72 – 1.21	Depositor EDS
% Data completeness (in resolution range)	98.7 (26.72-1.21) 98.7 (26.72-1.21)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.02	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.21Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.193 , 0.219 0.190 , 0.215	Depositor DCC
R_{free} test set	3157 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	13.7	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 40.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	1822	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, 63Y

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.70	15/1583 (0.9%)	1.37	5/2133 (0.2%)

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	14	MET	SD-CE	-13.13	1.46	1.79
1	A	126	ASN	CG-ND2	9.24	1.52	1.33
1	A	87	HIS	CE1-NE2	8.59	1.41	1.32
1	A	87	HIS	CD2-NE2	-8.43	1.28	1.37
1	A	74	VAL	CA-C	-7.31	1.44	1.52
1	A	72	ASN	C-O	6.98	1.32	1.24
1	A	126	ASN	CB-CG	6.78	1.69	1.52
1	A	60	ILE	C-N	6.22	1.40	1.33
1	A	70	ARG	CZ-NH1	5.66	1.40	1.32
1	A	49	LEU	N-CA	5.64	1.53	1.46
1	A	60	ILE	C-O	-5.49	1.17	1.24
1	A	88	PHE	N-CA	-5.42	1.39	1.45
1	A	87	HIS	C-O	5.40	1.31	1.23
1	A	41	SER	N-CA	5.35	1.52	1.46
1	A	87	HIS	CA-C	-5.35	1.45	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	126	ASN	CB-CG-ND2	7.61	127.81	116.40
1	A	132	LYS	CD-CE-NZ	-7.01	89.47	111.90
1	A	146	THR	OG1-CB-CG2	-6.78	95.74	109.30
1	A	70	ARG	NE-CZ-NH2	6.17	124.76	119.20
1	A	87	HIS	CB-CG-ND1	5.38	130.77	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1544	0	1549	21	0
2	A	48	0	26	5	0
3	A	42	0	0	9	0
4	A	188	0	0	4	0
All	All	1822	0	1575	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:PRO:HD2	3:A:202:63Y:FAE	1.28	1.23
2:A:201:NDP:H42N	3:A:202:63Y:CAA	1.71	1.19
1:A:61:PRO:CD	3:A:202:63Y:FAE	1.78	1.19
1:A:99:LEU:HD22	1:A:105:LEU:HD12	1.45	0.98
1:A:99:LEU:CD2	1:A:105:LEU:HD12	1.94	0.96
1:A:72:ASN:H	1:A:87:HIS:HD2	1.14	0.91
1:A:61:PRO:HD3	3:A:202:63Y:FAE	1.66	0.85
1:A:126:ASN:OD1	4:A:301:HOH:O	2.03	0.76
1:A:72:ASN:H	1:A:87:HIS:CD2	2.03	0.75
2:A:201:NDP:C4N	3:A:202:63Y:CAA	2.59	0.75
1:A:99:LEU:HD22	1:A:105:LEU:CD1	2.19	0.71
1:A:99:LEU:CD2	1:A:105:LEU:CD1	2.72	0.67
2:A:201:NDP:H42N	3:A:202:63Y:CAP	2.27	0.65
1:A:145:ASP:OD1	1:A:146:THR:HG23	1.95	0.65
1:A:14:MET:HA	1:A:14:MET:HE2	1.83	0.61
1:A:126:ASN:CG	4:A:301:HOH:O	2.44	0.60
1:A:16:ILE:O	2:A:201:NDP:H2N	2.10	0.51
1:A:61:PRO:HD2	3:A:202:63Y:CAR	2.37	0.50
1:A:132:LYS:HE2	4:A:340:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:LYS:CE	4:A:340:HOH:O	2.62	0.48
1:A:99:LEU:O	1:A:99:LEU:HD23	2.15	0.46
2:A:201:NDP:C5N	3:A:202:63Y:CAA	2.94	0.45
1:A:107:ASN:OD1	1:A:107:ASN:N	2.39	0.44
1:A:60:ILE:HG23	3:A:202:63Y:CAN	2.50	0.41
1:A:99:LEU:HD23	1:A:105:LEU:HD12	1.92	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/186 (102%)	187 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/168 (104%)	171 (98%)	3 (2%)	53	17

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	GLU
1	A	62	GLU
1	A	101	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	HIS
1	A	127	HIS
1	A	185	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	201	-	51,52,52	1.83	16 (31%)	71,80,80	1.99	12 (16%)
3	63Y	A	203	3	23,23,24	3.28	6 (26%)	28,34,36	2.61	11 (39%)
3	63Y	A	202	3	23,23,24	2.79	6 (26%)	28,34,36	2.56	10 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	201	-	-	6/34/77/77	0/5/5/5
3	63Y	A	203	3	-	0/4/4/4	0/3/3/3
3	63Y	A	202	3	-	0/4/4/4	0/3/3/3

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	203	63Y	CAQ-SAK	-13.47	1.60	1.78
3	A	202	63Y	C5-C6	-7.58	1.34	1.42
3	A	202	63Y	CAT-SAL	-5.65	1.60	1.75
3	A	202	63Y	C4-SAL	-5.26	1.65	1.74
3	A	202	63Y	C5-C4	-4.67	1.35	1.40
3	A	202	63Y	C5-CAP	-4.47	1.35	1.45
2	A	201	NDP	C6N-N1N	4.07	1.47	1.37
2	A	201	NDP	PA-O3	3.97	1.63	1.59
2	A	201	NDP	PN-O5D	-3.73	1.44	1.59
3	A	203	63Y	CAA-CAP	3.66	1.56	1.50
3	A	203	63Y	CAH-CAG	3.57	1.44	1.38
2	A	201	NDP	C2A-N1A	3.48	1.40	1.33
2	A	201	NDP	C7N-C3N	-3.21	1.41	1.48
2	A	201	NDP	C5D-C4D	3.05	1.60	1.51
3	A	202	63Y	CAP-CAT	-2.99	1.33	1.37
2	A	201	NDP	C8A-N7A	2.88	1.37	1.31
2	A	201	NDP	PN-O3	-2.70	1.56	1.59
2	A	201	NDP	C2D-C1D	-2.44	1.45	1.53
2	A	201	NDP	PA-O1A	-2.44	1.42	1.50
3	A	203	63Y	C2-N3	2.41	1.39	1.35
2	A	201	NDP	PA-O2A	-2.40	1.44	1.55
3	A	203	63Y	CAP-CAT	-2.37	1.34	1.37
2	A	201	NDP	PN-O2N	-2.28	1.44	1.55
2	A	201	NDP	C7N-N7N	2.25	1.39	1.33
2	A	201	NDP	P2B-O1X	-2.23	1.43	1.50
2	A	201	NDP	C6N-C5N	2.18	1.39	1.33
3	A	203	63Y	CAS-CAR	2.18	1.41	1.37
2	A	201	NDP	O4D-C4D	-2.15	1.40	1.45

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	NDP	N9A-C8A-N7A	-7.28	103.61	113.94
2	A	201	NDP	C4A-N9A-C8A	6.70	112.77	105.74
3	A	202	63Y	C5-C4-N3	-6.54	120.92	126.44
3	A	202	63Y	SAL-C4-N3	6.17	130.98	121.69
2	A	201	NDP	N3A-C2A-N1A	-5.99	119.51	128.58
3	A	203	63Y	C5-C4-N3	-5.88	121.47	126.44
3	A	203	63Y	CAQ-CAS-CAR	5.61	123.49	118.41
3	A	203	63Y	CAH-CAQ-CAS	-5.14	112.78	119.79
3	A	202	63Y	C5-C6-NAC	-4.85	117.00	123.40
3	A	203	63Y	CAG-CAH-CAQ	4.48	125.65	120.48
2	A	201	NDP	C5A-N7A-C8A	4.39	110.35	103.45
3	A	203	63Y	C4-SAL-CAT	-3.94	89.92	94.66
3	A	202	63Y	N3-C2-N1	-3.91	119.48	125.48
2	A	201	NDP	C2A-N1A-C6A	3.69	124.79	118.73
3	A	202	63Y	C5-C4-SAL	-3.48	108.11	112.40
3	A	202	63Y	SAK-CAT-SAL	3.39	125.56	118.82
3	A	203	63Y	NAB-C2-N1	3.25	122.10	117.22
2	A	201	NDP	O4D-C1D-N1N	-3.20	101.99	108.08
2	A	201	NDP	C1D-N1N-C2N	-3.18	115.91	121.14
3	A	203	63Y	NAC-C6-N1	-3.01	112.97	117.03
2	A	201	NDP	C6N-N1N-C2N	2.95	122.48	119.32
3	A	202	63Y	NAC-C6-N1	2.90	120.94	117.03
2	A	201	NDP	C2D-C1D-N1N	2.88	120.38	113.31
3	A	203	63Y	SAL-C4-N3	2.83	125.95	121.69
3	A	203	63Y	NAB-C2-N3	-2.82	112.99	117.22
3	A	202	63Y	C6-C5-CAP	-2.71	130.26	133.85
3	A	203	63Y	C6-C5-CAP	-2.64	130.35	133.85
2	A	201	NDP	O4B-C1B-C2B	-2.43	102.40	106.59
2	A	201	NDP	O4B-C4B-C3B	-2.22	100.74	105.15
2	A	201	NDP	C4B-O4B-C1B	2.22	114.37	109.47
3	A	203	63Y	FAD-CAN-CAG	2.06	123.44	118.65
3	A	202	63Y	NAB-C2-N1	2.03	120.26	117.22
3	A	202	63Y	NAB-C2-N3	2.02	120.25	117.22

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	NDP	C2D-C1D-N1N-C2N
2	A	201	NDP	C3D-C4D-C5D-O5D
2	A	201	NDP	O4D-C4D-C5D-O5D
2	A	201	NDP	O4D-C1D-N1N-C2N
2	A	201	NDP	PA-O3-PN-O1N

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	A	201	NDP	PA-O3-PN-O2N


There are no ring outliers.

2 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	NDP	5	0
3	A	202	63Y	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	186/186 (100%)	0.52	15 (8%)	18 19	6, 13, 33, 48	6 (3%)

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	ALA	4.5
1	A	42	SER	4.4
1	A	103	PRO	4.0
1	A	63	LYS	3.7
1	A	44	GLU	3.5
1	A	61	PRO	3.3
1	A	99	LEU	3.1
1	A	64	ASN	3.0
1	A	101	GLU	2.9
1	A	107	ASN	2.9
1	A	102	GLN	2.8
1	A	45	GLY	2.6
1	A	28	ARG	2.4
1	A	31[A]	PHE	2.3
1	A	43	VAL	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	63Y	A	202	21/22	0.79	0.16	11,14,27,28	21
3	63Y	A	203	21/22	0.95	0.09	7,11,16,21	21
2	NDP	A	201	48/48	0.96	0.08	10,14,18,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 63Y A 202:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 63Y A 203:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP A 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.