



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 11:29 PM UTC

PDB ID : 1HT5 / pdb_00001ht5
Title : THE 2.75 ANGSTROM RESOLUTION MODEL OF OVINE COX-1 COM-
PLEXED WITH METHYL ESTER FLURBIPROFEN
Authors : Selinsky, B.S.; Gupta, K.; Sharkey, C.T.; Loll, P.J.
Deposited on : 2000-12-28
Resolution : 2.75 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	
1	B	551	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	A	1671	-	-	X	-
5	FL2	B	2701	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 9428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROSTAGLANDIN H2 SYNTHASE-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	551	Total 4477	C 2903	N 758	O 788	S 28	0	0	0
1	B	551	Total 4477	C 2903	N 758	O 788	S 28	0	0	0

- Molecule 2 is octyl beta-D-glucopyranoside (CCD ID: BOG) (formula: C₁₄H₂₈O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 20	C 14	O 6	0	0
2	B	1	Total 20	C 14	O 6	0	0
2	B	1	Total 20	C 14	O 6	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	A	1	Total 14	8	1	5	0	0
3	B	1	Total 14	8	1	5	0	0
3	B	1	Total 14	8	1	5	0	0
3	B	1	Total 14	8	1	5	0	0
3	B	1	Total 14	8	1	5	0	0
3	B	1	Total 14	8	1	5	0	0

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


- Molecule 5 is FLURBIPROFEN METHYL ESTER (CCD ID: FL2) (formula: $C_{16}H_{15}FO_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
5	A	1	Total	C	F	O	0	0
			19	16	1	2		
5	B	1	Total	C	F	O	0	0
			19	16	1	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	79	Total 79	O 79	0	0
6	B	71	Total 71	O 71	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	98.89Å 209.45Å 223.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.89 – 2.75 50.89 – 2.75	Depositor EDS
% Data completeness (in resolution range)	88.2 (50.89-2.75) 88.2 (50.89-2.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.31 (at 2.69Å)	Xtrriage
Refinement program	CNS	Depositor
R, R_{free}	0.223 , 0.252 0.215 , 0.248	Depositor DCC
R_{free} test set	5438 reflections (9.41%)	wwPDB-VP
Wilson B-factor (Å ²)	42.7	Xtrriage
Anisotropy	0.082	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9428	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FL2, BOG, NAG, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/4615	0.97	24/6264 (0.4%)
1	B	0.46	0/4615	0.99	17/6264 (0.3%)
All	All	0.45	0/9230	0.98	41/12528 (0.3%)

There are no bond length outliers.

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	224	LEU	N-CA-C	-9.80	97.69	110.33
1	A	224	LEU	N-CA-C	-8.85	98.74	110.24
1	B	287	VAL	N-CA-C	8.24	121.11	113.20
1	A	287	VAL	N-CA-C	7.75	121.08	113.71
1	B	148	TYR	N-CA-C	-7.63	98.19	110.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4477	0	4386	201	0
1	B	4477	0	4386	185	0
2	A	20	0	28	8	0
2	B	40	0	56	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	70	0	65	11	0
3	B	70	0	65	9	0
4	A	43	0	30	5	0
4	B	43	0	30	8	0
5	A	19	0	15	4	0
5	B	19	0	15	7	0
6	A	79	0	0	0	0
6	B	71	0	0	0	0
All	All	9428	0	9076	380	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 380 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:144:ASN:HD21	3:B:2671:NAG:C1	1.80	0.94
1:B:410:ASN:HD21	3:B:2681:NAG:C1	1.81	0.94
1:A:391:MET:HG3	4:A:601:HEM:HAB	1.47	0.92
1:A:163:MET:HE3	1:A:171:LEU:HD21	1.51	0.90
1:B:185:ARG:HH21	1:B:438:ARG:NH1	1.70	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	549/551 (100%)	509 (93%)	36 (7%)	4 (1%)	18 34
1	B	549/551 (100%)	504 (92%)	42 (8%)	3 (0%)	24 43
All	All	1098/1102 (100%)	1013 (92%)	78 (7%)	7 (1%)	21 38

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	514	PRO
1	B	514	PRO
1	A	483	GLY
1	A	270	PRO
1	B	290	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	486/486 (100%)	452 (93%)	34 (7%)	14	26
1	B	486/486 (100%)	444 (91%)	42 (9%)	10	18
All	All	972/972 (100%)	896 (92%)	76 (8%)	11	21

5 of 76 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	300	LEU
1	B	534	LEU
1	B	316	LEU
1	B	460	LEU
1	B	578	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	386	HIS
1	B	400	GLN
1	B	443	HIS
1	A	386	HIS
1	A	375	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FL2	B	2701	-	20,20,20	3.17	14 (70%)	27,27,27	1.30	4 (14%)
3	NAG	A	1681	-	14,14,15	1.70	3 (21%)	17,19,21	0.92	0
3	NAG	B	1672	-	14,14,15	1.51	4 (28%)	17,19,21	1.13	1 (5%)
3	NAG	A	2672	-	14,14,15	1.86	6 (42%)	17,19,21	1.12	1 (5%)
2	BOG	B	2802	-	20,20,20	1.55	5 (25%)	25,25,25	0.96	1 (4%)
3	NAG	A	1661	-	14,14,15	1.35	3 (21%)	17,19,21	1.02	1 (5%)
4	HEM	A	601	1	50,50,50	1.44	5 (10%)	67,82,82	1.11	5 (7%)
2	BOG	A	1802	-	20,20,20	1.58	5 (25%)	25,25,25	0.88	2 (8%)
3	NAG	B	2662	-	14,14,15	1.70	5 (35%)	17,19,21	1.02	1 (5%)
3	NAG	A	1671	-	14,14,15	1.18	1 (7%)	17,19,21	0.97	1 (5%)
2	BOG	B	2801	-	20,20,20	1.73	6 (30%)	25,25,25	1.08	1 (4%)
3	NAG	B	2671	-	14,14,15	1.10	1 (7%)	17,19,21	0.94	1 (5%)
3	NAG	A	1662	-	14,14,15	1.66	5 (35%)	17,19,21	1.00	1 (5%)
5	FL2	A	1701	-	20,20,20	3.15	12 (60%)	27,27,27	1.32	4 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	2681	-	14,14,15	1.57	3 (21%)	17,19,21	1.03	1 (5%)
3	NAG	B	2661	-	14,14,15	1.40	2 (14%)	17,19,21	1.03	1 (5%)
4	HEM	B	601	1	50,50,50	1.43	5 (10%)	67,82,82	1.15	7 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	FL2	B	2701	-	-	3/14/14/14	0/2/2/2
3	NAG	A	1681	-	-	4/6/23/26	0/1/1/1
3	NAG	B	1672	-	-	4/6/23/26	0/1/1/1
3	NAG	A	2672	-	-	4/6/23/26	0/1/1/1
2	BOG	B	2802	-	-	4/11/31/31	0/1/1/1
3	NAG	A	1661	-	-	2/6/23/26	0/1/1/1
4	HEM	A	601	1	-	4/14/54/54	-
2	BOG	A	1802	-	-	4/11/31/31	0/1/1/1
3	NAG	B	2662	-	-	2/6/23/26	0/1/1/1
3	NAG	A	1671	-	-	4/6/23/26	0/1/1/1
2	BOG	B	2801	-	-	1/11/31/31	0/1/1/1
3	NAG	B	2671	-	-	2/6/23/26	0/1/1/1
3	NAG	A	1662	-	-	2/6/23/26	0/1/1/1
5	FL2	A	1701	-	-	3/14/14/14	0/2/2/2
3	NAG	B	2681	-	-	4/6/23/26	0/1/1/1
3	NAG	B	2661	-	-	0/6/23/26	0/1/1/1
4	HEM	B	601	1	-	7/14/54/54	-

The worst 5 of 85 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1701	FL2	C7-C6	5.83	1.48	1.40
5	B	2701	FL2	C7-C6	5.73	1.48	1.40
5	A	1701	FL2	C8-C9	5.34	1.47	1.39
5	B	2701	FL2	C8-C9	5.10	1.47	1.39
5	A	1701	FL2	C1-C2	4.74	1.48	1.39

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2801	BOG	C1'-O1-C1	4.44	121.26	113.68
2	B	2802	BOG	C1'-O1-C1	3.79	120.15	113.68
4	A	601	HEM	C2B-C1B-NB	3.58	113.96	109.84
4	B	601	HEM	C2B-C1B-NB	3.39	113.73	109.84
5	B	2701	FL2	C16-O1-C14	2.79	122.25	115.92

There are no chirality outliers.

5 of 54 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1681	NAG	C8-C7-N2-C2
3	A	1681	NAG	O7-C7-N2-C2
4	A	601	HEM	C2C-C3C-CAC-CBC
4	B	601	HEM	C2B-C3B-CAB-CBB
4	B	601	HEM	C4B-C3B-CAB-CBB



There are no ring outliers.


13 monomers are involved in 56 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	2701	FL2	7	0
3	A	1681	NAG	3	0
3	B	1672	NAG	2	0
2	B	2802	BOG	4	0
3	A	1661	NAG	1	0
4	A	601	HEM	5	0
2	A	1802	BOG	8	0
3	A	1671	NAG	7	0
2	B	2801	BOG	2	0
3	B	2671	NAG	5	0
5	A	1701	FL2	4	0
3	B	2681	NAG	2	0
4	B	601	HEM	8	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/551 (100%)	0.18	14 (2%) 58 56	22, 37, 53, 64	0
1	B	551/551 (100%)	0.15	14 (2%) 58 56	23, 37, 53, 64	0
All	All	1102/1102 (100%)	0.17	28 (2%) 58 56	22, 37, 53, 64	0

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	294	LEU	3.6
1	A	33	VAL	3.4
1	A	107	PHE	3.4
1	A	581	HIS	3.3
1	B	288	GLY	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	BOG	B	2801	20/20	0.40	0.31	37,63,68,68	0
3	NAG	B	2662	14/15	0.55	0.28	63,84,96,96	0
3	NAG	A	1662	14/15	0.57	0.27	63,84,96,96	0
3	NAG	A	1681	14/15	0.76	0.19	31,36,42,42	0
3	NAG	A	2672	14/15	0.78	0.20	32,39,65,76	0
3	NAG	B	2681	14/15	0.78	0.18	31,36,42,42	0
2	BOG	A	1802	20/20	0.79	0.19	37,54,64,67	0
2	BOG	B	2802	20/20	0.79	0.19	37,54,64,67	0
3	NAG	B	1672	14/15	0.83	0.18	32,39,65,76	0
3	NAG	A	1661	14/15	0.84	0.17	50,60,64,78	0
3	NAG	B	2661	14/15	0.84	0.17	50,60,64,78	0
3	NAG	A	1671	14/15	0.91	0.10	6,16,23,25	0
5	FL2	A	1701	19/19	0.91	0.16	50,51,56,59	0
5	FL2	B	2701	19/19	0.92	0.14	50,52,56,59	0
3	NAG	B	2671	14/15	0.94	0.08	6,16,23,25	0
4	HEM	A	601	43/43	0.95	0.12	36,41,53,61	0
4	HEM	B	601	43/43	0.96	0.11	36,40,55,65	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around BOG A 1802:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BOG B 2802:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around FL2 A 1701:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FL2 B 2701:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







6.5 Other polymers [i](#)

There are no such residues in this entry.