



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:55 AM UTC

PDB ID : 6HUH / pdb_00006huh
Title : CRYSTAL STRUCTURE OF OXA-427 class D BETA-LACTAMASE
Authors : Zavala, A.; Retailleau, P.; Bogaerts, P.; Glupczynski, Y.; Naas, T.; Iorga, B.
Deposited on : 2018-10-08
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	5% 70% 17% • 12%
1	B	274	6% 70% 16% • 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	KCX	A	54[B]	-	X	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3971 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	242	Total	C	N	O	S	0	1	0
			1912	1228	328	351	5			
1	B	242	Total	C	N	O	S	0	1	0
			1912	1228	328	351	5			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	265	VAL	-	expression tag	UNP A0A1Y6M6D7
A	266	GLU	-	expression tag	UNP A0A1Y6M6D7
A	267	HIS	-	expression tag	UNP A0A1Y6M6D7
A	268	HIS	-	expression tag	UNP A0A1Y6M6D7
A	269	HIS	-	expression tag	UNP A0A1Y6M6D7
A	270	HIS	-	expression tag	UNP A0A1Y6M6D7
A	271	HIS	-	expression tag	UNP A0A1Y6M6D7
A	272	HIS	-	expression tag	UNP A0A1Y6M6D7
A	273	HIS	-	expression tag	UNP A0A1Y6M6D7
A	274	HIS	-	expression tag	UNP A0A1Y6M6D7
B	265	VAL	-	expression tag	UNP A0A1Y6M6D7
B	266	GLU	-	expression tag	UNP A0A1Y6M6D7
B	267	HIS	-	expression tag	UNP A0A1Y6M6D7
B	268	HIS	-	expression tag	UNP A0A1Y6M6D7
B	269	HIS	-	expression tag	UNP A0A1Y6M6D7
B	270	HIS	-	expression tag	UNP A0A1Y6M6D7
B	271	HIS	-	expression tag	UNP A0A1Y6M6D7
B	272	HIS	-	expression tag	UNP A0A1Y6M6D7
B	273	HIS	-	expression tag	UNP A0A1Y6M6D7
B	274	HIS	-	expression tag	UNP A0A1Y6M6D7

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			3	2	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	69	Total	O	0	0
			69	69		
3	B	65	Total	O	0	0
			65	65		

- Molecule 1: Beta-lactamase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	143.79Å 42.58Å 99.71Å 90.00° 114.37° 90.00°	Depositor
Resolution (Å)	68.10 – 2.78 68.10 – 2.78	Depositor EDS
% Data completeness (in resolution range)	94.2 (68.10-2.78) 94.2 (68.10-2.78)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.26 (at 2.77Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.205 , 0.269 0.213 , 0.279	Depositor DCC
R_{free} test set	681 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.433	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 78.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3971	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 19.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.8380e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: KCX, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/1948	1.35	7/2644 (0.3%)
1	B	0.84	0/1948	1.39	14/2644 (0.5%)
All	All	0.84	0/3896	1.37	21/5288 (0.4%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	PHE	CA-CB-CG	-6.60	107.20	113.80
1	B	219	PHE	CA-CB-CG	-6.45	107.35	113.80
1	B	211	ASP	CA-C-N	6.38	129.69	120.38
1	B	211	ASP	C-N-CA	6.38	129.69	120.38
1	B	137	GLY	CA-C-N	6.30	129.58	120.38
1	B	137	GLY	C-N-CA	6.30	129.58	120.38
1	A	239	LYS	N-CA-C	-5.63	102.95	109.93
1	B	205	LEU	CA-C-N	5.45	131.95	121.54
1	B	205	LEU	C-N-CA	5.45	131.95	121.54
1	B	190	ASP	CA-CB-CG	5.44	118.04	112.60
1	B	118	PHE	CA-CB-CG	5.36	119.16	113.80
1	B	249	ALA	CA-C-N	5.30	127.38	120.28
1	B	249	ALA	C-N-CA	5.30	127.38	120.28
1	B	96	ARG	CA-C-N	5.24	127.25	120.44
1	B	96	ARG	C-N-CA	5.24	127.25	120.44
1	A	249	ALA	CA-C-N	5.21	127.26	120.28
1	A	249	ALA	C-N-CA	5.21	127.26	120.28
1	A	96	ARG	CA-C-N	5.11	127.08	120.44
1	A	96	ARG	C-N-CA	5.11	127.08	120.44
1	B	227	ASP	CA-CB-CG	5.06	117.66	112.60
1	A	227	ASP	CA-CB-CG	5.00	117.60	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1912	0	1881	16	0
1	B	1912	0	1881	18	0
2	A	8	0	0	0	0
2	B	5	0	0	0	0
3	A	69	0	0	0	0
3	B	65	0	0	0	0
All	All	3971	0	3762	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:185:ARG:HH22	1:B:193:GLN:HG2	1.43	0.81
1:B:205:LEU:HB3	1:B:210:LEU:HB3	1.77	0.65
1:B:220:VAL:HG12	1:B:233:VAL:HG13	1.82	0.61
1:A:124:ARG:HD3	1:A:171:PRO:HG3	1.84	0.59
1:A:220:VAL:HG12	1:A:233:VAL:HG13	1.84	0.59
1:A:198:THR:HA	1:A:218:TRP:O	2.04	0.58
1:B:198:THR:HB	1:B:246:SER:HB2	1.86	0.58
1:B:124:ARG:HD3	1:B:171:PRO:HG3	1.85	0.57
1:A:59:LEU:HD11	1:A:165:LEU:HD22	1.87	0.56
1:B:198:THR:HA	1:B:218:TRP:O	2.06	0.55
1:B:59:LEU:HD11	1:B:165:LEU:HD22	1.89	0.53
1:A:203:PRO:HG3	1:A:216:ILE:HD11	1.94	0.48
1:A:253:VAL:O	1:A:257:LEU:HB2	2.16	0.46
1:B:207:ASP:HB3	1:B:210:LEU:HG	1.97	0.45
1:A:198:THR:HB	1:A:246:SER:HB2	1.96	0.45
1:A:58:ALA:HB2	1:A:98:MET:HE2	1.98	0.45
1:A:70:GLU:HG3	1:A:183:ILE:HG23	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:217:GLY:O	1:A:235:THR:HA	2.16	0.44
1:B:205:LEU:HD13	1:B:212:ARG:HA	2.00	0.43
1:B:70:GLU:HG3	1:B:183:ILE:HG23	2.00	0.43
1:A:59:LEU:HD22	1:A:172:VAL:HG21	2.01	0.43
1:A:62:TYR:HB2	1:A:176:THR:HG23	2.01	0.43
1:B:62:TYR:HB2	1:B:176:THR:HG23	2.01	0.42
1:B:27:MET:HG2	1:B:234:HIS:CE1	2.54	0.42
1:B:210:LEU:HD13	1:B:211:ASP:H	1.85	0.41
1:B:30:ALA:HB3	1:B:231:ILE:HB	2.02	0.41
1:B:192:TRP:CH2	1:B:264:LEU:HD23	2.56	0.41
1:A:187:PRO:HA	1:B:71:GLN:OE1	2.20	0.41
1:A:243:GLN:HB2	1:A:248:ARG:NH2	2.35	0.41
1:A:54[B]:KCX:HZ	1:A:103:ILE:HA	1.86	0.41
1:B:192:TRP:HH2	1:B:264:LEU:HD23	1.86	0.41
1:A:58:ALA:O	1:A:62:TYR:HD2	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/274 (87%)	219 (92%)	17 (7%)	3 (1%)	9	28
1	B	239/274 (87%)	221 (92%)	17 (7%)	1 (0%)	30	57
All	All	478/548 (87%)	440 (92%)	34 (7%)	4 (1%)	16	40

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	204	LYS
1	B	208	GLY
1	A	205	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	208	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/226 (88%)	184 (92%)	16 (8%)	11	31
1	B	200/226 (88%)	184 (92%)	16 (8%)	11	31
All	All	400/452 (88%)	368 (92%)	32 (8%)	11	31

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	CYS
1	A	87	GLU
1	A	90	GLN
1	A	144	GLN
1	A	147	LEU
1	A	154	SER
1	A	162	LEU
1	A	165	LEU
1	A	166	VAL
1	A	206	LEU
1	A	210	LEU
1	A	215	GLN
1	A	226	GLN
1	A	227	ASP
1	A	238	GLN
1	A	262	LYS
1	B	26	CYS
1	B	27	MET
1	B	90	GLN
1	B	139	HIS
1	B	144	GLN
1	B	147	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	154	SER
1	B	162	LEU
1	B	165	LEU
1	B	166	VAL
1	B	204	LYS
1	B	206	LEU
1	B	210	LEU
1	B	227	ASP
1	B	248	ARG
1	B	262	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	120	GLN
1	A	139	HIS
1	A	193	GLN
1	A	215	GLN
1	A	228	GLN
1	A	238	GLN
1	B	90	GLN
1	B	120	GLN
1	B	139	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	KCX	A	54[A]	-	10,11,12	0.90	0	6,12,14	1.07	1 (16%)
1	KCX	B	54[B]	-	10,8,12	4.25	1 (10%)	6,8,14	13.94	2 (33%)
1	KCX	A	54[B]	-	10,8,12	6.88	1 (10%)	6,8,14	9.40	2 (33%)
1	KCX	B	54[A]	-	10,11,12	0.82	0	6,12,14	1.68	1 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	54[A]	-	-	3/9/10/12	-
1	KCX	B	54[B]	-	-	4/9/7/12	-
1	KCX	A	54[B]	-	-	7/9/7/12	-
1	KCX	B	54[A]	-	-	1/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	54[B]	KCX	CX-NZ	-21.67	0.96	1.35
1	B	54[B]	KCX	CX-NZ	-13.30	1.11	1.35

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	54[B]	KCX	OQ1-CX-NZ	-30.19	79.06	124.92
1	A	54[B]	KCX	OQ1-CX-NZ	22.68	159.38	124.92
1	B	54[B]	KCX	CE-NZ-CX	15.86	148.89	121.98
1	A	54[B]	KCX	CE-NZ-CX	3.54	127.99	121.98
1	B	54[A]	KCX	OQ1-CX-NZ	-3.30	119.90	124.92
1	A	54[A]	KCX	CE-NZ-CX	2.31	125.89	121.98

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	54[A]	KCX	C-CA-CB-CG
1	A	54[B]	KCX	C-CA-CB-CG
1	A	54[B]	KCX	CD-CE-NZ-CX
1	A	54[B]	KCX	OQ1-CX-NZ-CE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	A	54[B]	KCX	OQ2-CX-NZ-CE
1	B	54[B]	KCX	CD-CE-NZ-CX
1	B	54[B]	KCX	OQ1-CX-NZ-CE
1	B	54[B]	KCX	OQ2-CX-NZ-CE
1	A	54[B]	KCX	CG-CD-CE-NZ
1	A	54[B]	KCX	CA-CB-CG-CD
1	B	54[A]	KCX	C-CA-CB-CG
1	A	54[B]	KCX	N-CA-CB-CG
1	A	54[A]	KCX	CE-CD-CG-CB
1	B	54[B]	KCX	C-CA-CB-CG
1	A	54[A]	KCX	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	54[B]	KCX	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	301	-	4,4,4	0.30	0	6,6,6	0.17	0
2	SO4	B	301	-	4,4,4	0.33	0	6,6,6	0.07	0
2	SO4	A	302	-	0,2,4	-	-	0,1,6	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	241/274 (87%)	0.55	14 (5%) 29 24	23, 42, 84, 127	0
1	B	241/274 (87%)	0.60	16 (6%) 24 19	27, 49, 92, 130	0
All	All	482/548 (87%)	0.58	30 (6%) 26 21	23, 45, 92, 130	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	206	LEU	5.6
1	A	205	LEU	4.8
1	A	211	ASP	4.3
1	A	210	LEU	4.3
1	B	206	LEU	3.6
1	B	207	ASP	3.4
1	B	213	ASP	3.4
1	B	82	PRO	3.2
1	B	208	GLY	3.1
1	A	265	VAL	3.1
1	B	210	LEU	3.0
1	A	213	ASP	3.0
1	A	208	GLY	2.9
1	B	205	LEU	2.8
1	A	204	LYS	2.7
1	B	265	VAL	2.5
1	B	256	ALA	2.5
1	A	200	THR	2.4
1	A	209	SER	2.4
1	B	211	ASP	2.3
1	B	227	ASP	2.3
1	A	140	ASP	2.2
1	B	209	SER	2.2
1	B	99	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	139	HIS	2.2
1	A	138	LYS	2.2
1	B	238	GLN	2.2
1	B	212	ARG	2.1
1	B	241	GLY	2.1
1	A	160	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	B	54[A]	12/13	0.94	0.10	21,26,31,34	8
1	KCX	B	54[B]	9/13	0.94	0.10	28,28,31,34	5
1	KCX	A	54[A]	12/13	0.97	0.06	15,23,28,29	8
1	KCX	A	54[B]	9/13	0.97	0.06	26,26,28,29	5

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	302	3/5	0.90	0.14	45,45,46,46	3
2	SO4	B	301	5/5	0.90	0.11	87,88,90,90	0
2	SO4	A	301	5/5	0.93	0.14	88,88,88,90	0

6.5 Other polymers [i](#)

There are no such residues in this entry.