



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:59 PM UTC

PDB ID : 8HU4 / pdb_00008hu4
Title : Limosilactobacillus reuteri N1 GtfB
Authors : Dong, J.J.; Bai, Y.X.
Deposited on : 2022-12-22
Resolution : 2.76 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1009 (2.76-2.76)
Clashscore	190562	1044 (2.76-2.76)
Ramachandran outliers	187476	1024 (2.76-2.76)
Sidechain outliers	187428	1024 (2.76-2.76)
RSRZ outliers	180081	1009 (2.76-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	833	14% 95%
1	B	833	14% 94% 6%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13042 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dextransucrase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	827	Total	C	N	O	S	0	0	0
			6457	4025	1095	1315	22			
1	B	827	Total	C	N	O	S	0	0	0
			6457	4025	1095	1315	22			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	25	MET	-	initiating methionine	UNP A0A848PDI7
B	25	MET	-	initiating methionine	UNP A0A848PDI7

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		

- Molecule 3 is CITRIC ACID (CCD ID: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		
4	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	44	Total 44	O 44	0	0
5	B	42	Total 42	O 42	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: dextranucrase



- Molecule 1: dextranucrase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	204.42Å 100.73Å 106.89Å 90.00° 108.14° 90.00°	Depositor
Resolution (Å)	47.21 – 2.76 47.21 – 2.76	Depositor EDS
% Data completeness (in resolution range)	99.6 (47.21-2.76) 99.6 (47.21-2.76)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.40 (at 2.77Å)	Xtriage
Refinement program	REFMAC 5	Depositor
R, R_{free}	0.248 , 0.266 0.252 , 0.272	Depositor DCC
R_{free} test set	2666 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	13042	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, NA, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	0/6597	1.44	5/8984 (0.1%)
1	B	1.03	0/6597	1.44	5/8984 (0.1%)
All	All	1.03	0/13194	1.44	10/17968 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	820	GLY	CA-C-O	-6.47	118.02	122.22
1	A	820	GLY	CA-C-O	-6.37	118.08	122.22
1	A	506	GLY	CA-C-O	-6.11	118.05	122.45
1	B	506	GLY	CA-C-O	-6.09	118.07	122.45
1	B	527	GLY	CA-C-O	-6.08	118.27	122.22
1	A	527	GLY	CA-C-O	-5.67	118.36	122.45
1	A	52	THR	CA-C-N	5.52	125.34	120.10
1	A	52	THR	C-N-CA	5.52	125.34	120.10
1	B	52	THR	CA-C-N	5.45	125.28	120.10
1	B	52	THR	C-N-CA	5.45	125.28	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6457	0	6094	15	0
1	B	6457	0	6094	19	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	13	0	5	0	0
3	B	13	0	5	0	0
4	A	7	0	10	0	0
4	B	7	0	10	0	0
5	A	44	0	0	0	0
5	B	42	0	0	0	0
All	All	13042	0	12218	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:835:LEU:HD11	1:A:847:LEU:HD21	1.69	0.73
1:B:835:LEU:HD11	1:B:847:LEU:HD21	1.70	0.72
1:B:212:ASN:O	1:B:740:ARG:NH2	2.35	0.59
1:B:243:TYR:CE1	1:B:685:LEU:HD22	2.44	0.53
1:A:243:TYR:CE1	1:A:685:LEU:HD22	2.45	0.52
1:B:851:ILE:HG22	1:B:851:ILE:O	2.10	0.51
1:B:49:LEU:HD11	1:B:835:LEU:HD13	1.93	0.49
1:A:49:LEU:HD11	1:A:835:LEU:HD13	1.94	0.49
1:A:200:ILE:HA	1:A:308:MET:HE2	1.96	0.48
1:A:355:THR:O	1:A:427:LYS:NZ	2.47	0.47
1:B:355:THR:O	1:B:427:LYS:NZ	2.45	0.47
1:B:57:PRO:O	1:B:70:THR:OG1	2.29	0.47
1:A:57:PRO:O	1:A:70:THR:OG1	2.31	0.46
1:A:77:PRO:HG2	1:A:80:MET:HG3	1.98	0.45
1:B:416:VAL:HB	1:B:417:PRO:HD3	1.98	0.45
1:B:110:ALA:O	1:B:114:LYS:HB2	2.17	0.44
1:A:416:VAL:HB	1:A:417:PRO:HD3	1.99	0.43
1:A:76:ARG:HB3	1:A:80:MET:HE2	2.00	0.43
1:A:80:MET:SD	1:A:136:ILE:HG21	2.59	0.43
1:A:851:ILE:O	1:A:851:ILE:HG22	2.19	0.43
1:B:76:ARG:HB3	1:B:80:MET:HE2	2.02	0.42
1:B:368:ASN:HB2	1:B:371:GLN:HB2	2.02	0.41
1:A:62:GLN:HE22	1:A:141:VAL:HG21	1.85	0.41
1:A:473:MET:HE3	1:A:483:SER:HB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:536:ARG:NH2	1:A:538:LEU:HD12	2.35	0.41
1:B:243:TYR:CE1	1:B:247:MET:HG3	2.54	0.41
1:B:494:ASN:HD22	1:B:528:ARG:HD2	1.86	0.41
1:B:626:PHE:CG	1:B:662:ALA:HB2	2.56	0.41
1:B:266:ILE:HD12	1:B:270:VAL:HG21	2.03	0.41
1:B:77:PRO:HG2	1:B:80:MET:HG3	2.01	0.41
1:B:473:MET:HE3	1:B:483:SER:HB2	2.02	0.41
1:A:243:TYR:CE1	1:A:247:MET:HG3	2.56	0.40
1:B:236:TRP:CZ2	1:B:682:ARG:HG2	2.56	0.40
1:B:376:ILE:O	1:B:380:ILE:HG12	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	825/833 (99%)	783 (95%)	41 (5%)	1 (0%)	48	71
1	B	825/833 (99%)	783 (95%)	41 (5%)	1 (0%)	48	71
All	All	1650/1666 (99%)	1566 (95%)	82 (5%)	2 (0%)	48	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	200	ILE
1	B	200	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	695/700 (99%)	686 (99%)	9 (1%)	61	76
1	B	695/700 (99%)	687 (99%)	8 (1%)	63	78
All	All	1390/1400 (99%)	1373 (99%)	17 (1%)	63	78

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	VAL
1	A	260	ILE
1	A	287	ASN
1	A	295	LEU
1	A	297	TYR
1	A	392	TYR
1	A	679	VAL
1	A	680	THR
1	A	726	LEU
1	B	66	THR
1	B	90	VAL
1	B	260	ILE
1	B	287	ASN
1	B	297	TYR
1	B	392	TYR
1	B	679	VAL
1	B	680	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	98	ASN
1	A	151	ASN
1	A	334	ASN
1	A	335	ASN
1	A	351	GLN
1	A	410	GLN
1	A	419	GLN
1	A	479	ASN
1	A	638	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	674	ASN
1	A	765	ASN
1	B	334	ASN
1	B	335	ASN
1	B	419	GLN
1	B	494	ASN
1	B	608	ASN
1	B	638	ASN
1	B	765	ASN
1	B	840	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	903	-	6,6,6	0.14	0	5,5,5	0.08	0
3	CIT	A	902	-	12,12,12	1.18	1 (8%)	17,17,17	1.11	2 (11%)
3	CIT	B	902	-	12,12,12	1.20	1 (8%)	17,17,17	1.15	2 (11%)
4	PEG	B	903	-	6,6,6	0.13	0	5,5,5	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	903	-	-	2/4/4/4	-
3	CIT	A	902	-	-	0/16/16/16	-
3	CIT	B	902	-	-	3/16/16/16	-
4	PEG	B	903	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	902	CIT	C3-C6	2.34	1.55	1.53
3	A	902	CIT	C3-C6	2.18	1.55	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	902	CIT	O5-C6-C3	-2.47	117.31	122.09
3	A	902	CIT	O5-C6-C3	-2.45	117.36	122.09
3	B	902	CIT	O6-C6-C3	2.37	117.68	113.14
3	A	902	CIT	O6-C6-C3	2.23	117.42	113.14

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	902	CIT	C1-C2-C3-O7
3	B	902	CIT	C1-C2-C3-C4
3	B	902	CIT	C1-C2-C3-C6
4	B	903	PEG	O1-C1-C2-O2
4	A	903	PEG	O2-C3-C4-O4
4	A	903	PEG	C4-C3-O2-C2
4	B	903	PEG	C4-C3-O2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	827/833 (99%)	0.96	116 (14%) 6 5	19, 29, 63, 72	0
1	B	827/833 (99%)	0.95	114 (13%) 6 5	20, 29, 65, 76	0
All	All	1654/1666 (99%)	0.95	230 (13%) 6 5	19, 29, 65, 76	0

All (230) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	603	VAL	9.2
1	A	837	SER	7.5
1	B	26	GLY	6.8
1	A	838	SER	6.6
1	B	603	VAL	6.0
1	A	385	PRO	5.3
1	B	388	TYR	5.2
1	B	385	PRO	5.0
1	A	124	ASN	4.8
1	B	390	ALA	4.7
1	A	388	TYR	4.7
1	B	387	ALA	4.6
1	B	838	SER	4.3
1	A	387	ALA	4.3
1	A	386	GLY	4.3
1	B	386	GLY	4.3
1	B	111	GLY	4.3
1	B	839	ASP	4.2
1	A	28	GLY	4.2
1	B	141	VAL	4.2
1	B	69	PRO	4.0
1	B	849	SER	3.9
1	A	141	VAL	3.9
1	B	107	GLY	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	72	VAL	3.8
1	A	384	HIS	3.8
1	B	145	SER	3.8
1	B	100	GLY	3.8
1	A	842	SER	3.7
1	B	73	ALA	3.7
1	B	852	SER	3.7
1	A	38	ASP	3.7
1	A	68	TYR	3.7
1	A	26	GLY	3.7
1	B	601	ALA	3.6
1	A	835	LEU	3.6
1	A	123	ILE	3.6
1	B	747	ASN	3.6
1	B	550	ALA	3.6
1	B	112	SER	3.6
1	A	745	ASP	3.5
1	A	605	ALA	3.5
1	B	124	ASN	3.5
1	A	334	ASN	3.5
1	B	67	TRP	3.4
1	B	382	LYS	3.4
1	A	531	ALA	3.4
1	B	28	GLY	3.4
1	B	287	ASN	3.4
1	B	27	PRO	3.3
1	A	747	ASN	3.3
1	B	605	ALA	3.3
1	B	35	PHE	3.3
1	B	837	SER	3.3
1	A	841	LYS	3.3
1	A	113	VAL	3.3
1	A	840	ASN	3.3
1	A	839	ASP	3.3
1	B	74	ASP	3.3
1	B	836	ARG	3.3
1	B	61	SER	3.2
1	A	550	ALA	3.2
1	B	392	TYR	3.2
1	A	601	ALA	3.2
1	A	172	GLU	3.1
1	A	27	PRO	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	390	ALA	3.1
1	A	392	TYR	3.0
1	B	65	LYS	3.0
1	A	213	GLY	3.0
1	A	378	GLY	3.0
1	B	58	TYR	3.0
1	A	120	THR	3.0
1	B	123	ILE	3.0
1	B	389	LYS	3.0
1	B	136	ILE	3.0
1	A	122	SER	3.0
1	A	66	THR	3.0
1	B	110	ALA	3.0
1	B	835	LEU	2.9
1	A	843	PHE	2.9
1	B	113	VAL	2.9
1	A	850	GLU	2.9
1	B	384	HIS	2.9
1	B	842	SER	2.9
1	A	287	ASN	2.9
1	B	105	ASN	2.9
1	B	115	ASP	2.9
1	B	391	GLU	2.9
1	A	59	GLY	2.9
1	A	849	SER	2.8
1	A	109	THR	2.8
1	A	602	THR	2.8
1	A	391	GLU	2.8
1	A	62	GLN	2.8
1	B	131	ASN	2.8
1	B	108	LEU	2.8
1	A	110	ALA	2.8
1	B	143	ALA	2.8
1	B	175	TYR	2.8
1	B	597	THR	2.8
1	A	140	ILE	2.8
1	A	184	ASP	2.8
1	B	845	THR	2.8
1	A	389	LYS	2.8
1	B	40	SER	2.8
1	B	604	SER	2.8
1	B	38	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	308	MET	2.8
1	B	109	THR	2.7
1	A	32	ASN	2.7
1	B	469	GLY	2.7
1	A	142	ALA	2.7
1	B	97	VAL	2.7
1	B	101	TYR	2.7
1	A	174	GLY	2.7
1	A	836	ARG	2.7
1	A	393	ALA	2.7
1	B	62	GLN	2.6
1	A	111	GLY	2.6
1	B	491	ALA	2.6
1	B	99	HIS	2.6
1	B	841	LYS	2.6
1	B	30	TRP	2.6
1	B	146	THR	2.6
1	A	106	TYR	2.6
1	B	68	TYR	2.6
1	A	201	ASN	2.6
1	A	598	THR	2.6
1	A	674	ASN	2.6
1	A	604	SER	2.5
1	A	851	ILE	2.5
1	A	590	VAL	2.5
1	A	29	THR	2.5
1	A	112	SER	2.5
1	A	211	ASP	2.5
1	B	172	GLU	2.5
1	A	396	ALA	2.5
1	A	374	ASN	2.5
1	B	334	ASN	2.5
1	B	116	LEU	2.5
1	B	31	GLU	2.5
1	B	840	ASN	2.4
1	B	667	GLN	2.4
1	A	369	HIS	2.4
1	B	745	ASP	2.4
1	A	73	ALA	2.4
1	A	558	THR	2.4
1	A	852	SER	2.4
1	A	65	LYS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	382	LYS	2.4
1	B	134	TYR	2.4
1	A	597	THR	2.4
1	A	634	ILE	2.4
1	A	117	SER	2.4
1	A	380	ILE	2.3
1	A	303	SER	2.3
1	A	37	GLN	2.3
1	B	672	ARG	2.3
1	A	150	ALA	2.3
1	B	851	ILE	2.3
1	B	66	THR	2.3
1	A	94	GLN	2.3
1	A	487	GLY	2.3
1	A	36	ALA	2.3
1	A	752	ALA	2.3
1	B	93	ILE	2.3
1	B	140	ILE	2.3
1	B	81	TYR	2.3
1	B	120	THR	2.3
1	A	114	LYS	2.3
1	A	158	THR	2.3
1	A	521	THR	2.3
1	A	58	TYR	2.2
1	A	535	TYR	2.2
1	B	98	ASN	2.2
1	B	161	PRO	2.2
1	B	85	SER	2.2
1	A	45	ILE	2.2
1	B	843	PHE	2.2
1	A	833	ALA	2.2
1	A	127	GLU	2.2
1	B	846	LEU	2.2
1	B	29	THR	2.2
1	A	301	TYR	2.2
1	B	106	TYR	2.2
1	B	104	SER	2.2
1	A	33	MET	2.2
1	B	596	VAL	2.2
1	A	615	ALA	2.1
1	A	131	ASN	2.1
1	B	844	ASN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	134	TYR	2.1
1	B	673	TYR	2.1
1	A	93	ILE	2.1
1	A	189	ALA	2.1
1	B	600	ALA	2.1
1	A	299	GLU	2.1
1	A	398	GLN	2.1
1	B	125	LEU	2.1
1	A	107	GLY	2.1
1	A	527	GLY	2.1
1	B	45	ILE	2.1
1	B	396	ALA	2.1
1	B	752	ALA	2.1
1	A	102	GLU	2.1
1	B	71	THR	2.1
1	B	59	GLY	2.1
1	B	301	TYR	2.1
1	B	834	TYR	2.1
1	A	90	VAL	2.1
1	A	97	VAL	2.1
1	A	121	ALA	2.1
1	B	142	ALA	2.1
1	B	531	ALA	2.1
1	B	144	LYS	2.1
1	B	33	MET	2.1
1	A	206	ASN	2.1
1	B	599	ASN	2.1
1	A	100	GLY	2.1
1	A	35	PHE	2.1
1	A	383	ASP	2.0
1	B	563	ASP	2.0
1	B	607	SER	2.0
1	A	67	TRP	2.0
1	A	116	LEU	2.0
1	B	549	ASN	2.0
1	A	528	ARG	2.0
1	B	634	ILE	2.0
1	A	834	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CIT	B	902	13/13	0.75	0.18	46,46,46,46	0
3	CIT	A	902	13/13	0.77	0.19	47,48,48,48	0
4	PEG	B	903	7/7	0.80	0.16	29,29,29,29	0
4	PEG	A	903	7/7	0.82	0.16	36,36,36,36	0
2	NA	B	901	1/1	0.92	0.10	22,22,22,22	0
2	NA	A	901	1/1	0.98	0.04	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.