



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:26 PM UTC

PDB ID : 4HVM / pdb_00004hvm
Title : Crystal structure of tallysomycin biosynthesis protein TlmII
Authors : Chang, C.; Bigelow, L.; Bearden, J.; Babnigg, G.; Bingman, C.A.; Yennamalli, R.; Lohman, J.; Ma, M.; Shen, B.; Phillips Jr., G.N.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG); Enzyme Discovery for Natural Product Biosynthesis (NatPro)
Deposited on : 2012-11-06
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	493	3% 73% 7% 20%
1	B	493	5% 75% 8% 16%
1	C	493	4% 71% 11% 19%
1	D	493	6% 78% 7% 15%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	D	501	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12790 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TlmII.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	393	Total	C	N	O	S	Se	0	0	0
			3098	1954	575	561	1	7			
1	B	414	Total	C	N	O	S	Se	0	0	0
			3212	2017	594	592	2	7			
1	C	401	Total	C	N	O	S	Se	0	0	0
			3137	1980	580	568	2	7			
1	D	419	Total	C	N	O	S	Se	0	0	0
			3258	2046	603	600	2	7			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A4KUC0
A	-1	ASN	-	expression tag	UNP A4KUC0
A	0	ALA	-	expression tag	UNP A4KUC0
B	-2	SER	-	expression tag	UNP A4KUC0
B	-1	ASN	-	expression tag	UNP A4KUC0
B	0	ALA	-	expression tag	UNP A4KUC0
C	-2	SER	-	expression tag	UNP A4KUC0
C	-1	ASN	-	expression tag	UNP A4KUC0
C	0	ALA	-	expression tag	UNP A4KUC0
D	-2	SER	-	expression tag	UNP A4KUC0
D	-1	ASN	-	expression tag	UNP A4KUC0
D	0	ALA	-	expression tag	UNP A4KUC0

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	16	Total	O	0	0
			16	16		
3	B	16	Total	O	0	0
			16	16		
3	C	18	Total	O	0	0
			18	18		
3	D	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TlmII



• Molecule 1: TlmII



• Molecule 1: TlmII



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	123.80Å 128.01Å 129.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.71 – 2.70 44.71 – 2.70	Depositor EDS
% Data completeness (in resolution range)	77.6 (44.71-2.70) 90.7 (44.71-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.21 (at 2.69Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.182 , 0.232 0.195 , 0.246	Depositor DCC
R_{free} test set	2892 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	28.7	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.020 for -h,l,k 0.017 for -l,-k,-h 0.019 for k,h,-l 0.004 for k,l,h 0.004 for l,h,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12790	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.29	0/3161	0.70	0/4297
1	B	0.30	0/3277	0.74	2/4459 (0.0%)
1	C	0.29	0/3202	0.72	0/4357
1	D	0.30	0/3326	0.72	0/4528
All	All	0.29	0/12966	0.72	2/17641 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	229	GLU	CA-C-N	5.11	125.14	119.32
1	B	229	GLU	C-N-CA	5.11	125.14	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	3033	18	0
1	B	3212	0	3134	23	0
1	C	3137	0	3069	26	0
1	D	3258	0	3170	18	0
2	A	10	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	5	0	0	0	0
2	D	5	0	0	2	0
3	A	16	0	0	0	0
3	B	16	0	0	1	0
3	C	18	0	0	0	0
3	D	15	0	0	0	0
All	All	12790	0	12406	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:415:ASP:HB2	1:D:430:ASP:HB2	1.79	0.63
1:B:216:ARG:NH2	1:B:389:ALA:O	2.31	0.63
1:C:113:PRO:HA	1:C:118:LEU:HD21	1.81	0.63
1:B:247:ALA:HA	1:B:272:LEU:HD23	1.81	0.62
1:A:113:PRO:HA	1:A:118:LEU:HD21	1.82	0.61
1:C:73:LEU:HD21	1:C:140:MSE:HE1	1.83	0.61
1:D:274:LEU:HB3	1:D:356:VAL:HB	1.83	0.60
1:B:227:LEU:HD22	1:B:451:MSE:HE3	1.82	0.60
1:B:111:THR:OG1	1:B:120:ARG:NH2	2.35	0.60
1:C:227:LEU:HD22	1:C:451:MSE:HE3	1.84	0.59
1:A:415:ASP:HB3	1:A:430:ASP:HB2	1.85	0.59
1:B:168:ARG:HB3	1:B:289:VAL:HG22	1.86	0.58
1:A:73:LEU:HD21	1:A:140:MSE:HE1	1.87	0.57
1:B:414:THR:HG22	1:B:431:GLN:HG2	1.87	0.57
1:C:65:VAL:HG21	1:C:75:MSE:HE3	1.86	0.56
1:C:168:ARG:HB3	1:C:289:VAL:HG22	1.88	0.55
1:A:111:THR:OG1	1:A:120:ARG:NH2	2.43	0.52
1:D:397:ARG:NH2	2:D:501:SO4:O1	2.42	0.51
1:D:73:LEU:HD21	1:D:140:MSE:HE1	1.92	0.51
1:B:65:VAL:HG11	1:B:75:MSE:HE3	1.93	0.50
1:D:242:MSE:HE2	1:D:416:LEU:HD21	1.93	0.50
1:D:281:ARG:NH2	1:D:291:GLY:O	2.44	0.50
1:C:163:ARG:O	1:C:166:GLN:HG2	2.11	0.50
1:A:96:GLN:HG2	1:A:126:LEU:HD13	1.93	0.50
1:B:36:ARG:NH1	3:B:612:HOH:O	2.46	0.49
1:B:217:LEU:HD11	1:B:448:ASP:HB2	1.95	0.49
1:A:11:ARG:NH1	1:A:171:ALA:O	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:17:GLN:HB3	1:D:140:MSE:HE3	1.95	0.48
1:A:144:ALA:HA	1:A:147:ARG:NH1	2.28	0.48
1:B:113:PRO:HA	1:B:118:LEU:HD21	1.96	0.48
1:C:240:LEU:HB3	1:C:315:MSE:HE1	1.96	0.48
1:C:17:GLN:HB3	1:C:140:MSE:HE3	1.95	0.47
1:A:65:VAL:HG21	1:A:75:MSE:HE3	1.95	0.47
1:B:274:LEU:HB3	1:B:356:VAL:HB	1.95	0.47
1:D:217:LEU:HB2	1:D:444:LEU:HD13	1.96	0.47
1:B:67:PHE:HB3	1:B:73:LEU:HD11	1.96	0.47
1:C:415:ASP:HB3	1:C:430:ASP:HB2	1.96	0.47
1:A:190:ARG:HH21	1:A:194:ARG:HH12	1.64	0.46
1:A:306:LEU:HD22	1:A:309:PRO:HB3	1.97	0.46
1:C:144:ALA:HA	1:C:147:ARG:NH1	2.30	0.46
1:D:156:GLN:O	1:D:161:ARG:N	2.49	0.46
1:D:252:THR:O	1:D:253:GLN:HB2	2.16	0.46
1:A:302:ASP:O	1:A:317:ARG:HD3	2.16	0.46
1:C:64:ARG:HD3	1:C:81:ALA:HB2	1.98	0.45
1:B:163:ARG:O	1:B:166:GLN:HG2	2.16	0.45
1:A:215:SER:HB3	1:D:402:THR:HG22	1.99	0.45
1:A:89:LEU:HD13	1:A:98:ARG:HB2	1.98	0.45
1:A:24:LEU:HD22	1:A:32:ALA:HB2	1.98	0.44
1:C:24:LEU:HD22	1:C:32:ALA:HB2	2.00	0.44
1:C:177:HIS:HA	1:C:178:PRO:HD3	1.90	0.44
1:C:112:ASP:OD1	1:C:114:GLU:HG2	2.17	0.44
1:B:64:ARG:HD3	1:B:81:ALA:HB2	2.00	0.44
1:D:106:ASP:OD2	1:D:122:HIS:NE2	2.32	0.44
1:C:281:ARG:HA	1:C:286:LEU:HB2	2.00	0.44
1:D:418:VAL:HG12	1:D:427:ILE:HD13	1.99	0.43
1:C:28:PRO:HA	1:C:29:PRO:HD3	1.87	0.43
1:A:403:TRP:CZ3	1:B:401:ASP:HB2	2.54	0.43
1:B:229:GLU:HA	1:B:230:PRO:HD3	1.90	0.43
1:C:96:GLN:HG2	1:C:126:LEU:HD13	2.00	0.43
1:D:36:ARG:NH2	2:D:501:SO4:O4	2.49	0.43
1:A:313:ASP:O	1:A:317:ARG:HG3	2.18	0.43
1:B:223:ARG:HG2	1:B:455:VAL:HG21	2.01	0.43
1:C:94:ASP:O	1:C:98:ARG:HG2	2.19	0.42
1:D:182:GLU:O	1:D:186:VAL:HG13	2.19	0.42
1:B:105:ALA:HB1	1:C:452:ALA:O	2.19	0.42
1:B:210:GLY:O	1:B:432:ARG:NH1	2.45	0.42
1:D:163:ARG:O	1:D:166:GLN:HG2	2.19	0.42
1:C:182:GLU:HA	1:C:185:ARG:HG2	1.99	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:234:LEU:HB2	1:B:311:PHE:CE2	2.54	0.42
1:B:106:ASP:O	1:B:120:ARG:NH1	2.52	0.42
1:C:14:SER:O	1:C:18:ARG:HG3	2.19	0.42
1:C:45:ASP:HB3	1:C:48:ARG:HB3	2.02	0.42
1:A:112:ASP:OD1	1:A:114:GLU:HG2	2.20	0.42
1:B:22:ARG:NH1	1:B:337:ASP:OD2	2.53	0.41
1:D:21:TRP:HE3	1:D:140:MSE:SE	2.54	0.41
1:C:92:HIS:HA	1:C:93:PRO:HD3	1.90	0.41
1:C:143:ASP:O	1:C:147:ARG:HG3	2.21	0.41
1:C:330:LEU:HD12	1:C:331:PRO:HD2	2.03	0.41
1:A:145:ARG:O	1:A:149:MSE:HG3	2.21	0.41
1:D:144:ALA:HA	1:D:147:ARG:NH1	2.36	0.40
1:B:415:ASP:HB2	1:B:430:ASP:OD1	2.21	0.40
1:C:22:ARG:NH1	1:C:333:HIS:HB2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	379/493 (77%)	367 (97%)	12 (3%)	0	100	100
1	B	404/493 (82%)	390 (96%)	14 (4%)	0	100	100
1	C	389/493 (79%)	374 (96%)	13 (3%)	2 (0%)	24	48
1	D	409/493 (83%)	394 (96%)	15 (4%)	0	100	100
All	All	1581/1972 (80%)	1525 (96%)	54 (3%)	2 (0%)	48	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	304	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	463	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/386 (83%)	319 (100%)	0	100	100
1	B	330/386 (86%)	324 (98%)	6 (2%)	51	78
1	C	321/386 (83%)	319 (99%)	2 (1%)	78	91
1	D	334/386 (86%)	328 (98%)	6 (2%)	51	78
All	All	1304/1544 (84%)	1290 (99%)	14 (1%)	65	85

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	101	GLU
1	B	229	GLU
1	B	234	LEU
1	B	274	LEU
1	B	282	ASP
1	B	401	ASP
1	C	180	ASP
1	C	285	GLN
1	D	10	LEU
1	D	18	ARG
1	D	26	GLU
1	D	209	VAL
1	D	274	LEU
1	D	303	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	333	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	92	HIS
1	C	166	GLN
1	D	453	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	B	501	-	4,4,4	0.24	0	6,6,6	0.17	0
2	SO4	A	502	-	4,4,4	0.24	0	6,6,6	0.15	0
2	SO4	D	501	-	4,4,4	0.25	0	6,6,6	0.18	0
2	SO4	A	501	-	4,4,4	0.25	0	6,6,6	0.22	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	SO4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	386/493 (78%)	-0.07	13 (3%)	48 44	12, 30, 64, 92	0
1	B	407/493 (82%)	0.38	24 (5%)	28 25	19, 47, 83, 115	0
1	C	394/493 (79%)	0.08	22 (5%)	30 26	11, 35, 75, 104	0
1	D	412/493 (83%)	0.25	28 (6%)	23 20	11, 38, 80, 110	0
All	All	1599/1972 (81%)	0.17	87 (5%)	31 27	11, 37, 77, 115	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	360	LEU	7.4
1	B	270	ASP	5.8
1	D	207	GLY	5.0
1	B	403	TRP	4.9
1	B	433	ARG	4.8
1	C	393	GLY	4.4
1	B	406	THR	4.2
1	C	395	ASP	4.1
1	B	9	GLY	4.0
1	C	464	HIS	4.0
1	C	434	THR	4.0
1	C	396	THR	3.9
1	C	230	PRO	3.9
1	D	28	PRO	3.7
1	D	11	ARG	3.6
1	B	404	ASP	3.6
1	D	27	ALA	3.6
1	D	434	THR	3.5
1	A	433	ARG	3.5
1	D	305	GLY	3.5
1	B	251	TRP	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	210	GLY	3.5
1	C	208	PRO	3.5
1	A	230	PRO	3.4
1	D	209	VAL	3.4
1	B	408	THR	3.3
1	C	394	GLY	3.3
1	A	238	GLY	3.3
1	D	23	TRP	3.2
1	C	283	HIS	3.1
1	B	236	GLY	3.1
1	A	237	ASN	3.1
1	A	393	GLY	3.0
1	B	26	GLU	3.0
1	B	370	THR	3.0
1	B	368	SER	3.0
1	D	29	PRO	3.0
1	A	239	SER	2.9
1	B	401	ASP	2.9
1	C	400	THR	2.8
1	B	463	PRO	2.8
1	C	433	ARG	2.8
1	B	402	THR	2.7
1	D	26	GLU	2.7
1	D	234	LEU	2.7
1	B	307	ARG	2.7
1	D	10	LEU	2.7
1	B	210	GLY	2.7
1	D	236	GLY	2.7
1	B	235	GLY	2.7
1	C	238	GLY	2.7
1	D	370	THR	2.7
1	D	25	ARG	2.6
1	D	433	ARG	2.6
1	C	361	CYS	2.6
1	C	9	GLY	2.6
1	C	308	GLU	2.6
1	D	405	GLY	2.6
1	C	194	ARG	2.5
1	D	406	THR	2.5
1	D	270	ASP	2.4
1	A	401	ASP	2.4
1	A	441	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	403	TRP	2.4
1	D	304	THR	2.3
1	B	287	GLY	2.3
1	A	270	ASP	2.2
1	C	282	ASP	2.2
1	B	405	GLY	2.2
1	B	382	SER	2.2
1	C	11	ARG	2.2
1	A	115	ARG	2.1
1	A	228	THR	2.1
1	B	202	ARG	2.1
1	C	209	VAL	2.1
1	D	208	PRO	2.1
1	C	82	ASP	2.1
1	D	230	PRO	2.1
1	D	368	SER	2.1
1	D	218	ARG	2.1
1	D	408	THR	2.1
1	A	71	ASP	2.0
1	C	305	GLY	2.0
1	B	23	TRP	2.0
1	D	156	GLN	2.0
1	B	91	GLY	2.0
1	C	307	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	501	5/5	0.91	0.13	67,74,80,82	0
2	SO4	A	502	5/5	0.92	0.17	75,76,78,82	0
2	SO4	D	501	5/5	0.94	0.09	39,42,45,52	0
2	SO4	B	501	5/5	0.97	0.10	49,51,54,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.