



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:56 PM UTC

PDB ID : 5HVH / pdb_00005hvh
Title : Crystal Structure of Thrombin-activatable Fibrinolysis Inhibitor in Complex with two Inhibitory Nanobodies
Authors : Zhou, X.; Weeks, S.D.; Strelkov, S.V.; Declerck, P.J.
Deposited on : 2016-01-28
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	401	
2	B	128	
3	C	125	
4	D	2	
4	E	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	F	2	100%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5160 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Carboxypeptidase B2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	401	Total	C	N	O	S	0	0	0
			3253	2089	552	599	13			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	305	CYS	SER	engineered mutation	UNP Q96IY4
A	329	ILE	THR	engineered mutation	UNP Q96IY4
A	333	TYR	HIS	engineered mutation	UNP Q96IY4
A	335	GLN	HIS	engineered mutation	UNP Q96IY4

- Molecule 2 is a protein called VHH-a204.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	120	Total	C	N	O	S	0	0	0
			912	566	159	183	4			

- Molecule 3 is a protein called VHH-i83.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	118	Total	C	N	O	S	0	0	0
			896	556	159	177	4			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	F	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Carboxypeptidase B2



- Molecule 2: VHH-a204



- Molecule 3: VHH-i83



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:  100%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  100%

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 62 2 2	Depositor
Cell constants a, b, c, α , β , γ	196.99Å 196.99Å 147.51Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.25 – 3.00 49.25 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.25-3.00) 100.0 (49.25-3.00)	Depositor EDS
R_{merge}	0.27	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 3.01Å)	Xtriage
Refinement program	BUSTER-TNT 2.10.2	Depositor
R, R_{free}	0.190 , 0.202 0.200 , 0.208	Depositor DCC
R_{free} test set	1938 reflections (5.65%)	wwPDB-VP
Wilson B-factor (Å ²)	61.5	Xtriage
Anisotropy	0.641	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 70.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5160	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	1/3341 (0.0%)	1.30	13/4538 (0.3%)
2	B	0.79	0/933	1.16	0/1267
3	C	0.80	0/915	1.16	2/1241 (0.2%)
All	All	0.86	1/5189 (0.0%)	1.25	15/7046 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	327	LYS	CA-C	5.62	1.60	1.52

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	94	SER	N-CA-C	-10.38	95.15	110.52
3	C	48	VAL	N-CA-C	-9.76	102.31	111.48
1	A	342	LEU	CA-C-N	-6.58	116.26	122.37
1	A	342	LEU	C-N-CA	-6.58	116.26	122.37
1	A	348	ASP	CA-CB-CG	6.15	118.75	112.60
1	A	308	HIS	CA-C-N	6.08	131.22	122.40
1	A	308	HIS	C-N-CA	6.08	131.22	122.40
1	A	95	ALA	CA-C-N	5.63	127.83	120.28
1	A	95	ALA	C-N-CA	5.63	127.83	120.28
1	A	338	GLU	CB-CG-CD	5.52	121.98	112.60
3	C	100	ASP	CA-CB-CG	5.43	118.03	112.60
1	A	22	ASN	OD1-CG-ND2	5.40	128.00	122.60
1	A	366	ASP	CA-CB-CG	5.39	117.99	112.60
1	A	22	ASN	CB-CG-ND2	-5.25	108.53	116.40
1	A	240	LYS	N-CA-CB	5.02	118.03	110.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3253	0	3169	25	0
2	B	912	0	857	7	0
3	C	896	0	847	8	0
4	D	28	0	25	0	0
4	E	28	0	25	0	0
4	F	28	0	25	0	0
5	A	14	0	13	1	0
6	A	1	0	0	0	0
All	All	5160	0	4961	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:90:THR:HG22	2:B:120:VAL:H	1.30	0.94
1:A:2:GLN:HE22	1:A:213:ASN:HA	1.33	0.92
1:A:145:GLU:OE1	1:A:191:ARG:HA	1.83	0.78
2:B:82:MET:HE2	2:B:85:LEU:HD21	1.71	0.72
1:A:325:ILE:HG21	1:A:385:GLU:HA	1.84	0.59
1:A:139:LEU:HD12	1:A:173:ILE:HD13	1.85	0.59
2:B:90:THR:HG22	2:B:120:VAL:N	2.12	0.58
1:A:367:THR:HG21	3:C:104:GLY:HA2	1.88	0.55
3:C:89:THR:HG23	3:C:116:THR:HA	1.88	0.54
1:A:93:ALA:HB3	1:A:113:PHE:CZ	2.42	0.53
1:A:93:ALA:HB3	1:A:113:PHE:CE2	2.44	0.52
1:A:71:VAL:HG23	3:C:54:VAL:HG22	1.92	0.51
3:C:81:MET:HE2	3:C:84:LEU:HD21	1.92	0.51
2:B:40:ALA:HB3	2:B:43:LYS:HB2	1.92	0.51
1:A:246:GLY:HA3	1:A:342:LEU:HB3	1.94	0.49
1:A:329:ILE:HG13	1:A:365:ARG:NH2	2.26	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:29:PHE:CD2	3:C:31:PRO:HD3	2.47	0.49
1:A:302:ARG:HH21	1:A:342:LEU:HD21	1.79	0.48
1:A:240:LYS:O	1:A:240:LYS:HD3	2.14	0.48
2:B:34:MET:HB2	2:B:78:VAL:HG11	1.96	0.47
1:A:283:ALA:HB2	1:A:397:VAL:HG21	1.99	0.45
1:A:62:LEU:HD13	1:A:69:CYS:HB2	1.99	0.45
1:A:93:ALA:CB	1:A:113:PHE:CE2	2.99	0.44
1:A:93:ALA:CB	1:A:113:PHE:HE2	2.30	0.44
1:A:2:GLN:NE2	1:A:213:ASN:HA	2.16	0.44
1:A:93:ALA:HB2	1:A:113:PHE:HE2	1.83	0.43
1:A:295:VAL:HB	1:A:361:THR:HB	2.01	0.43
1:A:299:SER:O	1:A:303:SER:HB2	2.19	0.43
3:C:66:PHE:CE2	3:C:81:MET:HG2	2.54	0.43
1:A:297:PRO:HB3	1:A:349:ASP:HA	2.02	0.42
3:C:106:TYR:HB2	3:C:108:TYR:CE2	2.55	0.41
1:A:95:ALA:HB3	2:B:101:PRO:HB2	2.01	0.41
1:A:100:GLN:HE21	5:A:507:NAG:H81	1.86	0.41
1:A:209:SER:HA	1:A:213:ASN:O	2.20	0.41
1:A:397:VAL:O	1:A:401:VAL:HG22	2.21	0.41
3:C:27:SER:HB2	3:C:103:LEU:HD21	2.02	0.41
2:B:12:VAL:HG11	2:B:85:LEU:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/401 (100%)	382 (96%)	15 (4%)	2 (0%)	24	60
2	B	118/128 (92%)	114 (97%)	4 (3%)	0	100	100
3	C	116/125 (93%)	114 (98%)	2 (2%)	0	100	100
All	All	633/654 (97%)	610 (96%)	21 (3%)	2 (0%)	36	70

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	291	SER
1	A	305	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	354/354 (100%)	351 (99%)	3 (1%)	73	86
2	B	96/104 (92%)	94 (98%)	2 (2%)	47	75
3	C	92/99 (93%)	85 (92%)	7 (8%)	12	41
All	All	542/557 (97%)	530 (98%)	12 (2%)	45	74

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	69	CYS
1	A	245	GLU
2	B	95	CYS
2	B	106	VAL
3	C	11	LEU
3	C	12	VAL
3	C	13	GLN
3	C	25	SER
3	C	44	GLU
3	C	61	SER
3	C	83	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	57	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	118	HIS
1	A	280	GLN
1	A	328	ASN
1	A	396	HIS
2	B	13	GLN
2	B	117	GLN
3	C	99	GLN
3	C	105	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	1	1,4	14,14,15	0.39	0	17,19,21	1.74	2 (11%)
4	NAG	D	2	4	14,14,15	0.45	0	17,19,21	1.66	3 (17%)
4	NAG	E	1	1,4	14,14,15	0.34	0	17,19,21	0.89	1 (5%)
4	NAG	E	2	4	14,14,15	0.34	0	17,19,21	0.74	1 (5%)
4	NAG	F	1	1,4	14,14,15	0.30	0	17,19,21	0.96	1 (5%)
4	NAG	F	2	4	14,14,15	0.32	0	17,19,21	0.84	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1
4	NAG	E	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	E	2	4	-	0/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	F	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	C1-O5-C5	5.42	119.45	112.19
4	D	2	NAG	C1-O5-C5	4.21	117.83	112.19
4	D	2	NAG	O5-C1-C2	4.17	117.74	111.29
4	F	1	NAG	C1-O5-C5	3.37	116.70	112.19
4	D	1	NAG	O5-C1-C2	3.10	116.09	111.29
4	E	1	NAG	C1-O5-C5	3.08	116.31	112.19
4	D	2	NAG	C2-N2-C7	3.03	126.96	122.90
4	E	2	NAG	O5-C1-C2	-2.34	107.68	111.29
4	F	2	NAG	C1-O5-C5	2.31	115.28	112.19

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	F	2	NAG	O5-C5-C6-O6
4	E	1	NAG	C4-C5-C6-O6
4	F	2	NAG	C4-C5-C6-O6
4	D	2	NAG	C1-C2-N2-C7
4	D	2	NAG	C3-C2-N2-C7


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	507	1	14,14,15	0.35	0	17,19,21	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	507	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	507	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	401/401 (100%)	0.21	27 (6%) 24 12	39, 58, 107, 143	0
2	B	120/128 (93%)	1.44	29 (24%) 2 1	79, 135, 163, 168	0
3	C	118/125 (94%)	-0.10	2 (1%) 69 45	42, 71, 97, 130	0
All	All	639/654 (97%)	0.39	58 (9%) 15 8	39, 69, 150, 168	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	340	LEU	6.8
1	A	1	PHE	6.7
1	A	306	LYS	6.5
3	C	119	SER	5.7
1	A	304	LYS	4.8
1	A	327	LYS	4.3
2	B	14	PRO	4.2
2	B	122	SER	4.2
2	B	18	LEU	4.1
2	B	87	PRO	4.1
2	B	11	LEU	4.0
1	A	338	GLU	3.9
1	A	240	LYS	3.8
1	A	144	LYS	3.7
2	B	120	VAL	3.7
2	B	10	GLY	3.6
2	B	12	VAL	3.6
2	B	86	LYS	3.6
2	B	44	GLN	3.5
2	B	40	ALA	3.5
1	A	307	ASP	3.5
2	B	119	THR	3.5
2	B	64	LYS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	41	PRO	3.4
1	A	339	THR	3.2
1	A	147	ALA	3.2
3	C	28	ILE	3.1
1	A	337	SER	3.1
2	B	43	LYS	3.1
1	A	120	ASP	3.0
1	A	401	VAL	3.0
2	B	105	ASP	3.0
1	A	65	SER	2.8
2	B	75	LYS	2.7
1	A	342	LEU	2.7
2	B	7	SER	2.7
1	A	305	CYS	2.7
2	B	85	LEU	2.6
2	B	113	GLY	2.6
1	A	185	GLN	2.6
1	A	326	SER	2.6
2	B	121	SER	2.5
1	A	308	HIS	2.5
1	A	343	ALA	2.5
2	B	22	CYS	2.5
2	B	21	SER	2.4
2	B	95	CYS	2.3
2	B	5	GLN	2.3
1	A	188	ASN	2.3
1	A	145	GLU	2.2
1	A	349	ASP	2.2
2	B	68	THR	2.2
1	A	341	TYR	2.1
1	A	148	ALA	2.1
1	A	113	PHE	2.1
2	B	69	ILE	2.1
2	B	84	SER	2.0
2	B	67	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	D	2	14/15	0.22	0.23	135,142,145,145	0
4	NAG	F	2	14/15	0.52	0.20	124,130,132,133	0
4	NAG	D	1	14/15	0.55	0.26	121,135,140,140	0
4	NAG	E	2	14/15	0.71	0.17	117,125,127,127	0
4	NAG	F	1	14/15	0.88	0.15	107,110,115,122	0
4	NAG	E	1	14/15	0.88	0.14	104,109,115,120	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain E:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	A	507	14/15	0.73	0.22	131,133,135,136	0
6	ZN	A	508	1/1	0.98	0.05	59,59,59,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.