



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:04 PM UTC

PDB ID : 6HVF / pdb_00006hvf
Title : Kinase domain of cSrc in complex with compound 29B
Authors : Keul, M.; Mueller, M.P.; Rauh, D.
Deposited on : 2018-10-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	16% 69% 17% 14%
1	B	286	14% 70% 15% 15%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4165 atoms, of which 6 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	6	0
			1984	1275	329	362	18			
1	B	244	Total	C	N	O	S	0	1	0
			1918	1225	317	359	17			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
A	338	MET	THR	engineered mutation	UNP P00523
A	345	CYS	SER	engineered mutation	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523
B	338	MET	THR	engineered mutation	UNP P00523
B	345	CYS	SER	engineered mutation	UNP P00523

- Molecule 2 is {N}-[3-[3-ethyl-6-[4-(4-methylpiperazin-1-yl)phenyl]-4-oxidanylidene-7 {H}-pyrrolo[2,3-d]pyrimidin-5-yl]phenyl]prop-2-enamide (CCD ID: GUT) (formula: C₂₈H₃₀N₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			36	28	6	2		
2	B	1	Total	C	N	O	0	0
			36	28	6	2		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			10	2	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	76	Total	O	0	0
			76	76		
4	B	97	Total	O	0	0
			97	97		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.17Å 62.97Å 74.54Å 100.23° 91.64° 90.03°	Depositor
Resolution (Å)	43.66 – 2.10 43.66 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.5 (43.66-2.10) 97.6 (43.66-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.211 , 0.245 0.213 , 0.247	Depositor DCC
R_{free} test set	2148 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtriage
Anisotropy	0.543	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 58.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4165	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GUT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.16	0/2047	0.34	0/2770
1	B	0.16	0/1963	0.35	0/2662
All	All	0.16	0/4010	0.35	0/5432

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1984	0	1958	36	2
1	B	1918	0	1851	28	1
2	A	36	0	0	0	0
2	B	36	0	0	1	0
3	A	8	0	12	0	0
3	B	4	6	6	1	0
4	A	76	0	0	1	0
4	B	97	0	0	1	0
All	All	4159	6	3827	64	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:MET:HE3	1:A:291:ARG:HD2	1.42	0.98
1:B:318:ARG:HA	1:B:324:GLN:HE22	1.29	0.96
1:B:426:ILE:HD11	1:B:434:ALA:HB1	1.63	0.79
1:B:310:GLU:O	1:B:313:VAL:HG22	1.86	0.75
1:A:283:MET:CE	1:A:291:ARG:HD2	2.17	0.71
1:B:318:ARG:HG3	1:B:324:GLN:NE2	2.05	0.70
1:A:289:THR:HG23	1:A:290:THR:HG23	1.75	0.69
1:B:329:VAL:HG22	1:B:335:TYR:HB2	1.75	0.68
1:A:281:VAL:HG22	1:A:295:LYS:HG2	1.77	0.67
1:A:264:ARG:NH2	1:A:333:PRO:O	2.18	0.67
1:B:270:GLU:OE1	1:B:285:THR:HG23	1.98	0.62
1:B:388:ARG:HB3	1:B:428:TRP:CD1	2.36	0.60
1:B:362:GLN:O	1:B:366:MET:HG3	2.01	0.60
1:A:362:GLN:O	1:A:366:MET:HG3	2.03	0.59
1:A:513:GLN:OE1	4:A:701:HOH:O	2.17	0.59
1:B:466:MET:HE3	1:B:470:GLU:HG2	1.85	0.59
1:A:355:GLY:O	1:A:458:LYS:HD3	2.04	0.58
1:A:272:LYS:NZ	1:A:275:GLN:OE1	2.27	0.58
1:A:308:LEU:HD22	1:A:309:GLN:N	2.21	0.56
1:A:272:LYS:HE2	1:A:275:GLN:HG3	1.90	0.54
1:A:272:LYS:HZ1	1:A:275:GLN:CD	2.12	0.52
1:B:318:ARG:CA	1:B:324:GLN:HE22	2.14	0.52
1:A:314[A]:MET:HE2	1:A:325:LEU:HB2	1.92	0.52
1:A:511:TYR:HE2	1:B:351:LYS:O	1.93	0.52
1:A:493[B]:ASP:OD2	1:A:497:GLN:NE2	2.43	0.52
1:B:329:VAL:CG2	1:B:335:TYR:HB2	2.41	0.50
1:B:348:ASP:OD1	4:B:701:HOH:O	2.20	0.49
2:B:601:GUT:CAL	2:B:601:GUT:CAS	2.91	0.49
1:A:308:LEU:CD1	1:A:308:LEU:H	2.26	0.48
1:A:388:ARG:HB3	1:A:428:TRP:CD1	2.48	0.48
1:B:355:GLY:O	1:B:458:LYS:NZ	2.44	0.48
1:A:309:GLN:O	1:A:313:VAL:HG23	2.13	0.48
1:B:283:MET:HG3	1:B:340:TYR:CZ	2.49	0.47
1:A:262:ILE:HG23	1:A:327:ALA:HB1	1.95	0.47
1:B:261:GLU:HA	1:B:328:VAL:O	2.15	0.47
1:B:320:GLU:O	1:B:401:LYS:HE2	2.15	0.46
1:A:308:LEU:HD22	1:A:308:LEU:C	2.41	0.46
1:B:261:GLU:OE2	1:B:330:SER:OG	2.27	0.46
1:A:311:ALA:O	1:A:315:LYS:HG3	2.16	0.45
1:B:356:LYS:HE2	1:B:357:TYR:CE2	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:286:TRP:HB3	1:A:290:THR:OG1	2.16	0.45
1:B:356:LYS:HE2	1:B:357:TYR:CZ	2.52	0.45
1:B:527:TYR:OH	1:B:533:LEU:O	2.31	0.45
1:A:321:LYS:O	1:A:322:LEU:HD12	2.17	0.45
1:A:451:LEU:HD12	1:A:451:LEU:O	2.17	0.45
1:A:308:LEU:H	1:A:308:LEU:HD13	1.82	0.45
1:A:272:LYS:HE2	1:A:275:GLN:CG	2.47	0.44
1:A:332:GLU:HA	1:A:333:PRO:C	2.41	0.44
1:B:369:GLN:HA	3:B:602:EDO:H12	1.99	0.44
1:B:325:LEU:HD12	1:B:337:VAL:O	2.18	0.44
1:A:262:ILE:CG2	1:A:327:ALA:HB1	2.48	0.43
1:A:314[A]:MET:SD	1:A:338:MET:HE2	2.58	0.43
1:B:331:GLU:HA	1:B:331:GLU:OE1	2.18	0.43
1:A:349:PHE:CZ	1:A:354:MET:HG3	2.54	0.42
1:A:490:SER:OG	1:A:524:GLU:OE2	2.33	0.42
1:A:269:LEU:HD12	1:A:269:LEU:N	2.34	0.42
1:A:257:LYS:HB2	1:A:257:LYS:HE3	1.83	0.42
1:A:320:GLU:O	1:A:401:LYS:HE2	2.19	0.42
1:B:291:ARG:HG2	1:B:340:TYR:CD2	2.55	0.42
1:A:361:PRO:HA	1:A:520:PHE:CE2	2.55	0.41
1:B:318:ARG:HG3	1:B:324:GLN:HE21	1.82	0.41
1:B:403:ALA:O	1:B:404:ASP:C	2.63	0.41
1:A:308:LEU:HD13	1:A:308:LEU:N	2.36	0.41
1:B:262:ILE:HG21	1:B:286:TRP:CZ2	2.57	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:397:ASN:OD1	1:B:438:ARG:NE[1_655]	2.00	0.20
1:A:397:ASN:OD1	1:A:438:ARG:NE[1_455]	2.07	0.13
1:A:359[A]:ARG:NH1	1:A:436:TYR:OH[1_455]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/286 (85%)	240 (98%)	4 (2%)	0	100	100
1	B	237/286 (83%)	233 (98%)	4 (2%)	0	100	100
All	All	481/572 (84%)	473 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/245 (86%)	211 (100%)	1 (0%)	81	88
1	B	201/245 (82%)	199 (99%)	2 (1%)	68	76
All	All	413/490 (84%)	410 (99%)	3 (1%)	76	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	308	LEU
1	B	356	LYS
1	B	472	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	497	GLN
1	B	324	GLN
1	B	381	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GUT	A	601	1	40,40,40	2.75	13 (32%)	53,57,57	2.37	10 (18%)
2	GUT	B	601	1	40,40,40	2.76	13 (32%)	53,57,57	2.33	10 (18%)
3	EDO	A	603	-	3,3,3	0.46	0	2,2,2	0.37	0
3	EDO	B	602	-	3,3,3	0.43	0	2,2,2	0.33	0
3	EDO	A	602	-	3,3,3	0.43	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GUT	A	601	1	-	2/20/30/30	0/5/5/5
2	GUT	B	601	1	-	4/20/30/30	0/5/5/5
3	EDO	A	603	-	-	0/1/1/1	-
3	EDO	B	602	-	-	1/1/1/1	-
3	EDO	A	602	-	-	1/1/1/1	-

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	GUT	CAR-CAH	-8.60	1.34	1.47
2	B	601	GUT	CAR-CAH	-8.43	1.35	1.47
2	B	601	GUT	CAJ-CAI	-7.66	1.35	1.49
2	A	601	GUT	CAJ-CAI	-7.41	1.35	1.49
2	A	601	GUT	CBH-CBG	4.94	1.54	1.30
2	B	601	GUT	CBH-CBG	4.94	1.54	1.30
2	B	601	GUT	C2-N3	4.78	1.39	1.30
2	A	601	GUT	C2-N3	4.75	1.39	1.30
2	B	601	GUT	C5-C6	-4.75	1.34	1.44
2	A	601	GUT	C5-C4	-4.67	1.34	1.45
2	B	601	GUT	C5-C4	-4.67	1.34	1.45
2	A	601	GUT	C5-C6	-4.52	1.35	1.44
2	A	601	GUT	CBG-CBE	3.96	1.54	1.48
2	B	601	GUT	CAI-C5	-3.89	1.35	1.44
2	B	601	GUT	CBG-CBE	3.77	1.54	1.48
2	A	601	GUT	CAI-C5	-3.61	1.35	1.44
2	B	601	GUT	CAM-NAQ	-3.36	1.34	1.41
2	A	601	GUT	CAM-NAQ	-3.23	1.35	1.41
2	B	601	GUT	C6-N1	-2.90	1.35	1.39
2	A	601	GUT	C6-N1	-2.87	1.35	1.39
2	A	601	GUT	CAH-NAG	-2.79	1.32	1.37
2	B	601	GUT	CAH-NAG	-2.69	1.33	1.37
2	A	601	GUT	CAI-CAH	-2.65	1.35	1.39
2	A	601	GUT	C4-NAG	-2.55	1.32	1.37
2	B	601	GUT	CAI-CAH	-2.47	1.35	1.39
2	B	601	GUT	C4-NAG	-2.30	1.33	1.37

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GUT	CBH-CBG-CBE	-9.94	111.30	122.22
2	B	601	GUT	CBH-CBG-CBE	-9.54	111.74	122.22
2	A	601	GUT	N1-C2-N3	-9.50	117.18	126.45
2	B	601	GUT	N1-C2-N3	-9.15	117.52	126.45
2	B	601	GUT	CBC-NAX-CAY	4.11	120.81	111.57
2	B	601	GUT	C2-N1-C6	-3.99	120.21	123.06
2	A	601	GUT	C2-N1-C6	-3.85	120.31	123.06
2	A	601	GUT	CBC-NAX-CAY	3.84	120.21	111.57
2	A	601	GUT	CAM-NAQ-CBE	-3.47	121.53	128.07
2	B	601	GUT	CAS-CAR-CAH	3.28	124.67	120.76
2	A	601	GUT	CAW-CAR-CAH	3.25	124.62	120.76
2	A	601	GUT	CAS-CAR-CAH	-3.07	117.11	120.76
2	B	601	GUT	CAM-NAQ-CBE	-2.87	122.68	128.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	GUT	C2-N3-C4	2.67	120.10	113.43
2	B	601	GUT	C2-N3-C4	2.63	119.99	113.43
2	A	601	GUT	O6-C6-N1	-2.55	117.13	120.53
2	B	601	GUT	CAW-CAR-CAH	-2.54	117.75	120.76
2	A	601	GUT	C5-C6-N1	2.34	121.52	116.42
2	B	601	GUT	CAJ-CAL-CAM	2.22	123.29	120.44
2	B	601	GUT	C5-C6-N1	2.09	120.97	116.42

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	601	GUT	CBJ-CBI-N1-C6
2	B	601	GUT	CBJ-CBI-N1-C2
2	B	601	GUT	NAQ-CBE-CBG-CBH
2	B	601	GUT	OBFCBE-CBG-CBH
2	A	601	GUT	CAL-CAM-NAQ-CBE
2	A	601	GUT	CAN-CAM-NAQ-CBE
3	B	602	EDO	O1-C1-C2-O2
3	A	602	EDO	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	GUT	1	0
3	B	602	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	246/286 (86%)	1.02	45 (18%)	3 3	20, 41, 100, 119	6 (2%)
1	B	244/286 (85%)	0.96	41 (16%)	4 4	21, 41, 94, 111	1 (0%)
All	All	490/572 (85%)	0.99	86 (17%)	4 4	20, 41, 96, 119	7 (1%)

All (86) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	308	LEU	6.9
1	B	309	GLN	5.3
1	B	403	ALA	5.1
1	B	259	ALA	4.8
1	A	307	PHE	4.6
1	A	328	VAL	4.6
1	B	329	VAL	4.5
1	A	279	GLY	4.5
1	B	260	TRP	4.3
1	A	334	ILE	4.3
1	A	314[A]	MET	4.2
1	A	355	GLY	4.2
1	A	405	PHE	3.8
1	B	313	VAL	3.7
1	B	312	GLN	3.7
1	A	276	GLY	3.6
1	B	311	ALA	3.6
1	A	329	VAL	3.6
1	A	257	LYS	3.6
1	A	476	GLU	3.6
1	A	311	ALA	3.4
1	B	297	LEU	3.4
1	B	298	LYS	3.3
1	A	404	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	258	ASP	3.3
1	A	331	GLU	3.3
1	A	310	GLU	3.2
1	A	309	GLN	3.2
1	B	472	LEU	3.2
1	A	335	TYR	3.1
1	A	265	GLU	3.1
1	B	425	PRO	3.0
1	A	313	VAL	3.0
1	A	330	SER	2.9
1	A	296	THR	2.9
1	B	296	THR	2.9
1	A	280	GLU	2.9
1	B	272	LYS	2.9
1	B	276	GLY	2.9
1	A	273	LEU	2.8
1	A	354	MET	2.8
1	A	283	MET	2.8
1	B	285	THR	2.8
1	B	404	ASP	2.8
1	A	290	THR	2.8
1	A	271	VAL	2.8
1	A	333	PRO	2.7
1	A	275	GLN	2.7
1	B	265	GLU	2.6
1	A	263	PRO	2.6
1	B	271	VAL	2.6
1	A	269	LEU	2.6
1	B	315	LYS	2.6
1	B	314	MET	2.6
1	B	381	ASN	2.6
1	A	353	GLU	2.5
1	B	426	ILE	2.5
1	B	275	GLN	2.5
1	A	288	GLY	2.5
1	A	282	TRP	2.5
1	B	316	LYS	2.4
1	B	286	TRP	2.4
1	B	274	GLY	2.4
1	B	476	GLU	2.4
1	B	531	GLU	2.4
1	A	312	GLN	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	281	VAL	2.3
1	B	317	LEU	2.3
1	A	281	VAL	2.3
1	B	533	LEU	2.3
1	B	353	GLU	2.2
1	B	288	GLY	2.2
1	A	472	LEU	2.2
1	B	269	LEU	2.2
1	A	426	ILE	2.2
1	B	326	TYR	2.2
1	B	386	ASP	2.2
1	B	289	THR	2.1
1	A	262	ILE	2.1
1	A	294	ILE	2.1
1	A	267	LEU	2.1
1	B	331	GLU	2.1
1	A	268	ARG	2.1
1	B	268	ARG	2.1
1	A	316	LYS	2.1
1	A	287	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	B	602	4/4	0.82	0.18	65,78,79,82	0
2	GUT	B	601	36/36	0.87	0.15	38,49,62,65	0
3	EDO	A	603	4/4	0.88	0.18	44,46,49,50	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	A	602	4/4	0.90	0.15	57,58,60,61	0
2	GUT	A	601	36/36	0.91	0.12	33,50,68,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around GUT B 601:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.