



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 02:42 PM UTC

PDB ID : 8HVS / pdb_00008hvs
BMRB ID : 36534
Title : Solution Structure of the Antimicrobial Peptide HT-2
Authors : Li, S.; Yang, A.
Deposited on : 2022-12-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 54%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	10	90% 10%

2 Ensemble composition and analysis ⓘ

This entry contains 20 models.

Cyrange was unable to find well-defined residues.

Error message: Only domains with < 8 residues could be identified.

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 210 atoms, of which 106 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE.


Mol	Chain	Residues	Atoms					Trace
1	A	10	Total	C	H	N	O	0
			210	75	106	19	10	

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE



4.2.2 Score per residue for model 2

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE



There are no outlier residues in this chain.

4.2.3 Score per residue for model 3

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  90% 10%



4.2.4 Score per residue for model 4

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  80% 20%



4.2.5 Score per residue for model 5

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  70% 30%



4.2.6 Score per residue for model 6

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  70% 30%



4.2.7 Score per residue for model 7

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  70% 30%



4.2.8 Score per residue for model 8

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  80% 20%



4.2.9 Score per residue for model 9

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  100%

There are no outlier residues in this chain.

4.2.10 Score per residue for model 10

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  90% 10%



4.2.11 Score per residue for model 11


- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  70% 20% 10%



4.2.12 Score per residue for model 12

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  100%

There are no outlier residues in this chain.


4.2.13 Score per residue for model 13

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE



4.2.14 Score per residue for model 14

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE




4.2.15 Score per residue for model 15

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE




4.2.16 Score per residue for model 16

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE



4.2.17 Score per residue for model 17


- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE



4.2.18 Score per residue for model 18

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  70% 30%



4.2.19 Score per residue for model 19

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE


Chain A:  80% 20%



4.2.20 Score per residue for model 20

- Molecule 1: ARG-PHE-LEU-ARG-ARG-ILE-PHE-PHE-PHE-PHE

Chain A:  70% 30%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *distance geometry*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	106
Number of shifts mapped to atoms	106
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	54%

6 Model quality [i](#)

6.1 Standard geometry [i](#)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	2080	2120	2160	-

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is -.

There are no clashes.

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	8/10 (80%)	8±0 (96±6%)	0±0 (2±5%)	0±0 (1±4%)	12	60
All	All	160/200 (80%)	154 (96%)	4 (2%)	2 (1%)	12	60

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	2	PHE	2

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	10/10 (100%)	8±1 (78±13%)	2±1 (22±13%)	2	29
All	All	200/200 (100%)	155 (78%)	45 (22%)	2	29

All 8 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	3	LEU	11
1	A	7	PHE	9
1	A	1	ARG	7
1	A	5	ARG	6
1	A	4	ARG	4
1	A	8	PHE	4
1	A	2	PHE	3
1	A	6	ILE	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry

There are no ligands in this entry.

6.7 Other polymers

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 54% for the well-defined parts and 54% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *bmr_b_for_AS.txt*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	106
Number of shifts mapped to atoms	106
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 54%, i.e. 106 atoms were assigned a chemical shift out of a possible 195. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	27/50 (54%)	19/20 (95%)	8/20 (40%)	0/10 (0%)
Sidechain	69/95 (73%)	48/61 (79%)	21/25 (84%)	0/9 (0%)
Aromatic	10/50 (20%)	10/25 (40%)	0/25 (0%)	0/0 (—%)
Overall	106/195 (54%)	77/106 (73%)	29/70 (41%)	0/19 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 54%, i.e. 106 atoms were assigned a chemical shift out of a possible 195. 0 out of 1 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	27/50 (54%)	19/20 (95%)	8/20 (40%)	0/10 (0%)
Sidechain	69/95 (73%)	48/61 (79%)	21/25 (84%)	0/9 (0%)
Aromatic	10/50 (20%)	10/25 (40%)	0/25 (0%)	0/0 (—%)
Overall	106/195 (54%)	77/106 (73%)	29/70 (41%)	0/19 (0%)

7.1.4 Statistically unusual chemical shifts [i](#)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	6	ILE	CG2	28.30	10.93 – 24.12	8.2

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

