



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:37 PM UTC

PDB ID : 2HWO / pdb_00002hwo
Title : Crystal structure of Src kinase domain in complex with covalent inhibitor
Authors : Rauh, D.; Blair, J.A.; Shokat, K.M.
Deposited on : 2006-08-01
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	9% 69% 17% • 12%
1	B	286	4% 66% 16% • 15%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	251	Total	C	N	O	S	1	0	0
			1902	1225	317	345	15			
1	B	242	Total	C	N	O	S	0	0	0
			1887	1213	320	340	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	cloning artifact	UNP P00523
A	249	HIS	-	cloning artifact	UNP P00523
A	250	MET	-	cloning artifact	UNP P00523
A	345	CYS	SER	engineered mutation	UNP P00523
B	248	GLY	-	cloning artifact	UNP P00523
B	249	HIS	-	cloning artifact	UNP P00523
B	250	MET	-	cloning artifact	UNP P00523
B	345	CYS	SER	engineered mutation	UNP P00523

- Molecule 2 is N-(4-PHENYLAMINO-QUINAZOLIN-6-YL)-ACRYLAMIDE (CCD ID: RBS) (formula: C₁₇H₁₄N₄O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			22	17	4	1		
2	B	1	Total	C	N	O	0	0
			22	17	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	21	Total 21	O 21	0	0
4	B	19	Total 19	O 19	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.10Å 63.26Å 73.80Å 78.73° 88.04° 89.95°	Depositor
Resolution (Å)	72.36 – 2.50 72.33 – 2.50	Depositor EDS
% Data completeness (in resolution range)	97.5 (72.36-2.50) 97.5 (72.33-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.48 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.235 , 0.292 0.231 , 0.280	Depositor DCC
R_{free} test set	1278 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	31.2	Xtriage
Anisotropy	0.440	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.033 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3879	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.37% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, RBS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	3/1946 (0.2%)	1.15	7/2643 (0.3%)
1	B	0.92	0/1932	1.05	2/2623 (0.1%)
All	All	0.95	3/3878 (0.1%)	1.10	9/5266 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	463	TYR	C-O	6.68	1.27	1.23
1	A	270	GLU	CA-C	5.97	1.55	1.52
1	A	457	THR	CA-CB	5.54	1.61	1.53

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	483	CYS	CA-C-N	7.89	125.42	119.66
1	A	483	CYS	C-N-CA	7.89	125.42	119.66
1	A	333	PRO	N-CA-CB	7.80	111.58	103.00
1	A	330	SER	CA-C-O	-7.05	108.81	120.80
1	B	430	ALA	CA-C-N	5.46	124.92	119.24
1	B	430	ALA	C-N-CA	5.46	124.92	119.24
1	A	430	ALA	CA-C-N	5.40	124.85	119.24
1	A	430	ALA	C-N-CA	5.40	124.85	119.24
1	A	459	GLY	N-CA-C	5.09	122.34	115.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1902	0	1800	27	0
1	B	1887	0	1823	28	0
2	A	22	0	13	4	0
2	B	22	0	13	5	0
3	B	6	0	8	0	0
4	A	21	0	0	0	0
4	B	19	0	0	0	0
All	All	3879	0	3657	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:309:GLN:HA	1:A:310:GLU:C	2.06	0.81
1:B:338:THR:HG21	2:B:1345:RBS:H3	1.72	0.72
2:A:1345:RBS:H4	2:A:1345:RBS:N3	2.06	0.68
1:A:372:SER:HB2	1:A:513:GLN:NE2	2.08	0.67
1:B:388:ARG:HB3	1:B:428:TRP:CD1	2.32	0.65
2:B:1345:RBS:N3	2:B:1345:RBS:H4	2.14	0.61
1:A:329:VAL:HB	1:A:335:TYR:HB2	1.85	0.58
1:B:263:PRO:O	1:B:266:SER:OG	2.20	0.57
1:B:344:GLY:HA2	2:B:1345:RBS:C17	2.35	0.56
1:A:333:PRO:O	1:A:335:TYR:CD2	2.59	0.55
1:A:366:MET:HE2	1:A:400:CYS:SG	2.47	0.55
1:B:347:LEU:O	1:B:351:LYS:HG2	2.08	0.54
1:B:338:THR:HG22	1:B:339:GLU:O	2.08	0.53
2:A:1345:RBS:O61	2:A:1345:RBS:H8	2.07	0.53
1:B:262:ILE:HG12	1:B:327:ALA:HB1	1.91	0.52
1:A:428:TRP:HE1	1:A:454:GLU:CD	2.19	0.51
1:A:362:GLN:O	1:A:366:MET:HG3	2.11	0.51
1:A:312:GLN:O	1:A:316:LYS:HG3	2.10	0.50
1:A:320:GLU:O	1:A:401:LYS:HE2	2.11	0.50
1:A:323:VAL:HG11	1:A:393:LEU:HD12	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:338:THR:CG2	2:B:1345:RBS:H3	2.42	0.49
1:A:260:TRP:CD1	1:A:315:LYS:HG2	2.48	0.48
1:A:378:GLU:HG3	1:A:441:ILE:HG12	1.95	0.48
1:B:338:THR:HG21	2:B:1345:RBS:C3	2.42	0.47
1:A:262:ILE:HG12	1:A:327:ALA:HB1	1.95	0.47
1:A:262:ILE:HB	1:A:263:PRO:HD2	1.95	0.47
1:B:260:TRP:HH2	1:B:318:ARG:HH22	1.62	0.47
1:A:404:ASP:OD2	2:A:1345:RBS:H21	2.15	0.46
1:B:269:LEU:HB3	1:B:282:TRP:CE3	2.50	0.46
1:A:351:LYS:HE2	1:B:522:SER:CB	2.46	0.46
1:A:441:ILE:O	1:A:444:ASP:HB2	2.16	0.46
1:A:263:PRO:O	1:A:266:SER:OG	2.29	0.45
1:B:313:VAL:C	1:B:315:LYS:N	2.75	0.44
1:B:323:VAL:CG2	1:B:338:THR:HG23	2.48	0.44
1:A:518:ASP:O	1:A:519:TYR:C	2.60	0.44
1:B:524:GLU:C	1:B:526:GLN:H	2.26	0.43
1:A:309:GLN:CB	1:A:312:GLN:H	2.31	0.43
1:A:363:LEU:HD13	1:A:455:LEU:O	2.18	0.43
1:B:366:MET:HB3	1:B:400:CYS:SG	2.59	0.43
1:A:351:LYS:HE2	1:B:522:SER:HB3	2.01	0.42
1:B:280:GLU:O	1:B:295:LYS:HA	2.19	0.42
1:B:500:ARG:HD3	1:B:505:GLU:HB3	2.01	0.42
1:B:428:TRP:HE1	1:B:454:GLU:CD	2.27	0.42
1:A:388:ARG:HB3	1:A:428:TRP:CD1	2.54	0.42
1:B:313:VAL:C	1:B:315:LYS:H	2.27	0.42
1:A:372:SER:HB2	1:A:513:GLN:CD	2.45	0.41
1:B:440:THR:HG23	1:B:443:SER:N	2.36	0.41
1:B:453:THR:HG21	1:B:481:MET:HE3	2.03	0.41
1:B:265:GLU:H	1:B:265:GLU:CD	2.28	0.41
1:B:394:VAL:HA	1:B:399:VAL:O	2.21	0.41
1:A:339:GLU:OE2	1:A:401:LYS:NZ	2.43	0.41
2:A:1345:RBS:N3	2:A:1345:RBS:C4	2.74	0.41
1:B:491:LEU:HD23	1:B:491:LEU:HA	1.92	0.41
1:B:360:LEU:O	1:B:361:PRO:C	2.60	0.41
1:A:502:ASP:O	1:A:503:PRO:C	2.65	0.40
1:A:351:LYS:O	1:B:511:TYR:HE1	2.05	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/286 (84%)	221 (92%)	18 (8%)	2 (1%)	16	31
1	B	234/286 (82%)	220 (94%)	13 (6%)	1 (0%)	30	49
All	All	475/572 (83%)	441 (93%)	31 (6%)	3 (1%)	21	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	308	LEU
1	A	311	ALA
1	B	468	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/245 (76%)	174 (93%)	13 (7%)	14	29
1	B	193/245 (79%)	175 (91%)	18 (9%)	8	18
All	All	380/490 (78%)	349 (92%)	31 (8%)	10	23

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	295	LYS
1	A	323	VAL
1	A	328	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	339	GLU
1	A	351	LYS
1	A	353	GLU
1	A	356	LYS
1	A	360	LEU
1	A	372	SER
1	A	426	ILE
1	A	451	LEU
1	A	469	ARG
1	A	504	GLU
1	B	265	GLU
1	B	290	THR
1	B	322	LEU
1	B	323	VAL
1	B	334	ILE
1	B	338	THR
1	B	351	LYS
1	B	356	LYS
1	B	360	LEU
1	B	396	GLU
1	B	426	ILE
1	B	440	THR
1	B	467	VAL
1	B	470	GLU
1	B	471	VAL
1	B	491	LEU
1	B	528	GLN
1	B	531	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	381	ASN
1	A	532	ASN
1	B	381	ASN
1	B	397	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	B	100	-	5,5,5	0.72	0	5,5,5	0.65	0
2	RBS	A	1345	1	24,24,24	1.71	4 (16%)	31,32,32	2.72	9 (29%)
2	RBS	B	1345	1	24,24,24	2.02	5 (20%)	31,32,32	2.82	11 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	B	100	-	-	3/4/4/4	-
2	RBS	A	1345	1	-	2/10/10/10	0/3/3/3
2	RBS	B	1345	1	-	2/10/10/10	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1345	RBS	C51-C11	5.20	1.55	1.30
2	B	1345	RBS	C6-C7	-4.98	1.38	1.44
2	A	1345	RBS	C51-C11	4.88	1.53	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1345	RBS	C11-C10	3.88	1.54	1.48
2	A	1345	RBS	C11-C10	3.09	1.53	1.48
2	A	1345	RBS	C6-C7	-3.09	1.41	1.44
2	B	1345	RBS	C6-N3	2.25	1.37	1.34
2	A	1345	RBS	C9-N11	-2.13	1.37	1.41
2	B	1345	RBS	C5-N1	-2.08	1.36	1.40

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1345	RBS	C51-C11-C10	-8.50	112.88	122.22
2	A	1345	RBS	C19-N3-C6	6.20	121.41	116.60
2	B	1345	RBS	N2-C19-N3	-6.18	120.28	128.67
2	B	1345	RBS	C8-C7-C6	-5.66	120.24	124.84
2	B	1345	RBS	C19-N3-C6	5.62	120.96	116.60
2	A	1345	RBS	N2-C19-N3	-5.43	121.31	128.67
2	B	1345	RBS	C19-N2-C18	4.93	121.04	115.43
2	B	1345	RBS	N1-C6-N3	4.80	124.11	118.71
2	B	1345	RBS	C6-C7-C18	4.80	118.69	115.86
2	B	1345	RBS	C51-C11-C10	-4.63	117.13	122.22
2	A	1345	RBS	C6-C7-C18	4.39	118.45	115.86
2	A	1345	RBS	C19-N2-C18	4.35	120.38	115.43
2	B	1345	RBS	C11-C10-N11	3.76	116.39	113.84
2	A	1345	RBS	C7-C6-N3	-3.64	118.65	121.35
2	B	1345	RBS	C7-C6-N1	-3.20	116.52	119.77
2	A	1345	RBS	C8-C7-C6	-3.14	122.29	124.84
2	B	1345	RBS	C7-C18-N2	-2.99	119.65	122.82
2	A	1345	RBS	C7-C18-N2	-2.85	119.80	122.82
2	B	1345	RBS	C7-C6-N3	-2.68	119.37	121.35
2	A	1345	RBS	C11-C10-N11	2.42	115.48	113.84

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	1345	RBS	O61-C10-C11-C51
2	A	1345	RBS	N11-C10-C11-C51
2	B	1345	RBS	O61-C10-C11-C51
2	B	1345	RBS	N11-C10-C11-C51
3	B	100	GOL	C1-C2-C3-O3
3	B	100	GOL	O2-C2-C3-O3
3	B	100	GOL	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1345	RBS	4	0
2	B	1345	RBS	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	251/286 (87%)	0.54	26 (10%)	11 10	12, 25, 58, 65	3 (1%)
1	B	242/286 (84%)	0.35	11 (4%)	38 33	9, 26, 53, 65	0
All	All	493/572 (86%)	0.45	37 (7%)	20 18	9, 26, 57, 65	3 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	330	SER	5.3
1	A	277	CYS	5.2
1	A	287	ASN	4.1
1	A	276	GLY	4.1
1	A	307	PHE	4.1
1	A	278	PHE	4.1
1	B	312	GLN	3.6
1	A	333	PRO	3.5
1	B	298	LYS	3.3
1	A	308	LEU	3.2
1	B	425	PRO	2.9
1	A	312	GLN	2.9
1	A	262	ILE	2.8
1	A	406	GLY	2.7
1	A	271	VAL	2.6
1	B	313	VAL	2.5
1	A	299	PRO	2.5
1	A	335	TYR	2.5
1	B	260	TRP	2.4
1	A	269	LEU	2.4
1	A	286	TRP	2.4
1	A	297	LEU	2.3
1	A	502	ASP	2.3
1	A	314	MET	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	289	THR	2.3
1	A	267	LEU	2.2
1	A	274	GLY	2.2
1	B	269	LEU	2.2
1	B	275	GLN	2.2
1	B	258	ASP	2.2
1	B	332	GLU	2.1
1	A	310	GLU	2.1
1	B	276	GLY	2.1
1	A	290	THR	2.1
1	A	255	LEU	2.0
1	B	314	MET	2.0
1	A	425	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	RBS	B	1345	22/22	0.73	0.20	34,48,49,49	0
3	GOL	B	100	6/6	0.81	0.19	26,33,34,38	0
2	RBS	A	1345	22/22	0.86	0.14	29,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around RBS B 1345:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.