



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:32 PM UTC

PDB ID : 2HWP / pdb_00002hwp
Title : Crystal structure of Src kinase domain in complex with covalent inhibitor PD168393
Authors : Rauh, D.; Blair, J.A.; Shokat, K.M.
Deposited on : 2006-08-01
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	12% 70% 13% 14%
1	B	286	3% 65% 16% 16%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3763 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	245	Total	C	N	O	S	0	0	0
			1855	1196	313	330	16			
1	B	239	Total	C	N	O	S	0	0	0
			1832	1183	306	328	15			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	cloning artifact	UNP P00523
A	249	HIS	-	cloning artifact	UNP P00523
A	250	MET	-	cloning artifact	UNP P00523
A	345	CYS	SER	engineered mutation	UNP P00523
B	248	GLY	-	cloning artifact	UNP P00523
B	249	HIS	-	cloning artifact	UNP P00523
B	250	MET	-	cloning artifact	UNP P00523
B	345	CYS	SER	engineered mutation	UNP P00523

- Molecule 2 is N-[4-(3-BROMO-PHENYLAMINO)-QUINAZOLIN-6-YL]-ACRYLAMIDE (CCD ID: DJK) (formula: C₁₇H₁₃BrN₄O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 23	Br 1	C 17	N 4	O 1	0	0
2	B	1	Total 23	Br 1	C 17	N 4	O 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	15	Total O 15 15	0	0
3	B	15	Total O 15 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.06Å 63.01Å 73.69Å 100.74° 89.81° 90.20°	Depositor
Resolution (Å)	72.36 – 2.48 72.36 – 2.48	Depositor EDS
% Data completeness (in resolution range)	93.3 (72.36-2.48) 93.3 (72.36-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.57 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.225 , 0.276 0.221 , 0.272	Depositor DCC
R_{free} test set	1237 reflections (4.68%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.405	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 36.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3763	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DJK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/1897	0.95	2/2575 (0.1%)
1	B	0.74	0/1876	0.93	4/2554 (0.2%)
All	All	0.76	0/3773	0.94	6/5129 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	404	ASP	N-CA-C	8.06	122.70	112.86
1	A	273	LEU	N-CA-C	-6.48	105.00	114.39
1	B	430	ALA	CA-C-N	5.88	125.35	119.24
1	B	430	ALA	C-N-CA	5.88	125.35	119.24
1	B	461	VAL	CA-C-N	5.78	125.75	120.03
1	B	461	VAL	C-N-CA	5.78	125.75	120.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1855	0	1757	31	1
1	B	1832	0	1760	42	0
2	A	23	0	11	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	23	0	12	2	0
3	A	15	0	0	0	0
3	B	15	0	0	0	0
All	All	3763	0	3540	77	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:476:GLU:OE2	1:B:476:GLU:HA	1.33	1.13
1:A:476:GLU:HA	1:A:476:GLU:OE1	1.33	1.10
1:B:293:ALA:HB3	1:B:338:THR:HG22	1.50	0.94
1:A:476:GLU:OE1	1:A:476:GLU:CA	2.15	0.89
1:B:426:ILE:HD13	1:B:472:LEU:CD2	2.02	0.88
1:B:426:ILE:HD13	1:B:472:LEU:HD21	1.57	0.86
1:A:323:VAL:HG11	1:A:403:ALA:HB2	1.56	0.86
1:A:323:VAL:CG1	1:A:403:ALA:HB2	2.06	0.85
1:B:426:ILE:CD1	1:B:472:LEU:HD22	2.09	0.81
1:B:476:GLU:OE2	1:B:476:GLU:CA	2.21	0.78
1:B:424:PHE:N	1:B:425:PRO:HA	2.05	0.70
1:B:426:ILE:CD1	1:B:472:LEU:CD2	2.69	0.70
1:A:279:GLY:O	1:A:280:GLU:CB	2.39	0.70
1:B:424:PHE:N	1:B:426:ILE:H	1.93	0.67
1:A:287:ASN:O	1:A:289:THR:HG22	1.95	0.67
1:B:426:ILE:HD11	1:B:472:LEU:HD22	1.75	0.66
1:A:518:ASP:CG	1:B:461:VAL:HG23	2.21	0.66
1:A:257:LYS:CB	1:A:261:GLU:HG3	2.27	0.65
1:A:308:LEU:O	1:A:311:ALA:N	2.29	0.65
2:A:1345:DJK:N3	2:A:1345:DJK:H4	2.14	0.62
1:A:386:ASP:CB	1:A:388:ARG:NH1	2.63	0.61
1:B:424:PHE:N	1:B:425:PRO:CA	2.65	0.59
1:B:388:ARG:HB3	1:B:428:TRP:CD1	2.36	0.59
1:B:283:MET:HG3	1:B:340:TYR:CE1	2.38	0.58
2:B:1345:DJK:H4	2:B:1345:DJK:N3	2.19	0.57
2:B:1345:DJK:O61	2:B:1345:DJK:H8	2.04	0.57
1:A:386:ASP:CB	1:A:388:ARG:HH12	2.18	0.56
1:A:431:PRO:O	1:A:433:ALA:N	2.39	0.56
1:A:323:VAL:HG13	1:A:403:ALA:HB2	1.86	0.56
1:B:493:ASP:OD2	1:B:497:GLN:NE2	2.38	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:MET:HG2	1:A:405:PHE:CZ	2.42	0.55
1:A:314:MET:SD	1:A:405:PHE:CE1	3.00	0.54
1:A:308:LEU:O	1:A:309:GLN:C	2.49	0.53
1:B:283:MET:HG3	1:B:340:TYR:CZ	2.43	0.53
1:B:323:VAL:HG11	1:B:393:LEU:HD12	1.91	0.52
1:B:501:LYS:O	1:B:503:PRO:HD3	2.09	0.52
1:A:486:GLU:OE2	1:A:532:ASN:ND2	2.43	0.51
1:B:321:LYS:HA	1:B:401:LYS:HG2	1.93	0.51
1:B:283:MET:SD	1:B:291:ARG:NH1	2.85	0.49
1:A:431:PRO:C	1:A:433:ALA:H	2.21	0.49
1:B:388:ARG:CB	1:B:428:TRP:CD1	2.95	0.49
1:A:503:PRO:O	1:A:506:ARG:HB2	2.14	0.48
1:A:431:PRO:C	1:A:433:ALA:N	2.72	0.47
1:B:440:THR:HG23	1:B:443:SER:OG	2.14	0.47
1:A:404:ASP:OD1	2:A:1345:DJK:C20	2.64	0.46
1:A:501:LYS:O	1:A:503:PRO:HD3	2.15	0.45
1:B:424:PHE:N	1:B:426:ILE:N	2.63	0.45
1:A:289:THR:HG23	1:A:290:THR:OG1	2.16	0.45
2:A:1345:DJK:N3	2:A:1345:DJK:C4	2.79	0.45
1:B:493:ASP:O	1:B:497:GLN:HG3	2.16	0.45
2:A:1345:DJK:H8	2:A:1345:DJK:O61	2.17	0.45
1:B:502:ASP:O	1:B:503:PRO:C	2.60	0.45
1:B:424:PHE:N	1:B:426:ILE:HG22	2.32	0.44
1:B:343:LYS:HB2	1:B:394:VAL:HB	1.99	0.44
1:B:461:VAL:HA	1:B:462:PRO:HD3	1.85	0.43
1:B:518:ASP:O	1:B:519:TYR:C	2.61	0.42
1:A:388:ARG:HB3	1:A:428:TRP:CD1	2.54	0.42
1:A:308:LEU:O	1:A:309:GLN:O	2.38	0.42
1:B:329:VAL:HB	1:B:335:TYR:HB2	2.02	0.42
1:B:375:ALA:O	1:B:378:GLU:HB3	2.19	0.42
1:B:321:LYS:HD2	1:B:369:GLN:HB3	2.00	0.42
1:B:338:THR:HG23	1:B:339:GLU:O	2.19	0.42
1:B:503:PRO:O	1:B:506:ARG:HB2	2.19	0.42
1:A:452:LEU:HD23	1:A:452:LEU:HA	1.95	0.41
1:A:289:THR:HG23	1:A:290:THR:HG1	1.86	0.41
1:A:372:SER:HB2	1:A:513:GLN:NE2	2.35	0.41
1:B:354:MET:O	1:B:355:GLY:C	2.61	0.41
1:B:463:TYR:O	1:B:464:PRO:C	2.63	0.41
1:B:511:TYR:CD2	1:B:511:TYR:C	2.98	0.41
1:A:442:LYS:NZ	1:A:506:ARG:O	2.47	0.41
1:B:359:ARG:O	1:B:360:LEU:C	2.64	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:440:THR:OG1	1:B:442:LYS:N	2.51	0.41
1:B:485:PRO:O	1:B:486:GLU:HB2	2.21	0.41
1:A:372:SER:HB2	1:A:513:GLN:CD	2.45	0.41
1:B:269:LEU:HB3	1:B:282:TRP:CE3	2.56	0.40
1:A:258:ASP:C	1:A:258:ASP:OD2	2.65	0.40
1:B:283:MET:HE2	1:B:340:TYR:OH	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397:ASN:ND2	1:A:438:ARG:CD[1_455]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/286 (81%)	221 (96%)	6 (3%)	4 (2%)	7	12
1	B	231/286 (81%)	221 (96%)	10 (4%)	0	100	100
All	All	462/572 (81%)	442 (96%)	16 (4%)	4 (1%)	14	25

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	280	GLU
1	A	432	GLU
1	A	308	LEU
1	A	431	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	182/245 (74%)	169 (93%)	13 (7%)	13	26
1	B	185/245 (76%)	173 (94%)	12 (6%)	15	30
All	All	367/490 (75%)	342 (93%)	25 (7%)	14	28

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	277	CYS
1	A	290	THR
1	A	291	ARG
1	A	323	VAL
1	A	328	VAL
1	A	356	LYS
1	A	360	LEU
1	A	426	ILE
1	A	438	ARG
1	A	467	VAL
1	A	476	GLU
1	A	491	LEU
1	A	528	GLN
1	B	323	VAL
1	B	338	THR
1	B	356	LYS
1	B	360	LEU
1	B	426	ILE
1	B	440	THR
1	B	451	LEU
1	B	467	VAL
1	B	472	LEU
1	B	476	GLU
1	B	491	LEU
1	B	528	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	381	ASN
1	A	397	ASN
1	B	397	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DJK	B	1345	1	25,25,25	1.68	3 (12%)	33,34,34	2.54	10 (30%)
2	DJK	A	1345	1	25,25,25	1.67	4 (16%)	33,34,34	2.71	10 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DJK	B	1345	1	-	2/10/10/10	0/3/3/3
2	DJK	A	1345	1	-	2/10/10/10	0/3/3/3

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1345	DJK	C51-C11	5.12	1.55	1.30
2	A	1345	DJK	C51-C11	4.90	1.53	1.30
2	A	1345	DJK	C6-C7	-2.93	1.41	1.44
2	A	1345	DJK	C11-C10	2.91	1.52	1.48
2	B	1345	DJK	C11-C10	2.78	1.52	1.48
2	B	1345	DJK	C6-C7	-2.55	1.41	1.44
2	A	1345	DJK	C8-C9	2.05	1.41	1.37

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1345	DJK	C51-C11-C10	-8.38	113.01	122.22
2	B	1345	DJK	C51-C11-C10	-6.50	115.07	122.22
2	B	1345	DJK	C11-C10-N11	5.90	117.84	113.84
2	A	1345	DJK	C11-C10-N11	5.88	117.83	113.84
2	A	1345	DJK	C19-N3-C6	5.33	120.74	116.60
2	A	1345	DJK	N2-C19-N3	-5.27	121.52	128.67
2	B	1345	DJK	C6-C7-C18	4.83	118.71	115.86
2	B	1345	DJK	N2-C19-N3	-4.55	122.50	128.67
2	A	1345	DJK	C19-N2-C18	4.34	120.36	115.43
2	B	1345	DJK	C19-N3-C6	4.15	119.82	116.60
2	B	1345	DJK	C19-N2-C18	4.07	120.06	115.43
2	A	1345	DJK	C6-C7-C18	4.03	118.24	115.86
2	B	1345	DJK	C8-C7-C6	-3.34	122.12	124.84
2	B	1345	DJK	C7-C18-N2	-3.12	119.52	122.82
2	A	1345	DJK	C7-C18-N2	-2.82	119.84	122.82
2	A	1345	DJK	C7-C6-N3	-2.77	119.30	121.35
2	B	1345	DJK	C7-C6-N3	-2.65	119.39	121.35
2	A	1345	DJK	C8-C7-C6	-2.52	122.79	124.84
2	A	1345	DJK	O61-C10-C11	-2.44	119.03	122.70
2	B	1345	DJK	BR-C3-C4	2.34	122.55	119.23

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1345	DJK	O61-C10-C11-C51
2	A	1345	DJK	N11-C10-C11-C51
2	B	1345	DJK	O61-C10-C11-C51
2	B	1345	DJK	N11-C10-C11-C51


There are no ring outliers.

2 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1345	DJK	2	0
2	A	1345	DJK	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand DJK B 1345



Ligand DJK A 1345



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	245/286 (85%)	0.41	34 (13%) 6 5	10, 26, 62, 83	1 (0%)
1	B	239/286 (83%)	0.17	8 (3%) 49 45	11, 26, 53, 64	0
All	All	484/572 (84%)	0.29	42 (8%) 16 14	10, 26, 60, 83	1 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	271	VAL	5.5
1	A	308	LEU	5.4
1	A	309	GLN	5.3
1	A	424	PHE	5.2
1	A	307	PHE	4.7
1	A	331	GLU	4.2
1	A	405	PHE	4.1
1	B	312	GLN	4.0
1	B	258	ASP	3.6
1	A	276	GLY	3.5
1	A	268	ARG	3.5
1	A	273	LEU	3.5
1	A	279	GLY	3.3
1	A	256	ALA	3.2
1	A	311	ALA	3.2
1	A	277	CYS	3.2
1	A	335	TYR	3.1
1	A	264	ARG	3.0
1	A	314	MET	2.9
1	A	267	LEU	2.9
1	A	280	GLU	2.9
1	B	424	PHE	2.9
1	B	279	GLY	2.8
1	A	283	MET	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	275	GLN	2.6
1	B	275	GLN	2.6
1	A	282	TRP	2.5
1	A	333	PRO	2.4
1	A	281	VAL	2.4
1	A	272	LYS	2.4
1	A	286	TRP	2.3
1	A	329	VAL	2.3
1	A	287	ASN	2.3
1	A	266	SER	2.3
1	A	404	ASP	2.3
1	B	315	LYS	2.3
1	A	312	GLN	2.2
1	A	288	GLY	2.2
1	B	354	MET	2.2
1	A	262	ILE	2.2
1	A	265	GLU	2.2
1	B	317	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	DJK	A	1345	23/23	0.81	0.16	31,42,42,42	1
2	DJK	B	1345	23/23	0.83	0.14	35,42,43,43	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around DJK A 1345:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around DJK B 1345:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.