



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 04:10 PM UTC

PDB ID : 6HWM / pdb_00006hwm
Title : Structure of Thermus thermophilus ClpP in complex with bortezomib
Authors : Felix, J.; Schanda, P.; Fraga, H.; Morlot, C.
Deposited on : 2018-10-12
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	204	 2% 72% 19% 9%
1	B	204	 2% 73% 21% 6%
1	C	204	 2% 67% 23% 10%
1	D	204	 2% 68% 22% 10%
1	E	204	 2% 74% 22% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	204	 2% 69% 23% 7%
1	G	204	 % 68% 22% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	BO2	C	301	-	-	X	-
2	BO2	F	301	-	-	X	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10130 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent Clp protease proteolytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	185	Total	C	N	O	S	0	0	0
			1417	906	237	270	4			
1	B	192	Total	C	N	O	S	0	0	0
			1452	929	247	272	4			
1	C	183	Total	C	N	O	S	0	0	0
			1374	883	227	260	4			
1	D	184	Total	C	N	O	S	0	0	0
			1393	893	231	265	4			
1	E	195	Total	C	N	O	S	0	0	0
			1462	937	246	275	4			
1	F	189	Total	C	N	O	S	0	0	0
			1429	917	240	268	4			
1	G	183	Total	C	N	O	S	0	0	0
			1386	890	230	262	4			

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	LEU	-	expression tag	UNP Q5SKM8
A	196	GLU	-	expression tag	UNP Q5SKM8
A	197	HIS	-	expression tag	UNP Q5SKM8
A	198	HIS	-	expression tag	UNP Q5SKM8
A	199	HIS	-	expression tag	UNP Q5SKM8
A	200	HIS	-	expression tag	UNP Q5SKM8
A	201	HIS	-	expression tag	UNP Q5SKM8
A	202	HIS	-	expression tag	UNP Q5SKM8
A	203	HIS	-	expression tag	UNP Q5SKM8
A	204	HIS	-	expression tag	UNP Q5SKM8
B	195	LEU	-	expression tag	UNP Q5SKM8
B	196	GLU	-	expression tag	UNP Q5SKM8
B	197	HIS	-	expression tag	UNP Q5SKM8
B	198	HIS	-	expression tag	UNP Q5SKM8
B	199	HIS	-	expression tag	UNP Q5SKM8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	200	HIS	-	expression tag	UNP Q5SKM8
B	201	HIS	-	expression tag	UNP Q5SKM8
B	202	HIS	-	expression tag	UNP Q5SKM8
B	203	HIS	-	expression tag	UNP Q5SKM8
B	204	HIS	-	expression tag	UNP Q5SKM8
C	195	LEU	-	expression tag	UNP Q5SKM8
C	196	GLU	-	expression tag	UNP Q5SKM8
C	197	HIS	-	expression tag	UNP Q5SKM8
C	198	HIS	-	expression tag	UNP Q5SKM8
C	199	HIS	-	expression tag	UNP Q5SKM8
C	200	HIS	-	expression tag	UNP Q5SKM8
C	201	HIS	-	expression tag	UNP Q5SKM8
C	202	HIS	-	expression tag	UNP Q5SKM8
C	203	HIS	-	expression tag	UNP Q5SKM8
C	204	HIS	-	expression tag	UNP Q5SKM8
D	195	LEU	-	expression tag	UNP Q5SKM8
D	196	GLU	-	expression tag	UNP Q5SKM8
D	197	HIS	-	expression tag	UNP Q5SKM8
D	198	HIS	-	expression tag	UNP Q5SKM8
D	199	HIS	-	expression tag	UNP Q5SKM8
D	200	HIS	-	expression tag	UNP Q5SKM8
D	201	HIS	-	expression tag	UNP Q5SKM8
D	202	HIS	-	expression tag	UNP Q5SKM8
D	203	HIS	-	expression tag	UNP Q5SKM8
D	204	HIS	-	expression tag	UNP Q5SKM8
E	195	LEU	-	expression tag	UNP Q5SKM8
E	196	GLU	-	expression tag	UNP Q5SKM8
E	197	HIS	-	expression tag	UNP Q5SKM8
E	198	HIS	-	expression tag	UNP Q5SKM8
E	199	HIS	-	expression tag	UNP Q5SKM8
E	200	HIS	-	expression tag	UNP Q5SKM8
E	201	HIS	-	expression tag	UNP Q5SKM8
E	202	HIS	-	expression tag	UNP Q5SKM8
E	203	HIS	-	expression tag	UNP Q5SKM8
E	204	HIS	-	expression tag	UNP Q5SKM8
F	195	LEU	-	expression tag	UNP Q5SKM8
F	196	GLU	-	expression tag	UNP Q5SKM8
F	197	HIS	-	expression tag	UNP Q5SKM8
F	198	HIS	-	expression tag	UNP Q5SKM8
F	199	HIS	-	expression tag	UNP Q5SKM8
F	200	HIS	-	expression tag	UNP Q5SKM8
F	201	HIS	-	expression tag	UNP Q5SKM8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	202	HIS	-	expression tag	UNP Q5SKM8
F	203	HIS	-	expression tag	UNP Q5SKM8
F	204	HIS	-	expression tag	UNP Q5SKM8
G	195	LEU	-	expression tag	UNP Q5SKM8
G	196	GLU	-	expression tag	UNP Q5SKM8
G	197	HIS	-	expression tag	UNP Q5SKM8
G	198	HIS	-	expression tag	UNP Q5SKM8
G	199	HIS	-	expression tag	UNP Q5SKM8
G	200	HIS	-	expression tag	UNP Q5SKM8
G	201	HIS	-	expression tag	UNP Q5SKM8
G	202	HIS	-	expression tag	UNP Q5SKM8
G	203	HIS	-	expression tag	UNP Q5SKM8
G	204	HIS	-	expression tag	UNP Q5SKM8

- Molecule 2 is N-[(1R)-1-(DIHYDROXYBORYL)-3-METHYLBUTYL]-N-(PYRAZIN-2-YL CARBONYL)-L-PHENYLALANINAMIDE (CCD ID: BO2) (formula: C₁₉H₂₅BN₄O₄).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 28	B 1	C 19	N 4	O 4	0	0
2	B	1	Total 28	B 1	C 19	N 4	O 4	0	0
2	C	1	Total 28	B 1	C 19	N 4	O 4	0	0
2	D	1	Total 28	B 1	C 19	N 4	O 4	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	E	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
2	F	1	Total	B	C	N	O	0	0
			28	1	19	4	4		
2	G	1	Total	B	C	N	O	0	0
			28	1	19	4	4		



- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			7	4	3		
3	E	1	Total	C	O	0	0
			7	4	3		
3	G	1	Total	C	O	0	0
			7	4	3		




- Molecule 1: ATP-dependent Clp protease proteolytic subunit



- Molecule 1: ATP-dependent Clp protease proteolytic subunit



- Molecule 1: ATP-dependent Clp protease proteolytic subunit



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	135.14Å 168.74Å 166.08Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.29 – 2.70 46.29 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.29-2.70) 100.0 (46.29-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 2.69Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.205 , 0.236 0.205 , 0.238	Depositor DCC
R_{free} test set	5230 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	92.8	Xtriage
Anisotropy	0.407	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	10130	wwPDB-VP
Average B, all atoms (Å ²)	103.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: BO2, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	1/1440 (0.1%)	0.91	2/1958 (0.1%)
1	B	0.63	0/1480	0.83	2/2017 (0.1%)
1	C	0.58	0/1397	0.77	0/1904
1	D	0.54	0/1416	0.74	0/1928
1	E	0.58	0/1488	0.81	4/2029 (0.2%)
1	F	0.68	2/1452 (0.1%)	0.80	1/1976 (0.1%)
1	G	0.54	0/1409	0.76	0/1919
All	All	0.60	3/10082 (0.0%)	0.81	9/13731 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	166	LYS	C-O	-5.18	1.18	1.24
1	F	184	LEU	C-O	-5.15	1.17	1.24
1	A	155	LYS	N-CA	-5.01	1.40	1.46

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	3	ILE	CA-C-N	12.56	132.46	120.03
1	A	3	ILE	C-N-CA	12.56	132.46	120.03
1	F	194	ALA	N-CA-C	-5.64	103.92	111.87
1	B	161	LEU	CA-C-N	-5.49	111.56	120.72
1	B	161	LEU	C-N-CA	-5.49	111.56	120.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1417	0	1414	35	0
1	B	1452	0	1418	40	0
1	C	1374	0	1362	55	0
1	D	1393	0	1383	42	0
1	E	1462	0	1433	44	0
1	F	1429	0	1430	54	0
1	G	1386	0	1384	41	0
2	A	28	0	25	3	0
2	B	28	0	25	6	0
2	C	28	0	25	11	0
2	D	28	0	25	4	0
2	E	28	0	25	6	0
2	F	28	0	25	9	0
2	G	28	0	25	7	0
3	C	7	0	10	0	0
3	E	7	0	10	0	0
3	G	7	0	10	1	0
All	All	10130	0	10029	283	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 283 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:GLU:HG2	1:A:110:ARG:NH2	1.44	1.31
1:A:107:GLU:HG2	1:A:110:ARG:HH22	1.03	0.97
1:A:107:GLU:CG	1:A:110:ARG:NH2	2.30	0.94
1:A:107:GLU:CG	1:A:110:ARG:HH22	1.82	0.92
1:F:3:ILE:HD12	1:F:3:ILE:O	1.70	0.91

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/204 (89%)	179 (99%)	2 (1%)	0	100	100
1	B	188/204 (92%)	186 (99%)	2 (1%)	0	100	100
1	C	179/204 (88%)	177 (99%)	2 (1%)	0	100	100
1	D	180/204 (88%)	176 (98%)	4 (2%)	0	100	100
1	E	191/204 (94%)	188 (98%)	3 (2%)	0	100	100
1	F	185/204 (91%)	180 (97%)	5 (3%)	0	100	100
1	G	179/204 (88%)	177 (99%)	2 (1%)	0	100	100
All	All	1283/1428 (90%)	1263 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/167 (87%)	145 (99%)	1 (1%)	76	90
1	B	146/167 (87%)	146 (100%)	0	100	100
1	C	139/167 (83%)	139 (100%)	0	100	100
1	D	142/167 (85%)	140 (99%)	2 (1%)	59	82
1	E	146/167 (87%)	146 (100%)	0	100	100
1	F	145/167 (87%)	145 (100%)	0	100	100
1	G	142/167 (85%)	142 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1006/1169 (86%)	1003 (100%)	3 (0%)	86 94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ILE
1	D	191	ARG
1	D	193	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	201	HIS
1	F	46	GLN
1	G	46	GLN
1	C	46	GLN
1	D	46	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BO2	E	301	1	25,29,29	1.63	3 (12%)	33,38,38	1.22	4 (12%)
2	BO2	F	301	1	25,29,29	1.72	2 (8%)	33,38,38	1.16	4 (12%)
3	PEG	G	302	-	6,6,6	0.66	0	5,5,5	0.35	0
2	BO2	A	301	1	25,29,29	1.53	3 (12%)	33,38,38	1.57	7 (21%)
2	BO2	C	301	1	25,29,29	1.70	2 (8%)	33,38,38	1.42	5 (15%)
2	BO2	G	301	1	25,29,29	1.60	4 (16%)	33,38,38	1.32	4 (12%)
2	BO2	B	301	1	25,29,29	1.69	3 (12%)	33,38,38	1.67	9 (27%)
2	BO2	D	301	1	25,29,29	1.59	2 (8%)	33,38,38	1.53	5 (15%)
3	PEG	C	302	-	6,6,6	0.71	0	5,5,5	0.79	0
3	PEG	E	302	-	6,6,6	0.70	0	5,5,5	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BO2	E	301	1	-	8/23/28/28	0/2/2/2
2	BO2	F	301	1	-	8/23/28/28	0/2/2/2
3	PEG	G	302	-	-	2/4/4/4	-
2	BO2	A	301	1	-	10/23/28/28	0/2/2/2
2	BO2	C	301	1	-	6/23/28/28	0/2/2/2
2	BO2	G	301	1	-	9/23/28/28	0/2/2/2
2	BO2	B	301	1	-	8/23/28/28	0/2/2/2
2	BO2	D	301	1	-	5/23/28/28	0/2/2/2
3	PEG	C	302	-	-	3/4/4/4	-
3	PEG	E	302	-	-	3/4/4/4	-

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	BO2	C18-N20	5.70	1.46	1.34
2	F	301	BO2	C18-N20	5.58	1.46	1.34
2	C	301	BO2	C18-N20	5.54	1.45	1.34
2	E	301	BO2	C18-N20	5.53	1.45	1.34
2	D	301	BO2	C7-N9	5.32	1.46	1.34

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	BO2	C2-C7-N9	4.01	122.89	115.19
2	G	301	BO2	B26-C21-C22	3.98	119.83	112.31
2	D	301	BO2	B26-C21-C22	3.95	119.77	112.31
2	A	301	BO2	B26-C21-C22	3.61	119.12	112.31
2	B	301	BO2	C7-C2-N1	3.26	121.23	117.42

There are no chirality outliers.

5 of 62 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	BO2	C3-C2-C7-O8
2	A	301	BO2	C3-C2-C7-N9
2	A	301	BO2	C21-C22-C23-C24
2	A	301	BO2	C21-C22-C23-C25
2	B	301	BO2	C3-C2-C7-O8


There are no ring outliers.

8 monomers are involved in 47 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	301	BO2	6	0
2	F	301	BO2	9	0
3	G	302	PEG	1	0
2	A	301	BO2	3	0
2	C	301	BO2	11	0
2	G	301	BO2	7	0
2	B	301	BO2	6	0
2	D	301	BO2	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand BO2 E 301




Bond lengths



Bond angles




Torsions




Rings


Ligand BO2 F 301




Bond lengths



Bond angles




Torsions




Rings


Ligand BO2 A 301




Bond lengths



Bond angles




Torsions




Rings


Ligand BO2 C 301




Bond lengths



Bond angles




Torsions




Rings


Ligand BO2 G 301




Bond lengths



Bond angles




Torsions




Rings


Ligand BO2 B 301




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	185/204 (90%)	0.00	5 (2%) 56 53	81, 103, 145, 183	0
1	B	192/204 (94%)	-0.02	5 (2%) 57 54	75, 105, 158, 226	0
1	C	183/204 (89%)	0.07	4 (2%) 62 59	79, 101, 134, 181	0
1	D	184/204 (90%)	0.04	5 (2%) 56 53	73, 95, 133, 171	0
1	E	195/204 (95%)	0.10	5 (2%) 57 54	74, 93, 156, 191	0
1	F	189/204 (92%)	-0.08	5 (2%) 57 54	66, 88, 129, 162	0
1	G	183/204 (89%)	-0.07	3 (1%) 70 68	73, 97, 130, 167	0
All	All	1311/1428 (91%)	0.01	32 (2%) 59 56	66, 97, 145, 226	0

The worst 5 of 32 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	16	VAL	4.3
1	D	2	VAL	3.7
1	F	118	LYS	3.7
1	C	2	VAL	3.5
1	G	16	VAL	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PEG	E	302	7/7	0.78	0.14	106,110,131,132	0
3	PEG	G	302	7/7	0.78	0.13	107,111,117,119	0
3	PEG	C	302	7/7	0.79	0.10	104,112,119,119	0
2	BO2	F	301	28/28	0.88	0.13	102,123,144,148	0
2	BO2	A	301	28/28	0.88	0.12	94,112,129,142	0
2	BO2	D	301	28/28	0.89	0.13	86,117,141,148	0
2	BO2	B	301	28/28	0.90	0.13	98,115,135,138	0
2	BO2	G	301	28/28	0.91	0.12	93,116,149,152	0
2	BO2	E	301	28/28	0.91	0.12	82,119,138,141	0
2	BO2	C	301	28/28	0.93	0.10	91,111,126,130	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BO2 F 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BO2 A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BO2 D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around BO2 B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around BO2 G 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around BO2 E 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around BO2 C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.