



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:44 AM UTC

PDB ID : 2HXI / pdb_00002hxi
Title : Structural Genomics, the crystal structure of a putative transcriptional regulator from *Streptomyces coelicolor* A3(2)
Authors : Tan, K.; Xu, X.; Zheng, H.; Savchenko, A.; Edwards, A.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2006-08-03
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	4% 68% 12% • 18%
1	B	241	4% 69% 11% • 18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	197	Total	C	N	O	Se	0	0	0
			1515	952	282	278	3			
1	B	198	Total	C	N	O	Se	0	0	0
			1526	958	287	278	3			

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	expression tag	UNP O87854
A	-20	GLY	-	expression tag	UNP O87854
A	-19	SER	-	expression tag	UNP O87854
A	-18	SER	-	expression tag	UNP O87854
A	-17	HIS	-	expression tag	UNP O87854
A	-16	HIS	-	expression tag	UNP O87854
A	-15	HIS	-	expression tag	UNP O87854
A	-14	HIS	-	expression tag	UNP O87854
A	-13	HIS	-	expression tag	UNP O87854
A	-12	HIS	-	expression tag	UNP O87854
A	-11	SER	-	expression tag	UNP O87854
A	-10	SER	-	expression tag	UNP O87854
A	-9	GLY	-	expression tag	UNP O87854
A	-8	ARG	-	expression tag	UNP O87854
A	-7	GLU	-	expression tag	UNP O87854
A	-6	ASN	-	expression tag	UNP O87854
A	-5	LEU	-	expression tag	UNP O87854
A	-4	TYR	-	expression tag	UNP O87854
A	-3	PHE	-	expression tag	UNP O87854
A	-2	GLN	-	expression tag	UNP O87854
A	-1	GLY	-	expression tag	UNP O87854
A	0	HIS	-	expression tag	UNP O87854
A	67	MSE	MET	modified residue	UNP O87854
A	114	MSE	MET	modified residue	UNP O87854
A	115	MSE	MET	modified residue	UNP O87854

Continued on next page...



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	218	GLY	-	cloning artifact	UNP O87854
A	219	SER	-	cloning artifact	UNP O87854
B	-21	MET	-	expression tag	UNP O87854
B	-20	GLY	-	expression tag	UNP O87854
B	-19	SER	-	expression tag	UNP O87854
B	-18	SER	-	expression tag	UNP O87854
B	-17	HIS	-	expression tag	UNP O87854
B	-16	HIS	-	expression tag	UNP O87854
B	-15	HIS	-	expression tag	UNP O87854
B	-14	HIS	-	expression tag	UNP O87854
B	-13	HIS	-	expression tag	UNP O87854
B	-12	HIS	-	expression tag	UNP O87854
B	-11	SER	-	expression tag	UNP O87854
B	-10	SER	-	expression tag	UNP O87854
B	-9	GLY	-	expression tag	UNP O87854
B	-8	ARG	-	expression tag	UNP O87854
B	-7	GLU	-	expression tag	UNP O87854
B	-6	ASN	-	expression tag	UNP O87854
B	-5	LEU	-	expression tag	UNP O87854
B	-4	TYR	-	expression tag	UNP O87854
B	-3	PHE	-	expression tag	UNP O87854
B	-2	GLN	-	expression tag	UNP O87854
B	-1	GLY	-	expression tag	UNP O87854
B	0	HIS	-	expression tag	UNP O87854
B	67	MSE	MET	modified residue	UNP O87854
B	114	MSE	MET	modified residue	UNP O87854
B	115	MSE	MET	modified residue	UNP O87854
B	218	GLY	-	cloning artifact	UNP O87854
B	219	SER	-	cloning artifact	UNP O87854

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	230	Total O 230 230	0	0
2	B	212	Total O 212 212	0	0

- Molecule 1: Putative transcriptional regulator



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.34Å 94.87Å 51.84Å 90.00° 95.61° 90.00°	Depositor
Resolution (Å)	22.61 – 1.70 22.61 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.1 (22.61-1.70) 98.4 (22.61-1.70)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.187 , 0.232 0.188 , 0.194	Depositor DCC
R_{free} test set	2345 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3483	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.23	2/1540 (0.1%)	1.16	3/2077 (0.1%)
1	B	1.19	0/1550	1.16	1/2088 (0.0%)
All	All	1.21	2/3090 (0.1%)	1.16	4/4165 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	57	ALA	CA-CB	-7.01	1.42	1.53
1	A	182	ALA	CA-CB	5.36	1.61	1.53

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	7	TRP	N-CA-C	7.93	122.22	110.48
1	A	107	GLY	N-CA-C	6.31	123.73	115.40
1	A	22	ALA	N-CA-C	5.33	118.41	110.14
1	B	6	ARG	N-CA-C	5.30	117.45	109.23

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	21	LEU	Peptide
1	A	6	ARG	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1515	0	1516	21	0
1	B	1526	0	1530	23	0
2	A	230	0	0	13	0
2	B	212	0	0	5	0
All	All	3483	0	3046	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:HIS:HD2	1:A:138:ARG:HH11	1.29	0.81
1:A:137:HIS:CD2	1:A:138:ARG:HH11	2.02	0.78
1:B:5:ARG:O	1:B:5:ARG:HD2	1.90	0.72
1:A:22:ALA:HA	1:A:25:ALA:HB2	1.71	0.71
1:B:167:GLN:HG3	1:B:168:PHE:H	1.60	0.66
1:B:102:GLY:HA3	1:B:148:THR:HG21	1.77	0.66
1:A:21:LEU:HD12	2:A:398:HOH:O	1.95	0.65
1:B:170:VAL:HG13	1:B:173:LEU:HG	1.80	0.64
1:B:23:GLY:HA2	1:B:27:THR:HG21	1.82	0.61
1:B:170:VAL:CG1	1:B:173:LEU:HG	2.31	0.60
1:A:88:ARG:NH1	2:A:448:HOH:O	2.30	0.60
1:B:54:LEU:C	1:B:54:LEU:HD23	2.29	0.58
1:A:21:LEU:CD1	2:A:398:HOH:O	2.52	0.57
1:B:52:THR:HG23	2:B:378:HOH:O	2.05	0.57
1:A:17:ALA:O	1:A:21:LEU:HD13	2.05	0.56
1:A:30:VAL:HG11	1:A:45:TYR:CE1	2.42	0.55
1:B:22:ALA:HB3	2:B:383:HOH:O	2.05	0.55
1:B:170:VAL:HG22	2:B:248:HOH:O	2.05	0.55
1:A:48:PHE:O	1:A:49:ARG:HB2	2.06	0.55
1:B:98:ALA:O	1:B:148:THR:HG22	2.07	0.54
1:A:191:ARG:NH1	2:A:440:HOH:O	2.41	0.54
1:B:23:GLY:CA	1:B:27:THR:HG21	2.38	0.54
1:A:88:ARG:NH2	1:A:150:GLU:OE2	2.31	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:9:THR:OG1	1:B:47:HIS:HD2	1.92	0.53
1:A:185:HIS:HE1	2:A:341:HOH:O	1.92	0.53
1:A:96:GLN:HB3	2:A:345:HOH:O	2.09	0.52
1:A:89:GLU:HG2	2:A:338:HOH:O	2.09	0.52
2:A:369:HOH:O	1:B:169:ARG:HD3	2.11	0.51
1:A:27:THR:HG23	2:A:383:HOH:O	2.10	0.50
1:B:27:THR:HG23	1:B:28:PHE:N	2.27	0.49
1:B:13:LEU:HB3	1:B:61:ARG:HG3	1.95	0.47
1:A:88:ARG:HD3	2:A:448:HOH:O	2.15	0.47
1:A:51:LYS:NZ	2:A:417:HOH:O	2.48	0.46
1:A:21:LEU:HD22	2:A:312:HOH:O	2.16	0.46
1:A:217:PRO:HD3	2:A:324:HOH:O	2.16	0.45
1:B:65:SER:O	1:B:68:ASP:HB2	2.17	0.44
1:A:137:HIS:CD2	1:A:138:ARG:NH1	2.79	0.44
1:A:150:GLU:O	1:A:151:GLY:C	2.59	0.44
1:B:89:GLU:HG2	2:B:279:HOH:O	2.17	0.44
1:B:54:LEU:O	1:B:58:VAL:HG23	2.19	0.43
1:B:144:SER:O	1:B:148:THR:HG23	2.20	0.42
1:B:24:ASP:HA	1:B:25:ALA:HA	1.73	0.42
1:B:21:LEU:HD21	2:B:352:HOH:O	2.20	0.41
1:B:27:THR:CG2	1:B:28:PHE:N	2.84	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/241 (80%)	187 (97%)	4 (2%)	2 (1%)	12	3
1	B	192/241 (80%)	190 (99%)	1 (0%)	1 (0%)	24	12
All	All	385/482 (80%)	377 (98%)	5 (1%)	3 (1%)	16	5

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	21	LEU
1	A	23	GLY
1	A	25	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/180 (83%)	143 (95%)	7 (5%)	23	9
1	B	151/180 (84%)	145 (96%)	6 (4%)	28	12
All	All	301/360 (84%)	288 (96%)	13 (4%)	26	10

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	ASP
1	A	27	THR
1	A	29	SER
1	A	111	SER
1	A	168	PHE
1	A	178	GLU
1	A	211	HIS
1	B	52	THR
1	B	54	LEU
1	B	89	GLU
1	B	130	ASP
1	B	178	GLU
1	B	211	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	137	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	185	HIS
1	B	47	HIS
1	B	104	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	194/241 (80%)	-0.00	9 (4%) 37 41	13, 21, 43, 58	0
1	B	195/241 (80%)	0.14	10 (5%) 33 37	14, 23, 43, 69	0
All	All	389/482 (80%)	0.07	19 (4%) 35 38	13, 22, 43, 69	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	25	ALA	5.7
1	A	22	ALA	5.2
1	A	168	PHE	4.1
1	B	151	GLY	3.8
1	B	25	ALA	3.5
1	A	21	LEU	3.5
1	A	23	GLY	3.2
1	A	151	GLY	3.2
1	B	22	ALA	3.1
1	B	45	TYR	2.9
1	A	193	LEU	2.8
1	B	152	GLY	2.8
1	B	23	GLY	2.6
1	B	50	ASN	2.6
1	A	152	GLY	2.5
1	B	27	THR	2.5
1	A	6	ARG	2.1
1	B	52	THR	2.1
1	B	167	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.