



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 04:24 PM UTC

PDB ID : 3HXF / pdb_00003hxf
Title : Engineered RabGGTase in complex with a peptidomimetic inhibitor (compound 32)
Authors : Guo, Z.; Alexandrov, K.; Waldmann, H.; Goody, R.S.; Blankenfeldt, W.
Deposited on : 2009-06-20
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	6% 85% 7% 7%
2	B	331	% 85% 11% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5264 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Geranylgeranyl transferase type-2 subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	309	2416	1545	410	447	14	0	5	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP Q08602
A	238	ALA	-	linker	UNP Q08602
A	239	GLY	-	linker	UNP Q08602
A	240	SER	-	linker	UNP Q08602
A	241	GLY	-	linker	UNP Q08602

- Molecule 2 is a protein called Geranylgeranyl transferase type-2 subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	322	2492	1598	416	458	20	0	2	0


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is N-undecanoyl-L-histidyl-L-histidyl-N-methyl-N-(2-pyridin-2-ylethyl)-L-tyrosi
namide (CCD ID: BD9) (formula: C₄₀H₅₅N₉O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			54	40	9	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	124	Total	O	0	0
			124	124		
6	B	176	Total	O	0	0
			176	176		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Geranylgeranyl transferase type-2 subunit alpha



- Molecule 2: Geranylgeranyl transferase type-2 subunit beta



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.72Å 90.86Å 114.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.40 – 1.90 29.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.40-1.90) 99.8 (29.40-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.95 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.153 , 0.212 0.163 , 0.218	Depositor DCC
R_{free} test set	2788 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.6	Xtriage
Anisotropy	0.323	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5264	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA, BD9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	2/2489 (0.1%)	1.09	6/3393 (0.2%)
2	B	1.33	7/2556 (0.3%)	1.09	4/3469 (0.1%)
All	All	1.27	9/5045 (0.2%)	1.09	10/6862 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	315	ARG	CD-NE	-8.66	1.34	1.46
1	A	282	ARG	CD-NE	-6.88	1.36	1.46
2	B	243	TRP	N-CA	-6.80	1.38	1.47
2	B	183	CYS	CA-CB	6.41	1.63	1.53
2	B	246	LEU	N-CA	-6.25	1.38	1.46
2	B	175	MET	SD-CE	-5.89	1.64	1.79
2	B	243	TRP	C-O	-5.80	1.19	1.23
2	B	168	ILE	CA-CB	-5.34	1.48	1.54
2	B	51	TYR	CA-C	5.05	1.59	1.52

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	315	ARG	NE-CZ-NH2	-9.55	110.60	119.20
1	A	315	ARG	NE-CZ-NH1	9.10	130.60	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	282	ARG	NE-CZ-NH2	-7.57	112.39	119.20
2	B	182	GLY	CA-C-N	7.51	130.21	120.44
2	B	182	GLY	C-N-CA	7.51	130.21	120.44
1	A	282	ARG	CG-CD-NE	-6.58	97.53	112.00
2	B	232	ARG	N-CA-C	5.96	113.50	108.13
2	B	105	LEU	N-CA-C	5.61	117.39	111.28
1	A	45	ASP	N-CA-C	5.24	116.62	109.18
1	A	240	SER	N-CA-C	5.13	117.66	111.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	239	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2416	0	2292	12	0
2	B	2492	0	2437	23	0
3	B	1	0	0	0	0
4	B	1	0	0	0	0
5	B	54	0	52	1	0
6	A	124	0	0	0	2
6	B	176	0	0	2	1
All	All	5264	0	4781	35	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:94:HIS:HD2	2:B:96:LEU:H	1.23	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:23:HIS:HD1	2:B:277:GLY:H	1.28	0.81
2:B:190:HIS:HD2	2:B:192:GLY:H	1.30	0.76
2:B:323:VAL:HG23	2:B:325:VAL:HG12	1.68	0.75
2:B:190:HIS:CD2	2:B:192:GLY:H	2.08	0.71
1:A:45:ASP:OD2	1:A:47:SER:OG	2.05	0.70
2:B:115:ASN:HD22	2:B:115:ASN:C	2.02	0.67
1:A:119:ARG:HG3	1:A:119:ARG:HH21	1.62	0.64
1:A:70:ARG:HH11	1:A:112:HIS:HD2	1.45	0.63
2:B:19:LEU:H	2:B:271:GLN:HE22	1.46	0.62
2:B:190:HIS:HD2	2:B:192:GLY:N	2.03	0.55
2:B:323:VAL:HG23	2:B:325:VAL:CG1	2.36	0.55
1:A:119:ARG:HG3	1:A:119:ARG:NH2	2.19	0.55
1:A:92:GLU:OE2	1:A:112:HIS:HE1	1.92	0.53
2:B:37:TYR:O	2:B:41:MET:HG2	2.07	0.53
2:B:94:HIS:HE1	6:B:347:HOH:O	1.93	0.52
1:A:171:ILE:HA	1:A:177:ASN:ND2	2.27	0.50
1:A:220:PHE:CZ	1:A:259:SER:HB3	2.48	0.48
2:B:23:HIS:HD1	2:B:277:GLY:N	2.03	0.48
2:B:45:LEU:HD12	2:B:45:LEU:C	2.39	0.47
2:B:49:GLY:HA2	2:B:52:TRP:CE3	2.50	0.47
1:A:242:ARG:HA	1:A:245:LEU:HG	1.97	0.46
2:B:246:LEU:HG	2:B:255:LEU:HD11	1.97	0.46
2:B:54:LEU:HD11	2:B:71:ILE:HD13	1.98	0.46
2:B:19:LEU:H	2:B:271:GLN:NE2	2.14	0.45
2:B:37:TYR:C	2:B:39:TYR:H	2.27	0.43
5:B:334:BD9:H5	5:B:334:BD9:H2A	1.80	0.42
1:A:202:ARG:HA	1:A:238:ALA:O	2.19	0.42
2:B:52:TRP:HA	2:B:292:LEU:HD22	2.01	0.42
2:B:246:LEU:HG	2:B:255:LEU:CD1	2.51	0.41
1:A:111:HIS:HD2	6:B:372:HOH:O	2.04	0.41
2:B:243:TRP:CZ2	2:B:314:CYS:HB2	2.56	0.40
1:A:119:ARG:NH2	1:A:119:ARG:CG	2.85	0.40
1:A:226:GLN:HE21	2:B:233:PRO:HB3	1.86	0.40
2:B:310:SER:HB2	2:B:320:LEU:HD12	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:377:HOH:O	6:B:369:HOH:O[4_455]	1.00	1.20
6:A:363:HOH:O	6:A:417:HOH:O[2_554]	1.90	0.30

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/331 (94%)	301 (97%)	9 (3%)	0	100	100
2	B	320/331 (97%)	317 (99%)	3 (1%)	0	100	100
All	All	630/662 (95%)	618 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/293 (85%)	240 (96%)	9 (4%)	31	23
2	B	269/284 (95%)	261 (97%)	8 (3%)	36	30
All	All	518/577 (90%)	501 (97%)	17 (3%)	35	26

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	LEU
1	A	120	LEU
1	A	162	GLU
1	A	236	LEU
1	A	243	CYS
1	A	247	VAL
1	A	271	LYS
1	A	282	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	315	ARG
2	B	20	LEU
2	B	22	LYS
2	B	115	ASN
2	B	159	LEU
2	B	236	LEU
2	B	260[A]	ARG
2	B	260[B]	ARG
2	B	325	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	GLN
1	A	59	ASN
1	A	112	HIS
1	A	177	ASN
1	A	192	HIS
1	A	226	GLN
1	A	264	GLN
2	B	94	HIS
2	B	115	ASN
2	B	190	HIS
2	B	209	GLN
2	B	271	GLN
2	B	305	GLN
2	B	324	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BD9	B	334	3	57,57,57	1.09	4 (7%)	72,73,73	1.69	17 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BD9	B	334	3	-	12/55/55/55	0/4/4/4

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	334	BD9	C27-C28	4.06	1.55	1.49
5	B	334	BD9	C29-N6	-2.55	1.33	1.37
5	B	334	BD9	C14-N2	2.07	1.50	1.45
5	B	334	BD9	C20-C21	2.06	1.57	1.52

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	334	BD9	C27-C12-N1	4.37	118.97	110.64
5	B	334	BD9	C27-C12-C13	4.18	120.43	110.57
5	B	334	BD9	C12-C13-N2	3.86	124.87	116.63
5	B	334	BD9	C6-C7-C8	-3.80	102.65	113.19
5	B	334	BD9	C23-N5-C24	3.51	122.34	117.41
5	B	334	BD9	O2-C13-C12	-3.08	114.02	120.48
5	B	334	BD9	C13-C12-N1	-2.83	103.44	111.11
5	B	334	BD9	C22-C23-N5	-2.49	119.47	123.42
5	B	334	BD9	O4-C18-C16	-2.48	115.08	119.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	334	BD9	O1-C8-C7	-2.45	117.58	122.02
5	B	334	BD9	N7-C30-N6	-2.39	106.57	111.00
5	B	334	BD9	O3-C15-N3	2.31	127.10	122.96
5	B	334	BD9	C17-C16-C18	-2.29	105.13	109.94
5	B	334	BD9	C38-C17-C16	2.23	119.29	113.36
5	B	334	BD9	C34-N9-C32	2.22	110.84	105.80
5	B	334	BD9	C22-C26-C25	-2.05	117.71	120.24
5	B	334	BD9	N8-C34-N9	-2.01	106.56	111.49

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	334	BD9	C27-C12-C13-N2
5	B	334	BD9	C27-C12-C13-O2
5	B	334	BD9	C2-C3-C4-C5
5	B	334	BD9	C5-C6-C7-C8
5	B	334	BD9	C1-C2-C3-C4
5	B	334	BD9	C11-C10-C9-C1
5	B	334	BD9	C14-C31-C32-C33
5	B	334	BD9	C14-C31-C32-N9
5	B	334	BD9	C2-C1-C9-C10
5	B	334	BD9	C18-C16-C17-C38
5	B	334	BD9	C4-C5-C6-C7
5	B	334	BD9	C9-C1-C2-C3


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	334	BD9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	309/331 (93%)	0.22	20 (6%)	25 26	16, 34, 66, 99	5 (1%)
2	B	322/331 (97%)	-0.28	4 (1%)	76 79	14, 23, 45, 87	3 (0%)
All	All	631/662 (95%)	-0.04	24 (3%)	44 47	14, 27, 61, 99	8 (1%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	ALA	4.4
1	A	176	SER	4.3
2	B	37	TYR	4.3
1	A	240	SER	4.0
2	B	39	TYR	3.8
1	A	175	PHE	3.7
1	A	242	ARG	3.5
1	A	330	ALA	3.4
1	A	25	LEU	3.3
2	B	110	SER	3.1
1	A	16	ALA	2.9
1	A	195	PRO	2.8
1	A	220	PHE	2.8
1	A	241	GLY	2.6
1	A	28	TYR	2.4
1	A	193	PRO	2.3
1	A	243	CYS	2.3
1	A	27	LEU	2.3
1	A	221	THR	2.2
1	A	239	GLY	2.1
1	A	21	ARG	2.1
2	B	111	ILE	2.0
1	A	203	LEU	2.0
1	A	194	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	BD9	B	334	54/54	0.86	0.15	32,44,71,73	0
4	CA	B	333	1/1	0.99	0.09	29,29,29,29	0
3	ZN	B	332	1/1	1.00	0.04	25,25,25,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around BD9 B 334:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.