



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 07:09 AM UTC

PDB ID : 5HX4 / pdb_00005hx4
Title : Zinc-Free APOBEC3F Catalytic Domain Crystal Structure
Authors : Shaban, N.M.; Shi, K.; Aihara, H.; Harris, R.S.
Deposited on : 2016-01-29
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	199	4% 82% 7% 11%
1	B	199	7% 76% 8% 16%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5702 atoms, of which 2667 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA dC->dU-editing enzyme APOBEC-3F.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	177	Total	C	H	N	O	S	0	0	0
			2852	956	1365	246	276	9			
1	B	168	Total	C	H	N	O	S	0	3	0
			2733	924	1302	229	269	9			

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	175	GLY	-	expression tag	UNP Q8IUX4
A	176	PRO	-	expression tag	UNP Q8IUX4
A	177	LEU	-	expression tag	UNP Q8IUX4
A	178	GLY	-	expression tag	UNP Q8IUX4
A	179	SER	-	expression tag	UNP Q8IUX4
A	180	PRO	-	expression tag	UNP Q8IUX4
A	181	GLY	-	expression tag	UNP Q8IUX4
A	182	ILE	-	expression tag	UNP Q8IUX4
A	183	PRO	-	expression tag	UNP Q8IUX4
A	184	GLY	-	expression tag	UNP Q8IUX4
A	196	ASP	TYR	engineered mutation	UNP Q8IUX4
A	247	GLY	HIS	engineered mutation	UNP Q8IUX4
A	248	ARG	CYS	engineered mutation	UNP Q8IUX4
A	259	ALA	CYS	engineered mutation	UNP Q8IUX4
A	302	LYS	PHE	engineered mutation	UNP Q8IUX4
A	310	ASP	TRP	engineered mutation	UNP Q8IUX4
A	355	ASP	LYS	engineered mutation	UNP Q8IUX4
A	358	ASP	LYS	engineered mutation	UNP Q8IUX4
A	363	ASP	PHE	engineered mutation	UNP Q8IUX4
B	175	GLY	-	expression tag	UNP Q8IUX4
B	176	PRO	-	expression tag	UNP Q8IUX4
B	177	LEU	-	expression tag	UNP Q8IUX4
B	178	GLY	-	expression tag	UNP Q8IUX4
B	179	SER	-	expression tag	UNP Q8IUX4
B	180	PRO	-	expression tag	UNP Q8IUX4

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	181	GLY	-	expression tag	UNP Q8IUX4
B	182	ILE	-	expression tag	UNP Q8IUX4
B	183	PRO	-	expression tag	UNP Q8IUX4
B	184	GLY	-	expression tag	UNP Q8IUX4
B	196	ASP	TYR	engineered mutation	UNP Q8IUX4
B	247	GLY	HIS	engineered mutation	UNP Q8IUX4
B	248	ARG	CYS	engineered mutation	UNP Q8IUX4
B	259	ALA	CYS	engineered mutation	UNP Q8IUX4
B	302	LYS	PHE	engineered mutation	UNP Q8IUX4
B	310	ASP	TRP	engineered mutation	UNP Q8IUX4
B	355	ASP	LYS	engineered mutation	UNP Q8IUX4
B	358	ASP	LYS	engineered mutation	UNP Q8IUX4
B	363	ASP	PHE	engineered mutation	UNP Q8IUX4



- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	64	Total O 64 64	0	0
2	B	53	Total O 53 53	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3F



- Molecule 1: DNA dC->dU-editing enzyme APOBEC-3F



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	36.10Å 51.27Å 99.77Å 90.00° 90.85° 90.00°	Depositor
Resolution (Å)	36.10 – 1.92 36.10 – 1.92	Depositor EDS
% Data completeness (in resolution range)	92.2 (36.10-1.92) 87.7 (36.10-1.92)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 1.92Å)	Xtriage
Refinement program	PHENIX (dev_2356: ???)	Depositor
R, R_{free}	0.185 , 0.214 0.187 , 0.217	Depositor DCC
R_{free} test set	1299 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	33.1	Xtriage
Anisotropy	0.249	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.048 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5702	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/1536	0.57	0/2084
1	B	0.48	0/1487	0.58	0/2020
All	All	0.51	0/3023	0.57	0/4104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1487	1365	1365	11	0
1	B	1431	1302	1289	15	0
2	A	64	0	0	5	0
2	B	53	0	0	3	0
All	All	3035	2667	2654	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:215:GLU:OE2	1:B:239:ARG:NH2	2.09	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:LYS:O	2:A:401:HOH:O	1.95	0.84
1:B:359:TYR:OH	2:B:401:HOH:O	2.02	0.76
1:A:209:LYS:NZ	2:A:403:HOH:O	2.25	0.70
1:A:190:ASN:ND2	2:A:404:HOH:O	2.29	0.66
1:B:278:SER:HB3	1:B:314[A]:TYR:CZ	2.33	0.64
1:B:278:SER:HB3	1:B:314[A]:TYR:OH	1.98	0.63
1:B:334:LYS:NZ	2:B:403:HOH:O	2.36	0.59
1:B:309:PHE:O	1:B:315:GLN:NE2	2.33	0.54
1:B:313:ASP:OD1	1:B:314[A]:TYR:N	2.43	0.52
1:A:361:PHE:CD1	1:A:361:PHE:C	2.90	0.49
1:B:278:SER:CB	1:B:314[A]:TYR:OH	2.60	0.49
1:A:278:SER:HB3	1:A:306:LEU:O	2.11	0.49
1:B:306:LEU:HA	1:B:314[A]:TYR:OH	2.12	0.49
1:B:278:SER:HB3	1:B:314[A]:TYR:CE2	2.48	0.49
1:A:233:TRP:CE3	1:B:235:ARG:HD2	2.49	0.48
1:B:313:ASP:OD1	1:B:314[B]:TYR:N	2.48	0.46
1:A:193:GLU:OE2	2:A:402:HOH:O	2.20	0.46
1:B:277:TRP:CE3	1:B:307:TYR:CB	3.00	0.44
1:A:369:GLN:O	1:A:373:GLU:HG2	2.18	0.43
1:A:190:ASN:N	1:A:191:PRO:HD2	2.34	0.43
1:A:306:LEU:HB2	2:A:406:HOH:O	2.19	0.42
1:A:309:PHE:HB3	1:A:311:ASP:OD1	2.18	0.42
1:B:357:LEU:C	1:B:357:LEU:HD23	2.45	0.41
1:B:266:ASN:HB2	2:B:402:HOH:O	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/199 (87%)	168 (97%)	5 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	165/199 (83%)	160 (97%)	5 (3%)	0	100	100
All	All	338/398 (85%)	328 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	160/178 (90%)	159 (99%)	1 (1%)	78	75
1	B	157/178 (88%)	156 (99%)	1 (1%)	78	75
All	All	317/356 (89%)	315 (99%)	2 (1%)	78	75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	289	GLU
1	B	347	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	294	HIS
1	B	214	ASN
1	B	268	ASN
1	B	298	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	177/199 (88%)	0.23	8 (4%) 38 43	29, 45, 93, 110	0
1	B	168/199 (84%)	0.39	14 (8%) 17 21	16, 46, 99, 121	2 (1%)
All	All	345/398 (86%)	0.31	22 (6%) 25 29	16, 46, 97, 121	2 (0%)

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	277	TRP	4.3
1	A	207	LEU	3.9
1	A	227	HIS	3.2
1	B	207	LEU	3.0
1	B	206	ASN	3.0
1	B	202	PHE	2.9
1	B	308	TYR	2.9
1	B	307	TYR	2.7
1	A	202	PHE	2.7
1	B	201	TYR	2.6
1	B	309	PHE	2.6
1	B	281	PRO	2.6
1	A	314	TYR	2.6
1	B	279	PRO	2.6
1	A	355	ASP	2.5
1	A	228	HIS	2.4
1	A	208	ARG	2.3
1	B	334	LYS	2.3
1	B	359	TYR	2.3
1	B	266	ASN	2.2
1	A	373	GLU	2.1
1	B	278	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.