



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 06:05 PM EDT

PDB ID : 3HYS / pdb_00003hys
Title : Structure of Rv0554 from Mycobacterium tuberculosis complexed with Malonic Acid
Authors : Johnston, J.M.; Baker, E.N.
Deposited on : 2009-06-23
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	293	
1	B	293	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MLA	B	5188	-	X	-	-
3	TRS	A	263	-	X	X	-
4	EDO	A	266	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4556 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called protein Rv0554, putative Bromoperoxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	1	0
			2119	1348	370	390	11			
1	B	261	Total	C	N	O	S	0	2	0
			1997	1269	348	370	10			

There are 64 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-30	MET	-	expression tag	UNP O06420
A	-29	SER	-	expression tag	UNP O06420
A	-28	TYR	-	expression tag	UNP O06420
A	-27	TYR	-	expression tag	UNP O06420
A	-26	HIS	-	expression tag	UNP O06420
A	-25	HIS	-	expression tag	UNP O06420
A	-24	HIS	-	expression tag	UNP O06420
A	-23	HIS	-	expression tag	UNP O06420
A	-22	HIS	-	expression tag	UNP O06420
A	-21	HIS	-	expression tag	UNP O06420
A	-20	ASP	-	expression tag	UNP O06420
A	-19	TYR	-	expression tag	UNP O06420
A	-18	ASP	-	expression tag	UNP O06420
A	-17	ILE	-	expression tag	UNP O06420
A	-16	PRO	-	expression tag	UNP O06420
A	-15	THR	-	expression tag	UNP O06420
A	-14	THR	-	expression tag	UNP O06420
A	-13	GLU	-	expression tag	UNP O06420
A	-12	ASN	-	expression tag	UNP O06420
A	-11	LEU	-	expression tag	UNP O06420
A	-10	TYR	-	expression tag	UNP O06420
A	-9	PHE	-	expression tag	UNP O06420
A	-8	GLN	-	expression tag	UNP O06420
A	-7	GLY	-	expression tag	UNP O06420
A	-6	ALA	-	expression tag	UNP O06420

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	MET	-	expression tag	UNP O06420
A	-4	ASP	-	expression tag	UNP O06420
A	-3	PRO	-	expression tag	UNP O06420
A	-2	GLU	-	expression tag	UNP O06420
A	-1	PHE	-	expression tag	UNP O06420
A	0	ARG	-	expression tag	UNP O06420
A	1	VAL	-	expression tag	UNP O06420
B	-30	MET	-	expression tag	UNP O06420
B	-29	SER	-	expression tag	UNP O06420
B	-28	TYR	-	expression tag	UNP O06420
B	-27	TYR	-	expression tag	UNP O06420
B	-26	HIS	-	expression tag	UNP O06420
B	-25	HIS	-	expression tag	UNP O06420
B	-24	HIS	-	expression tag	UNP O06420
B	-23	HIS	-	expression tag	UNP O06420
B	-22	HIS	-	expression tag	UNP O06420
B	-21	HIS	-	expression tag	UNP O06420
B	-20	ASP	-	expression tag	UNP O06420
B	-19	TYR	-	expression tag	UNP O06420
B	-18	ASP	-	expression tag	UNP O06420
B	-17	ILE	-	expression tag	UNP O06420
B	-16	PRO	-	expression tag	UNP O06420
B	-15	THR	-	expression tag	UNP O06420
B	-14	THR	-	expression tag	UNP O06420
B	-13	GLU	-	expression tag	UNP O06420
B	-12	ASN	-	expression tag	UNP O06420
B	-11	LEU	-	expression tag	UNP O06420
B	-10	TYR	-	expression tag	UNP O06420
B	-9	PHE	-	expression tag	UNP O06420
B	-8	GLN	-	expression tag	UNP O06420
B	-7	GLY	-	expression tag	UNP O06420
B	-6	ALA	-	expression tag	UNP O06420
B	-5	MET	-	expression tag	UNP O06420
B	-4	ASP	-	expression tag	UNP O06420
B	-3	PRO	-	expression tag	UNP O06420
B	-2	GLU	-	expression tag	UNP O06420
B	-1	PHE	-	expression tag	UNP O06420
B	0	ARG	-	expression tag	UNP O06420
B	1	VAL	-	expression tag	UNP O06420

- Molecule 2 is MALONIC ACID (CCD ID: MLA) (formula: C₃H₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	3	4		
2	B	1	Total	C	O	0	0
			7	3	4		

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Na 1 1	0	0
5	B	1	Total Na 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	222	Total 222	O 222	0	0
6	B	166	Total 166	O 166	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: protein Rv0554, putative Bromoperoxidase



- Molecule 1: protein Rv0554, putative Bromoperoxidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	100.44Å 100.44Å 135.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.30 50.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-2.30) 99.8 (50.00-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.67 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.166 , 0.223 0.167 , 0.224	Depositor DCC
R_{free} test set	1579 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	28.0	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4556	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TRS, EDO, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.47	8/2172 (0.4%)	1.19	5/2958 (0.2%)
1	B	1.37	3/2049 (0.1%)	1.22	6/2793 (0.2%)
All	All	1.42	11/4221 (0.3%)	1.20	11/5751 (0.2%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	78	ILE	CA-CB	7.75	1.62	1.54
1	B	200	ALA	CA-CB	6.91	1.64	1.53
1	A	145	ALA	CA-CB	6.17	1.63	1.53
1	B	70	ALA	CA-CB	5.69	1.62	1.53
1	A	169	PHE	C-O	-5.49	1.17	1.24
1	A	185	ASP	N-CA	5.42	1.53	1.46
1	A	90	ALA	CA-CB	5.25	1.61	1.53
1	A	214	THR	CA-C	5.15	1.58	1.52
1	A	67	ALA	CA-CB	5.12	1.61	1.53
1	A	19	ALA	C-O	5.10	1.30	1.23
1	A	86	VAL	CA-CB	5.00	1.60	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	215	PRO	CA-C-N	8.34	128.83	119.32
1	B	215	PRO	C-N-CA	8.34	128.83	119.32
1	A	260	VAL	N-CA-C	7.33	119.55	109.80
1	B	215	PRO	O-C-N	5.98	124.06	121.31
1	B	174	ILE	CB-CA-C	-5.67	103.62	110.99
1	A	153	ARG	CB-CA-C	-5.66	101.99	110.88
1	B	136	GLN	CB-CA-C	-5.41	101.43	110.19
1	B	169	PHE	N-CA-C	5.25	117.74	111.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	33	VAL	CA-C-N	-5.24	113.23	119.05
1	A	33	VAL	C-N-CA	-5.24	113.23	119.05
1	A	259	SER	N-CA-C	5.05	120.06	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2119	0	2087	29	0
1	B	1997	0	1971	17	0
2	A	7	0	2	0	0
2	B	7	0	2	1	0
3	A	8	0	12	7	0
4	A	12	0	18	6	0
4	B	16	0	24	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	222	0	0	2	0
6	B	166	0	0	2	0
All	All	4556	0	4116	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:VAL:O	1:A:261:LYS:HB2	1.71	0.88
1:A:21:ARG:HB2	3:A:263:TRS:H22	1.57	0.86
1:A:21:ARG:CB	3:A:263:TRS:H22	2.05	0.85
1:B:147:LEU:HD23	1:B:156:LEU:CD2	2.06	0.85
1:A:65:MET:HB3	4:A:266:EDO:H21	1.65	0.79
1:A:65:MET:HE2	4:A:266:EDO:H22	1.71	0.70

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:153:ARG:HD3	1:B:153:ARG:HH22	1.56	0.69
1:A:260:VAL:O	1:A:261:LYS:CB	2.42	0.68
1:B:21:ARG:HB3	2:B:5188:MLA:HC22	1.78	0.66
1:B:147:LEU:CD2	1:B:156:LEU:CD2	2.75	0.64
1:A:21:ARG:HB3	3:A:263:TRS:C2	2.28	0.63
1:A:21:ARG:CB	3:A:263:TRS:C2	2.78	0.61
1:B:147:LEU:HD23	1:B:156:LEU:HD23	1.82	0.60
1:A:48:ASN:HA	4:A:266:EDO:H11	1.84	0.60
1:B:147:LEU:CD2	1:B:156:LEU:HD23	2.32	0.59
1:A:153:ARG:CD	1:B:153:ARG:HH22	2.17	0.57
1:B:136:GLN:HB3	6:B:373:HOH:O	2.04	0.56
1:A:120:ARG:CZ	1:A:215:PRO:HG3	2.38	0.53
1:B:199:ILE:O	1:B:227:PRO:HD2	2.07	0.53
1:A:65:MET:HE2	4:A:266:EDO:C2	2.36	0.53
1:A:21:ARG:HB3	3:A:263:TRS:H21	1.90	0.52
1:A:88:MET:HB2	3:A:263:TRS:H32	1.91	0.52
1:A:97:MET:CE	1:A:201:ALA:HB3	2.41	0.51
3:A:263:TRS:H31	6:A:476:HOH:O	2.11	0.50
1:A:138:PRO:HB2	1:B:129:GLU:OE1	2.11	0.50
1:B:160:VAL:HG23	6:B:400:HOH:O	2.13	0.48
1:A:97:MET:HE2	1:A:201:ALA:HB3	1.96	0.48
1:A:65:MET:CB	4:A:266:EDO:H21	2.39	0.47
1:B:82:ARG:HD3	1:B:257:PHE:CD1	2.51	0.45
1:A:153:ARG:NE	1:B:153:ARG:HH22	2.16	0.44
1:A:126:ALA:HB1	1:B:141:TYR:HA	2.00	0.43
1:A:208:PHE:HB2	1:A:211:ASP:HB2	2.01	0.42
1:A:123:PHE:CD1	1:B:144:ARG:HG3	2.54	0.42
1:A:122:PHE:CZ	1:B:140:THR:HG23	2.55	0.41
1:A:261:LYS:HA	1:A:261:LYS:HD2	1.83	0.41
1:A:175:LYS:HD2	1:A:177:THR:CG2	2.51	0.41
1:A:46:PHE:HE2	4:A:266:EDO:C1	2.34	0.41
1:A:220[A]:ARG:HG2	6:A:289:HOH:O	2.19	0.41
1:B:29:HIS:N	1:B:30:PRO:CD	2.83	0.41
1:A:120:ARG:NH2	1:A:215:PRO:HG3	2.36	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/293 (94%)	270 (98%)	4 (2%)	0	100	100
1	B	261/293 (89%)	258 (99%)	3 (1%)	0	100	100
All	All	535/586 (91%)	528 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/233 (93%)	212 (98%)	5 (2%)	44	63
1	B	205/233 (88%)	194 (95%)	11 (5%)	20	29
All	All	422/466 (91%)	406 (96%)	16 (4%)	29	44

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	120	ARG
1	A	148	LEU
1	A	166	ILE
1	A	212	VAL
1	A	261	LYS
1	B	2	ILE
1	B	4	LEU
1	B	63	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	99	VAL
1	B	135	VAL
1	B	136	GLN
1	B	147	LEU
1	B	155	THR
1	B	189	GLN
1	B	243	PHE
1	B	260	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-12	ASN
1	A	32	GLN
1	A	150	ASN
1	B	32	GLN
1	B	136	GLN
1	B	189	GLN
1	B	191	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	B	265	-	3,3,3	0.56	0	2,2,2	0.48	0
4	EDO	B	263	-	3,3,3	0.61	0	2,2,2	0.41	0
4	EDO	B	266	-	3,3,3	0.71	0	2,2,2	0.22	0
3	TRS	A	263	-	7,7,7	0.87	0	9,9,9	2.54	4 (44%)
4	EDO	A	265	-	3,3,3	0.41	0	2,2,2	0.84	0
4	EDO	A	266	-	3,3,3	0.71	0	2,2,2	0.90	0
4	EDO	B	264	-	3,3,3	1.04	0	2,2,2	0.91	0
2	MLA	A	5188	-	6,6,6	4.19	3 (50%)	7,7,7	1.40	2 (28%)
2	MLA	B	5188	-	6,6,6	4.56	4 (66%)	7,7,7	2.81	4 (57%)
4	EDO	A	264	-	3,3,3	0.51	0	2,2,2	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	265	-	-	1/1/1/1	-
4	EDO	B	263	-	-	1/1/1/1	-
4	EDO	B	266	-	-	0/1/1/1	-
3	TRS	A	263	-	-	9/9/9/9	-
4	EDO	A	265	-	-	0/1/1/1	-
4	EDO	A	266	-	-	1/1/1/1	-
4	EDO	B	264	-	-	0/1/1/1	-
2	MLA	A	5188	-	-	0/4/4/4	-
2	MLA	B	5188	-	-	2/4/4/4	-
4	EDO	A	264	-	-	1/1/1/1	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	5188	MLA	C2-C3	-10.33	1.35	1.51
2	A	5188	MLA	C2-C3	-9.48	1.36	1.51
2	B	5188	MLA	O1A-C1	-2.39	1.22	1.30
2	B	5188	MLA	O1B-C1	2.20	1.29	1.22
2	A	5188	MLA	O3B-C3	-2.14	1.23	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	5188	MLA	C2-C1	-2.12	1.47	1.51
2	B	5188	MLA	C2-C1	-2.03	1.47	1.51

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	5188	MLA	O1B-C1-C2	-4.18	110.21	122.11
3	A	263	TRS	O3-C3-C	4.00	122.02	110.88
3	A	263	TRS	C3-C-C1	3.95	121.17	110.66
2	B	5188	MLA	O3B-C3-C2	3.79	126.26	114.51
3	A	263	TRS	O2-C2-C	3.40	120.35	110.88
2	B	5188	MLA	O1A-C1-C2	3.24	124.54	114.51
3	A	263	TRS	C2-C-C1	-3.00	102.66	110.66
2	B	5188	MLA	O3A-C3-C2	-2.55	114.84	122.11
2	A	5188	MLA	C3-C2-C1	2.43	121.51	112.95
2	A	5188	MLA	O3B-C3-O3A	-2.37	117.23	123.33

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	263	TRS	N-C-C1-O1
3	A	263	TRS	C1-C-C2-O2
3	A	263	TRS	C3-C-C2-O2
3	A	263	TRS	N-C-C2-O2
3	A	263	TRS	C1-C-C3-O3
3	A	263	TRS	C2-C-C3-O3
3	A	263	TRS	N-C-C3-O3
3	A	263	TRS	C2-C-C1-O1
4	A	266	EDO	O1-C1-C2-O2
4	B	265	EDO	O1-C1-C2-O2
2	B	5188	MLA	O1A-C1-C2-C3
2	B	5188	MLA	O1B-C1-C2-C3
4	A	264	EDO	O1-C1-C2-O2
4	B	263	EDO	O1-C1-C2-O2
3	A	263	TRS	C3-C-C1-O1

There are no ring outliers.

3 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	263	TRS	7	0
4	A	266	EDO	6	0
2	B	5188	MLA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/293 (93%)	-0.66	3 (1%) 78 79	12, 21, 36, 62	2 (0%)
1	B	261/293 (89%)	-0.33	2 (0%) 82 83	14, 29, 42, 65	3 (1%)
All	All	536/586 (91%)	-0.50	5 (0%) 81 82	12, 24, 40, 65	5 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2	ILE	3.6
1	A	225	ALA	2.9
1	B	225	ALA	2.9
1	A	260	VAL	2.6
1	A	-13	GLU	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	B	266	4/4	0.80	0.16	49,51,53,53	0
4	EDO	B	264	4/4	0.85	0.20	38,38,40,41	0
4	EDO	A	264	4/4	0.85	0.17	53,56,57,58	0
3	TRS	A	263	8/8	0.87	0.13	40,46,47,51	0
5	NA	A	267	1/1	0.87	0.08	39,39,39,39	0
4	EDO	B	265	4/4	0.88	0.14	43,45,46,51	0
2	MLA	A	5188	7/7	0.89	0.12	49,54,57,59	0
4	EDO	A	265	4/4	0.90	0.11	40,43,44,44	0
4	EDO	B	263	4/4	0.90	0.13	52,53,54,54	0
5	NA	B	267	1/1	0.91	0.11	40,40,40,40	0
2	MLA	B	5188	7/7	0.93	0.11	18,25,34,40	0
4	EDO	A	266	4/4	0.96	0.22	21,23,25,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.