



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:31 AM UTC

PDB ID : 2HZI / pdb_00002hzi
Title : Abl kinase domain in complex with PD180970
Authors : Cowan-Jacob, S.W.; Fendrich, G.; Liebetanz, J.; Fabbro, D.; Manley, P.
Deposited on : 2006-08-09
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	5% 87% 9% • •
1	B	277	16% 82% 12% • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4925 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	8	0
			2210	1425	353	413	19			
1	B	264	Total	C	N	O	S	0	2	0
			2117	1374	341	385	17			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	GLY	-	cloning artifact	UNP P00519
A	225	ALA	-	cloning artifact	UNP P00519
A	226	MET	-	cloning artifact	UNP P00519
A	227	ASP	-	cloning artifact	UNP P00519
A	228	PRO	-	cloning artifact	UNP P00519
B	224	GLY	-	cloning artifact	UNP P00519
B	225	ALA	-	cloning artifact	UNP P00519
B	226	MET	-	cloning artifact	UNP P00519
B	227	ASP	-	cloning artifact	UNP P00519
B	228	PRO	-	cloning artifact	UNP P00519

- Molecule 2 is 6-(2,6-DICHLOROPHENYL)-2-[(4-FLUORO-3-METHYLPHENYL)AMINO]-8-METHYLPYRIDO[2,3-D]PYRIMIDIN-7(8H)-ONE (CCD ID: JIN) (formula: C₂₁H₁₅Cl₂FN₄O).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Cl	F	N	O	0	0
			29	21	2	1	4	1		
2	B	1	Total	C	Cl	F	N	O	0	0
			29	21	2	1	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	317	Total	O	0	2
			317	317		
3	B	223	Total	O	0	0
			223	223		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase ABL1



- Molecule 1: Proto-oncogene tyrosine-protein kinase ABL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	106.47Å 131.53Å 56.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.65 – 1.70 81.65 – 1.70	Depositor EDS
% Data completeness (in resolution range)	(Not available) (81.65-1.70) 99.4 (81.65-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.175 , 0.204 (Not available) , 0.205	Depositor DCC
R_{free} test set	4365 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	18.7	Xtriage
Anisotropy	0.034	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4925	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: JIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.00	3/2287 (0.1%)	1.00	2/3097 (0.1%)
1	B	0.90	4/2183 (0.2%)	1.01	5/2957 (0.2%)
All	All	0.95	7/4470 (0.2%)	1.01	7/6054 (0.1%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	437	MET	SD-CE	-9.75	1.55	1.79
1	A	496	MET	SD-CE	-9.01	1.57	1.79
1	A	278	MET	SD-CE	-7.49	1.60	1.79
1	B	496	MET	SD-CE	-7.17	1.61	1.79
1	B	472	MET	SD-CE	-7.07	1.61	1.79
1	B	350	ALA	CA-CB	-5.86	1.44	1.53
1	B	343	MET	SD-CE	-5.62	1.65	1.79

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	382	PHE	N-CA-C	-8.20	98.93	110.50
1	A	461	PRO	CB-CA-C	-6.91	102.21	111.12
1	B	319	THR	N-CA-C	6.43	120.39	112.54
1	B	403	ILE	N-CA-C	5.77	116.42	110.36
1	B	242	ILE	N-CA-C	5.73	116.38	108.12
1	A	309	PRO	O-C-N	5.42	123.70	121.15
1	B	263	LYS	N-CA-C	5.29	117.96	111.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2210	0	2129	22	0
1	B	2117	0	2035	25	0
2	A	29	0	15	1	0
2	B	29	0	15	0	0
3	A	317	0	0	4	0
3	B	223	0	0	8	0
All	All	4925	0	4194	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:237:MET:HE2	1:B:314:ILE:HD13	1.37	1.04
1:B:237:MET:CE	1:B:314:ILE:HD13	1.99	0.92
1:B:356:LYS:HG3	3:B:705:HOH:O	1.81	0.78
1:B:273:LEU:C	3:B:795:HOH:O	2.26	0.78
1:A:437:MET:HE3	3:A:850:HOH:O	1.81	0.78
1:B:386:ARG:NH1	3:B:729:HOH:O	2.23	0.71
1:B:306:THR:HA	1:B:311:PHE:HD2	1.56	0.69
1:A:479[A]:ASN:OD1	3:A:770:HOH:O	2.10	0.69
1:B:237:MET:HE2	1:B:314:ILE:CD1	2.18	0.68
1:A:331:ASN:HD22	1:A:331:ASN:C	2.02	0.67
1:A:286:GLU:O	1:A:290[B]:MET:HG3	1.95	0.66
1:A:239:ARG:HG3	1:A:312:TYR:CE1	2.34	0.63
1:A:490:HIS:HE1	3:A:758:HOH:O	1.81	0.63
1:B:404:LYS:HE3	3:B:757:HOH:O	2.01	0.61
1:A:245:LYS:HE2	1:A:260:VAL:CG2	2.31	0.60
1:A:281:GLU:OE2	1:B:490:HIS:HE1	1.85	0.60
1:A:320[B]:TYR:HB2	1:A:371:VAL:O	2.02	0.59
1:B:237:MET:HE1	1:B:242:ILE:HD11	1.85	0.58
1:A:331:ASN:HD21	1:A:333:GLN:HB2	1.71	0.55
1:A:404:LYS:NZ	3:A:829:HOH:O	2.41	0.54
1:B:336:ASN:OD1	1:B:338:VAL:HG12	2.09	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:331:ASN:C	1:A:331:ASN:ND2	2.66	0.53
1:B:273:LEU:O	3:B:804:HOH:O	2.18	0.53
1:B:306:THR:HA	1:B:311:PHE:CD2	2.40	0.53
1:B:263:LYS:HG2	1:B:264:TYR:N	2.22	0.52
1:B:236:GLU:OE2	1:B:306:THR:OG1	2.27	0.52
1:B:295:HIS:CG	1:B:296:PRO:HD2	2.46	0.51
1:A:404:LYS:HE3	1:A:445:LEU:HD23	1.94	0.50
1:B:238:GLU:O	1:B:239:ARG:C	2.58	0.47
1:B:352:GLU:OE2	1:B:356:LYS:NZ	2.46	0.47
1:A:245:LYS:CE	1:A:260:VAL:CG2	2.94	0.46
1:B:237:MET:CE	1:B:242:ILE:HD11	2.46	0.46
1:B:261:TRP:CE2	1:B:264:TYR:CE1	3.06	0.44
1:B:490:HIS:HD2	3:B:613:HOH:O	1.99	0.44
1:A:331:ASN:ND2	1:A:333:GLN:H	2.15	0.44
1:B:240:THR:C	1:B:242:ILE:H	2.26	0.44
1:A:295:HIS:CG	1:A:296:PRO:HD2	2.53	0.43
1:A:401:PHE:HB3	1:A:402:PRO:HD2	2.00	0.43
1:A:290[B]:MET:SD	2:A:600:JIN:H16	2.59	0.43
1:A:319:THR:OG1	1:A:320[B]:TYR:CD1	2.71	0.43
1:B:440:TYR:N	1:B:441:PRO:CD	2.82	0.42
1:B:278:MET:C	3:B:816:HOH:O	2.62	0.42
1:A:233:ASP:OD2	1:A:291:LYS:NZ	2.49	0.42
1:B:285:LYS:HG3	3:B:780:HOH:O	2.20	0.41
1:A:290[A]:MET:HE3	1:A:290[A]:MET:HB3	2.00	0.41
1:A:404:LYS:HE3	1:A:445:LEU:CD2	2.51	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	274/277 (99%)	263 (96%)	10 (4%)	1 (0%)	30 16

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	262/277 (95%)	250 (95%)	11 (4%)	1 (0%)	30	16
All	All	536/554 (97%)	513 (96%)	21 (4%)	2 (0%)	30	16

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	234	LYS
1	B	234	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/241 (97%)	232 (100%)	1 (0%)	84	80
1	B	216/241 (90%)	214 (99%)	2 (1%)	70	62
All	All	449/482 (93%)	446 (99%)	3 (1%)	76	69

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	331	ASN
1	B	414	ASN
1	B	453	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	331	ASN
1	A	396	HIS
1	A	490	HIS
1	B	297	ASN
1	B	300	GLN
1	B	414	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	490	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	JIN	A	600	-	32,32,32	1.85	10 (31%)	44,47,47	2.90	21 (47%)
2	JIN	B	600	-	32,32,32	1.62	6 (18%)	44,47,47	2.98	17 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JIN	A	600	-	-	1/8/8/8	0/4/4/4
2	JIN	B	600	-	-	2/8/8/8	0/4/4/4

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	JIN	C06-N11	-4.75	1.33	1.38
2	B	600	JIN	C09-C12	-3.90	1.40	1.46
2	B	600	JIN	C10-C09	3.64	1.40	1.35
2	A	600	JIN	C09-C12	-3.60	1.40	1.46
2	A	600	JIN	C12-N11	-3.08	1.34	1.38
2	A	600	JIN	C10-C09	3.02	1.40	1.35
2	B	600	JIN	C01-C10	-2.90	1.37	1.43
2	B	600	JIN	C12-N11	-2.78	1.34	1.38
2	B	600	JIN	C06-N11	-2.75	1.35	1.38
2	B	600	JIN	C27-N08	-2.61	1.35	1.40
2	A	600	JIN	C20-CL5	2.42	1.79	1.73
2	A	600	JIN	C16-C17	2.34	1.42	1.38
2	A	600	JIN	C01-C06	-2.29	1.37	1.42
2	A	600	JIN	C01-C10	-2.26	1.39	1.43
2	A	600	JIN	C04-N03	-2.24	1.31	1.34
2	A	600	JIN	C13-N11	2.14	1.50	1.46

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	600	JIN	C19-C09-C12	8.64	127.69	119.36
2	A	600	JIN	C28-C29-C30	7.56	123.23	115.88
2	B	600	JIN	C09-C12-N11	6.74	125.07	116.21
2	B	600	JIN	C13-N11-C12	6.30	123.42	117.34
2	B	600	JIN	C28-C29-C30	6.02	121.74	115.88
2	A	600	JIN	C02-C01-C06	5.82	119.69	114.55
2	A	600	JIN	C19-C09-C12	5.79	124.94	119.36
2	A	600	JIN	C34-C29-C30	-5.05	115.31	121.94
2	A	600	JIN	C13-N11-C12	5.05	122.20	117.34
2	B	600	JIN	O14-C12-C09	-4.74	119.17	125.37
2	A	600	JIN	C31-C30-C29	-4.56	117.25	124.46
2	B	600	JIN	C26-C27-C28	-4.55	114.16	119.66
2	B	600	JIN	C10-C09-C12	-4.40	115.25	119.36
2	B	600	JIN	C31-C30-C29	-4.10	117.98	124.46
2	A	600	JIN	C01-C06-N11	4.09	122.71	119.53
2	A	600	JIN	F35-C30-C29	4.06	123.68	117.66
2	A	600	JIN	C09-C12-N11	4.00	121.46	116.21
2	A	600	JIN	C26-C27-C28	-3.89	114.95	119.66
2	A	600	JIN	C19-C18-CL4	3.81	124.53	119.62
2	A	600	JIN	C18-C19-C09	-3.81	118.12	121.86
2	B	600	JIN	O14-C12-N11	-3.80	115.74	121.22
2	B	600	JIN	C02-N03-C04	3.70	120.89	115.81

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	600	JIN	C31-C26-C27	3.62	124.47	120.30
2	B	600	JIN	N03-C04-N05	-3.59	122.95	126.42
2	B	600	JIN	C34-C29-C30	-3.40	117.48	121.94
2	B	600	JIN	F35-C30-C29	3.19	122.38	117.66
2	A	600	JIN	O14-C12-C09	-3.19	121.20	125.37
2	B	600	JIN	C02-C01-C06	3.00	117.20	114.55
2	A	600	JIN	C31-C26-C27	2.79	123.51	120.30
2	A	600	JIN	C13-N11-C06	-2.71	117.59	120.09
2	A	600	JIN	O14-C12-N11	-2.71	117.32	121.22
2	A	600	JIN	C26-C27-N08	2.69	129.71	120.61
2	A	600	JIN	C10-C09-C12	-2.43	117.09	119.36
2	B	600	JIN	C26-C27-N08	2.36	128.62	120.61
2	A	600	JIN	C02-N03-C04	2.36	119.05	115.81
2	A	600	JIN	C17-C18-CL4	-2.31	113.87	118.42
2	A	600	JIN	C18-C19-C20	2.23	119.82	116.70
2	B	600	JIN	C01-C06-N11	2.19	121.23	119.53

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600	JIN	C12-C09-C19-C18
2	B	600	JIN	C12-C09-C19-C20
2	B	600	JIN	C12-C09-C19-C18


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	JIN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/277 (96%)	0.18	14 (5%) 33 36	10, 19, 33, 39	8 (2%)
1	B	264/277 (95%)	0.86	43 (16%) 4 4	15, 24, 52, 57	2 (0%)
All	All	532/554 (96%)	0.52	57 (10%) 11 11	10, 21, 45, 57	10 (1%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	389[A]	THR	6.9
1	A	390[A]	GLY	6.3
1	B	242	ILE	5.4
1	B	264	TYR	5.4
1	B	397	ALA	5.1
1	B	250	GLY	4.8
1	B	278	MET	4.8
1	B	246	HIS	4.6
1	B	233	ASP	4.5
1	A	246	HIS	4.4
1	A	276	ASP	4.2
1	B	241	ASP	4.0
1	B	309	PRO	4.0
1	A	444	ASP	4.0
1	B	280	VAL	3.9
1	B	263	LYS	3.7
1	B	265	SER	3.6
1	B	307	ARG	3.6
1	A	250	GLY	3.4
1	B	266	LEU	3.4
1	B	273	LEU	3.4
1	B	240	THR	3.4
1	A	320[A]	TYR	3.3
1	B	444	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	262	LYS	3.3
1	B	272	THR	3.2
1	B	396	HIS	3.2
1	B	308	GLU	3.1
1	B	279	GLU	3.1
1	B	237	MET	3.0
1	B	390	GLY	3.0
1	B	261	TRP	2.9
1	B	245	LYS	2.9
1	B	260	VAL	2.9
1	B	305	CYS	2.9
1	B	238	GLU	2.8
1	B	234	LYS	2.8
1	B	399	ALA	2.7
1	B	239	ARG	2.6
1	B	281	GLU	2.5
1	B	310	PRO	2.5
1	B	249	GLY	2.5
1	B	398	GLY	2.5
1	A	398	GLY	2.4
1	B	306	THR	2.4
1	A	397	ALA	2.4
1	B	320	TYR	2.4
1	B	311	PHE	2.3
1	B	235	TRP	2.3
1	A	498	GLN	2.3
1	A	251	GLY	2.3
1	B	401	PHE	2.2
1	B	446	SER	2.2
1	A	275	GLU	2.2
1	B	383	GLY	2.1
1	A	239	ARG	2.1
1	A	274	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	JIN	A	600	29/29	0.97	0.06	11,15,20,23	0
2	JIN	B	600	29/29	0.97	0.07	20,24,29,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around JIN A 600:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around JIN B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.