



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:45 PM UTC

PDB ID : 3HZJ / pdb_00003hzj
Title : Crystal structure of the RabGAP domain of the RABGAP1L protein
Authors : Nedyalkova, L.; Tempel, W.; Tong, Y.; Zhong, N.; MacKenzie, F.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.; Bochkarev, A.; Park, H.; Structural Genomics Consortium (SGC)
Deposited on : 2009-06-23
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	310	
1	B	310	
1	C	310	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	UNX	A	1	-	-	-	X
2	UNX	A	2	-	-	-	X
2	UNX	A	4	-	-	-	X
2	UNX	A	901	-	-	-	X
2	UNX	A	902	-	-	-	X
2	UNX	A	903	-	-	-	X
2	UNX	A	904	-	-	-	X
2	UNX	A	905	-	-	-	X
2	UNX	A	906	-	-	-	X
2	UNX	A	907	-	-	-	X
2	UNX	A	908	-	-	-	X
2	UNX	A	909	-	-	-	X
2	UNX	A	910	-	-	-	X
2	UNX	A	911	-	-	-	X
2	UNX	B	901	-	-	-	X
2	UNX	B	902	-	-	-	X
2	UNX	B	908	-	-	-	X
2	UNX	B	910	-	-	-	X
2	UNX	C	3	-	-	-	X
2	UNX	C	5	-	-	-	X
2	UNX	C	6	-	-	-	X
2	UNX	C	902	-	-	-	X
2	UNX	C	903	-	-	-	X
2	UNX	C	904	-	-	-	X
2	UNX	C	905	-	-	-	X
2	UNX	C	906	-	-	-	X
2	UNX	C	907	-	-	-	X
2	UNX	C	908	-	-	-	X
2	UNX	C	909	-	-	-	X
2	UNX	C	911	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6473 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAB GTPase-activating protein 1-like.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	287	Total	C	N	O	S	Se	0	0	0
			2270	1466	378	410	8	8			
1	B	256	Total	C	N	O	S	Se	0	0	0
			1887	1231	312	330	7	7			
1	C	288	Total	C	N	O	S	Se	0	0	0
			2271	1467	382	406	8	8			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	506	GLY	-	expression tag	UNP Q5R372
B	506	GLY	-	expression tag	UNP Q5R372
C	506	GLY	-	expression tag	UNP Q5R372

- Molecule 2 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	14	Total	X	0	0
			14	14		
2	B	4	Total	X	0	0
			4	4		
2	C	14	Total	X	0	0
			14	14		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	O	0	0
			3	3		
3	C	10	Total	O	0	0
			10	10		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RAB GTPase-activating protein 1-like



- Molecule 1: RAB GTPase-activating protein 1-like



- Molecule 1: RAB GTPase-activating protein 1-like



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.09Å 64.57Å 290.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.30 20.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	92.3 (20.00-2.30) 92.1 (20.00-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.89 (at 2.30Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.244 , 0.274 (Not available) , 0.265	Depositor DCC
R_{free} test set	1287 reflections (3.12%)	wwPDB-VP
Wilson B-factor (Å ²)	49.4	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 46.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6473	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/2317	0.82	1/3130 (0.0%)
1	B	0.66	0/1923	0.83	2/2609 (0.1%)
1	C	0.77	0/2318	0.85	1/3132 (0.0%)
All	All	0.73	0/6558	0.83	4/8871 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	712	MSE	CG-SE-CE	-8.86	79.43	98.92
1	A	634	MSE	N-CA-C	6.02	113.64	108.22
1	B	708	PHE	CA-C-N	5.17	125.17	119.90
1	B	708	PHE	C-N-CA	5.17	125.17	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2270	0	2159	22	0
1	B	1887	0	1686	17	0
1	C	2271	0	2159	18	0
2	A	14	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	4	0	0	0	0
2	C	14	0	0	0	0
3	A	3	0	0	0	0
3	C	10	0	0	0	0
All	All	6473	0	6004	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:703:LEU:HD13	1:A:720:LEU:HD22	1.62	0.81
1:C:712:MSE:SE	1:C:716:ILE:HD11	2.33	0.77
1:C:712:MSE:SE	1:C:716:ILE:CD1	2.90	0.69
1:C:708:PHE:CD2	1:C:712:MSE:HE1	2.28	0.68
1:A:712:MSE:SE	1:A:716:ILE:HD11	2.52	0.59
1:C:617:ILE:HD12	1:C:620:CYS:SG	2.42	0.59
1:B:715:HIS:HB3	1:B:773:MSE:HG3	1.85	0.58
1:B:793:TYR:CE1	1:B:796:MSE:HE1	2.38	0.58
1:B:793:TYR:CZ	1:B:796:MSE:HE1	2.38	0.57
1:B:703:LEU:HD13	1:B:720:LEU:HD22	1.86	0.56
1:C:524:LEU:O	1:C:527:ARG:NH1	2.38	0.56
1:A:686:SER:HB3	1:B:686:SER:HB3	1.88	0.56
1:C:703:LEU:HD13	1:C:720:LEU:HD22	1.87	0.56
1:B:793:TYR:CE1	1:B:796:MSE:CE	2.89	0.54
1:B:673:LEU:HD11	1:B:786:LEU:HD21	1.88	0.54
1:A:704:PHE:CD1	1:A:716:ILE:HD13	2.43	0.54
1:A:657:ARG:HG2	1:A:658:ASN:CG	2.34	0.52
1:B:582:ILE:HD11	1:B:604:TYR:HA	1.91	0.52
1:B:704:PHE:CD1	1:B:716:ILE:HD13	2.43	0.52
1:A:650:TYR:O	1:A:725:LEU:HD12	2.10	0.51
1:C:712:MSE:SE	1:C:716:ILE:HD12	2.61	0.51
1:B:548:TRP:CZ3	1:B:630:LEU:HD23	2.47	0.50
1:A:582:ILE:HD11	1:A:604:TYR:HA	1.92	0.49
1:A:712:MSE:SE	1:A:716:ILE:CD1	3.10	0.49
1:A:741:GLU:O	1:A:745:GLN:HG2	2.12	0.49
1:A:772:LEU:HD23	1:A:773:MSE:HE2	1.94	0.49
1:C:712:MSE:HG2	1:C:773:MSE:HE3	1.94	0.49
1:A:657:ARG:HG2	1:A:658:ASN:ND2	2.27	0.48
1:A:647:MSE:HE3	1:A:653:ARG:HB3	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:704:PHE:HD1	1:A:716:ILE:HD13	1.80	0.47
1:A:620:CYS:O	1:A:623:GLN:HB2	2.14	0.47
1:B:677:GLN:NE2	1:B:780:LYS:O	2.48	0.45
1:B:712:MSE:HE2	1:B:772:LEU:CD2	2.47	0.45
1:C:555:HIS:N	1:C:555:HIS:ND1	2.65	0.45
1:C:723:GLU:OE1	1:C:727:ILE:HD11	2.17	0.44
1:A:662:ASP:O	1:A:666:LYS:HG3	2.18	0.44
1:A:652:LEU:HD13	1:A:725:LEU:HD21	2.00	0.43
1:C:704:PHE:CD1	1:C:716:ILE:HD13	2.54	0.43
1:C:517:LEU:HD23	1:C:550:LEU:HD12	2.01	0.43
1:B:705:THR:HA	1:B:713:VAL:HG11	2.01	0.43
1:C:545:ALA:HB2	1:C:633:HIS:O	2.19	0.43
1:A:704:PHE:CD1	1:A:716:ILE:CD1	3.03	0.42
1:A:545:ALA:HB2	1:A:633:HIS:O	2.20	0.42
1:A:561:LEU:HD12	1:A:637:GLU:HG3	2.01	0.42
1:C:582:ILE:HD11	1:C:604:TYR:HA	2.02	0.42
1:A:534:LEU:HD23	1:A:534:LEU:HA	1.90	0.41
1:B:566:ILE:O	1:B:566:ILE:HG22	2.19	0.41
1:C:690:LEU:HD11	1:C:752:LEU:HD21	2.02	0.41
1:C:512:SER:HB3	1:C:534:LEU:HD11	2.01	0.41
1:A:705:THR:HA	1:A:713:VAL:HG11	2.02	0.41
1:C:619:TYR:CE1	1:C:623:GLN:HB3	2.56	0.41
1:B:625:PHE:O	1:B:629:VAL:HG23	2.21	0.41
1:B:640:PHE:O	1:B:644:VAL:HG23	2.22	0.40
1:A:759:LEU:HB3	1:A:760:PRO:HD3	2.04	0.40
1:B:655:LEU:O	1:B:660:PHE:HA	2.21	0.40
1:C:647:MSE:HE3	1:C:653:ARG:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/310 (92%)	278 (98%)	7 (2%)	0	100	100
1	B	252/310 (81%)	242 (96%)	10 (4%)	0	100	100
1	C	286/310 (92%)	280 (98%)	5 (2%)	1 (0%)	36	46
All	All	823/930 (88%)	800 (97%)	22 (3%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	657	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/266 (86%)	225 (98%)	4 (2%)	53	72
1	B	164/266 (62%)	162 (99%)	2 (1%)	63	79
1	C	227/266 (85%)	224 (99%)	3 (1%)	61	77
All	All	620/798 (78%)	611 (98%)	9 (2%)	57	75

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	537	SER
1	A	572	SER
1	A	658	ASN
1	A	662	ASP
1	B	712	MSE
1	B	725	LEU
1	C	527	ARG
1	C	608	LYS
1	C	796	MSE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	583	HIS
1	A	600	GLN
1	A	658	ASN
1	A	659	ASN
1	A	675	GLN
1	B	623	GLN
1	C	621	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 32 ligands modelled in this entry, 32 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/310 (90%)	0.16	1 (0%) 88 89	28, 47, 67, 83	0
1	B	249/310 (80%)	0.75	25 (10%) 12 14	35, 68, 114, 157	0
1	C	280/310 (90%)	0.22	9 (3%) 50 52	30, 46, 68, 83	0
All	All	808/930 (86%)	0.36	35 (4%) 40 41	28, 50, 94, 157	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	648	TYR	4.0
1	C	525	GLY	3.8
1	B	590	ASP	3.5
1	B	579	THR	3.4
1	B	532	SER	3.3
1	C	526	ALA	3.2
1	B	564	TYR	3.2
1	B	588	ALA	3.1
1	C	532	SER	3.0
1	B	531	LEU	2.9
1	B	539	VAL	2.8
1	C	522	SER	2.7
1	B	540	PRO	2.6
1	A	711	CYS	2.5
1	B	571	ASP	2.5
1	B	566	ILE	2.5
1	B	594	ASP	2.5
1	B	612	VAL	2.3
1	B	567	LEU	2.3
1	B	535	VAL	2.3
1	B	586	PHE	2.2
1	B	629	VAL	2.3
1	C	711	CYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	555	HIS	2.2
1	B	604	TYR	2.2
1	B	576	SER	2.2
1	B	613	TYR	2.1
1	B	611	SER	2.1
1	B	568	ILE	2.1
1	B	596	GLY	2.1
1	C	530	GLY	2.1
1	B	610	TYR	2.1
1	C	520	TRP	2.1
1	C	584	ARG	2.0
1	B	534	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	UNX	A	905	1/1	-0.59	1.88	2,2,2,2	1
2	UNX	C	3	1/1	-0.58	2.14	2,2,2,2	1
2	UNX	C	909	1/1	-0.54	1.65	2,2,2,2	1
2	UNX	C	911	1/1	-0.43	1.45	2,2,2,2	1
2	UNX	A	4	1/1	-0.38	1.38	2,2,2,2	1
2	UNX	A	903	1/1	-0.37	1.43	2,2,2,2	1
2	UNX	C	6	1/1	-0.37	0.93	2,2,2,2	1
2	UNX	C	904	1/1	-0.35	1.38	2,2,2,2	1
2	UNX	A	906	1/1	-0.34	1.46	2,2,2,2	1
2	UNX	C	907	1/1	-0.33	1.19	2,2,2,2	1
2	UNX	A	910	1/1	-0.28	0.82	2,2,2,2	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UNX	A	908	1/1	-0.27	0.97	2,2,2,2	1
2	UNX	C	903	1/1	-0.27	1.27	2,2,2,2	1
2	UNX	C	908	1/1	-0.24	1.49	2,2,2,2	1
2	UNX	B	908	1/1	-0.22	1.00	2,2,2,2	1
2	UNX	C	5	1/1	-0.05	1.12	2,2,2,2	1
2	UNX	A	902	1/1	-0.04	1.29	2,2,2,2	1
2	UNX	A	907	1/1	0.06	1.76	2,2,2,2	1
2	UNX	C	905	1/1	0.06	1.76	2,2,2,2	1
2	UNX	B	910	1/1	0.20	1.24	2,2,2,2	1
2	UNX	A	2	1/1	0.29	1.19	2,2,2,2	1
2	UNX	A	909	1/1	0.30	1.34	2,2,2,2	1
2	UNX	B	902	1/1	0.31	1.86	2,2,2,2	1
2	UNX	A	911	1/1	0.33	1.52	2,2,2,2	1
2	UNX	A	1	1/1	0.34	1.25	2,2,2,2	1
2	UNX	C	902	1/1	0.34	1.51	2,2,2,2	1
2	UNX	C	906	1/1	0.53	1.53	2,2,2,2	1
2	UNX	A	904	1/1	0.59	1.41	2,2,2,2	1
2	UNX	A	901	1/1	0.67	1.27	2,2,2,2	1
2	UNX	B	901	1/1	0.72	1.18	2,2,2,2	1
2	UNX	C	901	1/1	0.87	1.22	2,2,2,2	1
2	UNX	C	910	1/1	0.89	1.20	2,2,2,2	1

6.5 Other polymers

There are no such residues in this entry.