



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:58 AM UTC

PDB ID : 3HZL / pdb_00003hzl
Title : Tyr258Phe mutant of NikD, an unusual amino acid oxidase essential for nikkomycin biosynthesis: open form at 1.55Å resolution
Authors : Mathews, F.S.; Jorns, M.S.; Carrell, C.J.
Deposited on : 2009-06-23
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	8% 86% 12% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3590 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NikD protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	394	3088	1959	544	575	10	0	0	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	258	PHE	TYR	engineered mutation	UNP Q9X9P9
A	390	LEU	-	expression tag	UNP Q9X9P9
A	391	GLU	-	expression tag	UNP Q9X9P9
A	392	HIS	-	expression tag	UNP Q9X9P9
A	393	HIS	-	expression tag	UNP Q9X9P9
A	394	HIS	-	expression tag	UNP Q9X9P9
A	395	HIS	-	expression tag	UNP Q9X9P9
A	396	HIS	-	expression tag	UNP Q9X9P9
A	397	HIS	-	expression tag	UNP Q9X9P9

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is PYRIDINE-2-CARBOXYLIC ACID (CCD ID: 6PC) (formula: $C_6H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			9	6	1	2		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	416	Total	O	0	0
			416	416		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NikD protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.39Å 90.53Å 85.97Å 90.00° 118.36° 90.00°	Depositor
Resolution (Å)	38.89 – 1.55 38.89 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.5 (38.89-1.55) 99.6 (38.89-1.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.18 (at 1.55Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.225 , 0.244 0.209 , 0.210	Depositor DCC
R_{free} test set	4324 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.024 for k,h,-1/2*h-1/2*k-l 0.014 for -k,-h,-1/2*h+1/2*k-l 0.024 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3590	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, FAD, 6PC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.78	0/3172	1.09	15/4326 (0.3%)

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	356	ALA	N-CA-C	9.72	125.52	112.30
1	A	129	ARG	N-CA-C	7.24	119.25	111.36
1	A	238	THR	N-CA-C	-6.60	100.99	111.02
1	A	307	HIS	N-CA-C	6.19	121.77	113.72
1	A	57	THR	N-CA-C	-6.13	105.84	113.38
1	A	38	PHE	N-CA-C	-5.77	102.37	110.50
1	A	355	TRP	N-CA-C	-5.73	96.45	107.57
1	A	350	VAL	N-CA-C	5.71	116.09	107.75
1	A	198	GLY	N-CA-C	-5.62	105.69	112.49
1	A	144	GLN	N-CA-C	-5.50	98.79	108.69
1	A	95	THR	N-CA-C	5.41	119.62	113.19
1	A	49	GLU	N-CA-C	5.40	117.40	109.24
1	A	312	ASP	N-CA-C	-5.39	101.45	109.42
1	A	242	PHE	N-CA-C	-5.09	102.85	110.23
1	A	315	PRO	N-CA-C	5.00	118.95	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3088	0	2970	35	0
2	A	53	0	30	1	0
3	A	9	0	4	0	0
4	A	24	0	42	3	0
5	A	416	0	0	8	0
All	All	3590	0	3046	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:400:MPD:H11	5:A:1174:HOH:O	1.90	0.71
1:A:390:LEU:HD12	1:A:390:LEU:H	1.56	0.70
1:A:213:GLU:HB3	1:A:214:PRO:HD3	1.79	0.64
1:A:268:GLY:HA2	5:A:1109:HOH:O	1.98	0.64
1:A:255:ASN:HD22	1:A:299:ARG:HH12	1.47	0.62
1:A:55:GLN:HE21	1:A:55:GLN:H	1.48	0.61
1:A:255:ASN:ND2	1:A:299:ARG:HH12	1.99	0.60
1:A:55:GLN:H	1:A:55:GLN:NE2	2.00	0.59
1:A:390:LEU:HD12	1:A:390:LEU:N	2.17	0.59
1:A:85:HIS:HD2	5:A:1323:HOH:O	1.89	0.55
1:A:53:ARG:HD2	1:A:55:GLN:O	2.08	0.54
1:A:122:LYS:HE2	5:A:1285:HOH:O	2.09	0.52
1:A:123:ALA:O	1:A:127:GLU:HG3	2.09	0.52
1:A:331:ARG:NE	1:A:386:GLN:HG3	2.26	0.50
1:A:57:THR:HA	1:A:110:MET:HG2	1.94	0.49
1:A:84:ILE:CD1	1:A:150:ILE:HG12	2.43	0.49
1:A:232:ARG:HD2	5:A:1164:HOH:O	2.13	0.49
1:A:331:ARG:CD	1:A:386:GLN:HG3	2.42	0.49
1:A:49:GLU:OE2	1:A:272:ARG:HD2	2.13	0.49
1:A:213:GLU:HG2	5:A:1363:HOH:O	2.13	0.48
1:A:390:LEU:H	1:A:390:LEU:CD1	2.24	0.48
1:A:84:ILE:HD12	1:A:150:ILE:HG12	1.95	0.48
1:A:199:LYS:HD2	1:A:371:ALA:O	2.15	0.47
1:A:73:ARG:HH11	4:A:402:MPD:H52	1.80	0.47
1:A:103:GLN:HE21	1:A:106:GLY:H	1.62	0.47
1:A:21:TRP:CD1	1:A:25:GLU:HG3	2.51	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:297:MET:HA	1:A:297:MET:HE2	1.97	0.45
1:A:35:ARG:HD2	2:A:398:FAD:C5A	2.49	0.43
1:A:55:GLN:O	1:A:55:GLN:HG2	2.18	0.43
1:A:132:PHE:HA	1:A:307:HIS:O	2.19	0.43
1:A:395:HIS:HA	5:A:1169:HOH:O	2.19	0.42
1:A:274:GLY:HA2	1:A:300:LEU:HD22	2.02	0.41
1:A:53:ARG:HD2	1:A:53:ARG:HH11	1.74	0.41
1:A:101:GLU:OE2	4:A:400:MPD:O4	2.37	0.41
1:A:137:ARG:HG3	5:A:1245:HOH:O	2.21	0.41
1:A:235:THR:HG23	1:A:236:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/397 (99%)	387 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/322 (99%)	316 (99%)	3 (1%)	70	51

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	GLU
1	A	55	GLN
1	A	237	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	55	GLN
1	A	85	HIS
1	A	103	GLN
1	A	248	GLN
1	A	252	GLN
1	A	255	ASN
1	A	263	ASN
1	A	296	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MPD	A	401	-	7,7,7	0.50	0	9,10,10	0.36	0
4	MPD	A	400	-	7,7,7	0.65	0	9,10,10	0.26	0
2	FAD	A	398	1	58,58,58	1.53	10 (17%)	85,89,89	1.06	4 (4%)
4	MPD	A	402	-	7,7,7	0.84	0	9,10,10	0.41	0
3	6PC	A	399	-	9,9,9	2.39	3 (33%)	11,11,11	1.35	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	A	401	-	-	0/5/5/5	-
4	MPD	A	400	-	-	0/5/5/5	-
2	FAD	A	398	1	-	0/34/50/50	0/6/6/6
4	MPD	A	402	-	-	0/5/5/5	-
3	6PC	A	399	-	-	4/4/4/4	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	398	FAD	C4X-N5	4.78	1.41	1.30
3	A	399	6PC	C3-N2	4.01	1.43	1.34
3	A	399	6PC	C1-N2	3.93	1.41	1.34
2	A	398	FAD	C5'-C4'	-3.37	1.47	1.51
2	A	398	FAD	C2A-N1A	3.14	1.39	1.33
2	A	398	FAD	C1'-C2'	-3.13	1.48	1.52
2	A	398	FAD	C9A-C5X	3.09	1.46	1.41
2	A	398	FAD	C2A-N3A	2.70	1.38	1.33
2	A	398	FAD	P-O3P	-2.66	1.56	1.59
2	A	398	FAD	C10-N10	2.39	1.42	1.37
2	A	398	FAD	O4-C4	2.39	1.28	1.23
3	A	399	6PC	C6-C1	2.30	1.43	1.39
2	A	398	FAD	C9-C8	2.02	1.42	1.39

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	398	FAD	C4-N3-C2	-2.97	120.37	125.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	399	6PC	O2-C2-C1	2.61	120.91	114.71
2	A	398	FAD	O5B-PA-O1A	-2.29	99.88	108.94
2	A	398	FAD	O2A-PA-O1A	2.28	123.05	112.44
2	A	398	FAD	C9A-C9-C8	2.26	123.77	119.22
3	A	399	6PC	O2-C2-O1	-2.19	118.64	123.35

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	399	6PC	C6-C1-C2-O1
3	A	399	6PC	C6-C1-C2-O2
3	A	399	6PC	N2-C1-C2-O1
3	A	399	6PC	N2-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	400	MPD	2	0
2	A	398	FAD	1	0
4	A	402	MPD	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/397 (99%)	0.74	32 (8%) 18 21	12, 23, 42, 53	1 (0%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	237	VAL	5.4
1	A	389	LEU	4.3
1	A	372	VAL	4.3
1	A	390	LEU	4.2
1	A	395	HIS	3.7
1	A	115	SER	3.4
1	A	388	ALA	3.4
1	A	4	SER	3.3
1	A	238	THR	3.3
1	A	374	ASP	3.2
1	A	120	TRP	3.1
1	A	130	PHE	2.8
1	A	114	LEU	2.7
1	A	121	LEU	2.7
1	A	123	ALA	2.7
1	A	116	VAL	2.7
1	A	137	ARG	2.7
1	A	218	ARG	2.6
1	A	108	ALA	2.5
1	A	236	PRO	2.5
1	A	59	GLU	2.4
1	A	373	GLU	2.4
1	A	118	TYR	2.3
1	A	392	HIS	2.2
1	A	251	PRO	2.1
1	A	124	THR	2.1
1	A	391	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	384	ALA	2.1
1	A	113	LYS	2.1
1	A	371	ALA	2.0
1	A	283	ASP	2.0
1	A	279	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MPD	A	402	8/8	0.62	0.30	42,45,48,49	0
4	MPD	A	400	8/8	0.80	0.20	35,37,40,46	0
4	MPD	A	401	8/8	0.88	0.17	30,35,38,38	0
3	6PC	A	399	9/9	0.95	0.08	22,23,26,28	0
2	FAD	A	398	53/53	0.98	0.06	12,15,21,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.