



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 09:55 AM UTC

PDB ID : 2I0L / pdb_00002i0l
Title : X-ray crystal structure of Sap97 PDZ2 bound to the C-terminal peptide of HPV18 E6.
Authors : Chen, X.S.; Zhang, Y.; Dasgupta, J.; Banks, L.; Thomas, M.
Deposited on : 2006-08-10
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	84	
1	B	84	
2	C	7	
2	D	7	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1472 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Disks large homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	83	Total	C	N	O	S	0	0	0
			609	387	104	117	1			
1	B	83	Total	C	N	O	S	0	0	0
			602	382	102	117	1			

- Molecule 2 is a protein called peptide E6.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	6	Total	C	N	O	0	0	0
			49	28	10	11			
2	D	5	Total	C	N	O	0	0	0
			44	25	9	10			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	88	Total	O	0	0
			88	88		
3	C	8	Total	O	0	0
			8	8		
3	B	66	Total	O	0	0
			66	66		
3	D	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Disks large homolog 1




- Molecule 1: Disks large homolog 1



- Molecule 2: peptide E6



- Molecule 2: peptide E6



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	90.90Å 44.18Å 52.48Å 90.00° 121.85° 90.00°	Depositor
Resolution (Å)	26.20 – 2.31 26.20 – 2.31	Depositor EDS
% Data completeness (in resolution range)	88.5 (26.20-2.31) 88.5 (26.20-2.31)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.32Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.239 , 0.302 0.239 , 0.299	Depositor DCC
R_{free} test set	387 reflections (5.54%)	wwPDB-VP
Wilson B-factor (Å ²)	33.2	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 67.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1472	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	4/617 (0.6%)	1.32	10/831 (1.2%)
1	B	1.81	3/610 (0.5%)	1.53	10/824 (1.2%)
2	C	1.81	2/48 (4.2%)	1.76	2/62 (3.2%)
2	D	0.50	0/43	1.36	1/55 (1.8%)
All	All	1.40	9/1318 (0.7%)	1.44	23/1772 (1.3%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	391	LYS	C-N	-41.06	0.76	1.33
1	A	363	LYS	CA-C	8.16	1.63	1.52
1	A	364	LEU	N-CA	6.96	1.55	1.46
2	C	2002	ARG	CA-C	6.67	1.60	1.52
1	B	375	SER	N-CA	-6.39	1.39	1.46
2	C	2003	GLU	N-CA	6.31	1.53	1.46
1	B	374	ASN	CA-C	-5.15	1.45	1.52
1	A	363	LYS	N-CA	5.12	1.52	1.46
1	A	363	LYS	C-N	5.07	1.40	1.33

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	391	LYS	O-C-N	-19.01	97.52	122.42
1	B	391	LYS	CA-C-N	14.84	149.88	121.54
1	B	391	LYS	C-N-CA	14.84	149.88	121.54
1	A	376	VAL	N-CA-C	-11.74	90.00	107.37
1	A	365	GLN	OE1-CD-NE2	-9.35	113.25	122.60
1	B	390	LEU	CA-C-N	-6.96	110.22	122.26
1	B	390	LEU	C-N-CA	-6.96	110.22	122.26
2	C	2004	THR	N-CA-C	6.92	120.68	109.40
1	A	365	GLN	CG-CD-NE2	6.73	126.49	116.40
1	A	363	LYS	CA-C-N	6.67	131.97	122.09
1	A	363	LYS	C-N-CA	6.67	131.97	122.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	374	ASN	N-CA-C	-6.66	101.78	111.17
2	D	2004	THR	N-CA-C	6.16	119.44	109.40
1	A	324	GLY	CA-C-N	5.82	127.12	119.84
1	A	324	GLY	C-N-CA	5.82	127.12	119.84
2	C	2002	ARG	N-CA-C	5.66	117.63	108.41
1	B	362	GLY	N-CA-C	5.49	122.41	114.67
1	A	332	ILE	N-CA-C	5.38	117.63	108.86
1	B	397	VAL	N-CA-C	5.38	116.03	108.17
1	B	340	HIS	N-CA-C	-5.36	105.09	111.69
1	A	327	GLY	N-CA-C	5.34	118.06	110.74
1	B	375	SER	CA-C-N	-5.30	115.89	123.10
1	B	375	SER	C-N-CA	-5.30	115.89	123.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	609	0	621	44	1
1	B	602	0	600	31	0
2	C	49	0	44	8	0
2	D	44	0	42	6	0
3	A	88	0	0	13	0
3	B	66	0	0	4	0
3	C	8	0	0	2	0
3	D	6	0	0	0	0
All	All	1472	0	1307	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 31.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:373:VAL:HG21	1:B:390:LEU:HD23	1.35	1.01
1:B:355:GLY:H	1:B:359:HIS:HD2	1.20	0.88
1:A:373:VAL:HG23	1:A:378:LEU:HD11	1.59	0.85
1:A:373:VAL:O	1:A:376:VAL:HG22	1.76	0.85
1:A:361:ASP:O	1:A:361:ASP:CG	2.24	0.80
1:B:355:GLY:H	1:B:359:HIS:CD2	2.00	0.79
1:A:350:THR:O	1:A:366:ILE:HD13	1.85	0.76
1:B:373:VAL:HG21	1:B:390:LEU:CD2	2.13	0.75
1:B:390:LEU:HD13	2:D:2006:VAL:HG21	1.71	0.72
1:A:328:LEU:HD12	2:C:2006:VAL:HG12	1.71	0.71
1:A:326:LYS:HB2	3:A:2029:HOH:O	1.91	0.70
1:A:393:THR:O	1:A:394:SER:O	2.11	0.68
1:A:355:GLY:HA3	3:A:2089:HOH:O	1.94	0.66
1:B:373:VAL:O	1:B:375:SER:N	2.29	0.65
2:D:2006:VAL:HG12	2:D:2006:VAL:O	1.98	0.63
1:A:391:LYS:HD2	3:A:2242:HOH:O	1.97	0.62
1:A:373:VAL:CG2	1:A:378:LEU:HD11	2.29	0.61
1:B:373:VAL:HG22	1:B:399:LEU:CD2	2.30	0.61
1:B:373:VAL:HG23	1:B:378:LEU:HD11	1.83	0.61
1:B:371:LEU:H	1:B:400:LYS:HB2	1.65	0.61
1:A:390:LEU:HB3	2:C:2006:VAL:HG11	1.82	0.61
1:B:338:ASN:HB2	3:B:2061:HOH:O	2.01	0.60
1:A:383:HIS:O	1:A:387:VAL:HG23	2.00	0.60
1:B:354:GLU:HG3	3:B:2011:HOH:O	2.01	0.60
1:A:376:VAL:O	1:A:376:VAL:HG23	2.02	0.59
1:A:387:VAL:HG12	3:A:2242:HOH:O	2.02	0.59
1:A:361:ASP:O	1:A:361:ASP:OD1	2.20	0.59
1:A:336:VAL:HG22	1:A:382:THR:HA	1.85	0.58
1:B:390:LEU:HB3	2:D:2006:VAL:HG11	1.86	0.58
1:B:373:VAL:CG2	1:B:390:LEU:HD23	2.24	0.56
1:A:347:ILE:HD12	1:A:378:LEU:HB3	1.88	0.55
1:A:373:VAL:O	1:A:374:ASN:C	2.48	0.55
1:A:326:LYS:HB3	1:A:326:LYS:NZ	2.23	0.54
1:B:329:GLY:HA2	1:B:353:ILE:HD12	1.89	0.54
1:A:350:THR:O	1:A:366:ILE:CD1	2.56	0.53
1:A:362:GLY:O	1:A:364:LEU:N	2.41	0.53
1:B:340:HIS:C	1:B:341:ILE:HD12	2.33	0.53
1:A:384:GLU:OE1	2:C:2002:ARG:NH2	2.41	0.53
1:A:326:LYS:HG2	1:A:326:LYS:O	2.10	0.51
1:A:347:ILE:HD11	1:A:386:ALA:HB2	1.92	0.51
2:C:2002:ARG:NH1	3:C:2184:HOH:O	2.33	0.50
1:A:388:THR:HA	3:A:2242:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:318:GLU:HG3	3:B:2125:HOH:O	2.12	0.50
1:A:374:ASN:HA	3:A:2087:HOH:O	2.11	0.49
1:A:376:VAL:O	1:A:376:VAL:CG2	2.61	0.49
1:B:373:VAL:O	1:B:376:VAL:N	2.29	0.49
1:A:372:ALA:HB1	1:A:375:SER:HA	1.94	0.48
1:B:390:LEU:HB3	2:D:2006:VAL:CG1	2.44	0.48
1:A:395:ASP:HA	3:A:2103:HOH:O	2.14	0.47
1:A:355:GLY:C	3:A:2089:HOH:O	2.57	0.47
1:A:359:HIS:CE1	3:A:2078:HOH:O	2.66	0.47
1:A:355:GLY:CA	3:A:2089:HOH:O	2.59	0.47
1:A:364:LEU:O	1:A:365:GLN:NE2	2.47	0.47
1:B:328:LEU:HD12	2:D:2006:VAL:HG12	1.97	0.47
1:A:390:LEU:HB3	2:C:2006:VAL:CG1	2.44	0.46
2:C:2004:THR:HG22	2:C:2005:GLN:N	2.30	0.45
1:B:360:LYS:HB3	1:B:360:LYS:HE3	1.73	0.45
1:A:346:SER:OG	1:A:369:LYS:HE3	2.17	0.45
1:A:359:HIS:HE1	3:A:2078:HOH:O	2.00	0.45
1:B:349:VAL:HG21	1:B:364:LEU:HD11	1.98	0.45
1:B:373:VAL:HG22	1:B:399:LEU:HD21	1.99	0.45
1:B:341:ILE:HD12	1:B:341:ILE:N	2.32	0.44
1:A:318:GLU:HA	1:A:399:LEU:O	2.16	0.44
1:A:374:ASN:HB3	3:A:2124:HOH:O	2.17	0.44
1:A:394:SER:N	3:A:2146:HOH:O	2.51	0.44
1:B:340:HIS:HB3	1:B:341:ILE:HD12	2.00	0.44
1:B:382:THR:HG23	1:B:385:GLU:OE2	2.17	0.44
1:A:330:PHE:HE1	1:A:332:ILE:CG2	2.31	0.43
1:B:365:GLN:HB2	1:B:368:ASP:OD1	2.18	0.43
1:B:382:THR:OG1	1:B:385:GLU:HG3	2.18	0.43
1:A:390:LEU:CB	2:C:2006:VAL:HG11	2.48	0.42
1:B:336:VAL:HA	1:B:345:ASN:OD1	2.19	0.42
1:B:373:VAL:CG2	1:B:390:LEU:CD2	2.92	0.42
2:D:2006:VAL:O	2:D:2006:VAL:CG1	2.67	0.41
1:A:351:LYS:HA	1:A:366:ILE:HD11	2.02	0.41
1:B:379:GLU:O	1:B:380:GLU:HB2	2.19	0.41
1:A:361:ASP:OD1	1:A:361:ASP:C	2.59	0.41
1:B:384:GLU:HA	3:B:2118:HOH:O	2.21	0.41
1:A:320:LYS:O	1:A:321:LEU:HD13	2.22	0.40
2:C:2001:ARG:N	3:C:2160:HOH:O	2.54	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:GLU:OE2	1:A:384:GLU:OE2[2_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	81/84 (96%)	70 (86%)	7 (9%)	4 (5%)	1	1
1	B	81/84 (96%)	70 (86%)	10 (12%)	1 (1%)	10	11
2	C	4/7 (57%)	4 (100%)	0	0	100	100
2	D	3/7 (43%)	3 (100%)	0	0	100	100
All	All	169/182 (93%)	147 (87%)	17 (10%)	5 (3%)	3	2

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	363	LYS
1	A	394	SER
1	B	374	ASN
1	A	375	SER
1	A	325	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	64/66 (97%)	56 (88%)	8 (12%)	4	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	62/66 (94%)	60 (97%)	2 (3%)	34	50
2	C	5/7 (71%)	3 (60%)	2 (40%)	0	0
2	D	5/7 (71%)	3 (60%)	2 (40%)	0	0
All	All	136/146 (93%)	122 (90%)	14 (10%)	7	8

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	321	LEU
1	A	325	PRO
1	A	326	LYS
1	A	338	ASN
1	A	354	GLU
1	A	361	ASP
1	A	366	ILE
1	A	379	GLU
2	C	2002	ARG
2	C	2005	GLN
1	B	319	ILE
1	B	365	GLN
2	D	2002	ARG
2	D	2004	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	392	ASN
2	C	2005	GLN
1	B	359	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	391:LYS	C	392:ASN	N	0.76

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	83/84 (98%)	0.24	1 (1%) 76 78	16, 26, 39, 45	0
1	B	83/84 (98%)	0.12	0 100 100	19, 28, 39, 48	0
2	C	6/7 (85%)	0.81	1 (16%) 4 5	30, 36, 61, 64	0
2	D	5/7 (71%)	0.88	1 (20%) 3 3	33, 37, 45, 55	0
All	All	177/182 (97%)	0.22	3 (1%) 69 71	16, 28, 44, 64	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	362	GLY	2.5
2	C	2001	ARG	2.2
2	D	2006	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.