



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:58 AM UTC

PDB ID : 2I0Q / pdb_00002i0q
Title : Crystal structure of a telomere single-strand DNA-protein complex from O. nova with full-length alpha and beta telomere proteins
Authors : Horvath, M.P.; Buczek, P.
Deposited on : 2006-08-11
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	11	7% 55% 36% 9%
2	A	495	7% 82% 10% 7%
3	B	385	21% 47% 8% 44%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	A	497	-	-	X	-
5	EDO	A	502	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6019 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*GP*GP*TP*TP*TP*TP*GP*GP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	11	Total	C	N	O	P	0	0	0
			231	110	43	68	10			

- Molecule 2 is a protein called Telomere-binding protein alpha subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	461	Total	C	N	O	S	0	12	0
			3750	2381	638	729	2			

- Molecule 3 is a protein called Telomere-binding protein beta subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	B	216	Total	C	N	O	S	0	3	0
			1705	1095	283	326	1			

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	Cl	0	0
			4	4		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	19	Total	O	0	0
			19	19		
6	A	241	Total	O	0	2
			243	243		
6	B	46	Total	O	0	1
			47	47		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*GP*GP*GP*TP*TP*TP*TP*GP*GP*GP*G)-3'

Chain D: 





- Molecule 2: Telomere-binding protein alpha subunit

Chain A: 



- Molecule 3: Telomere-binding protein beta subunit

Chain B: 



THR	ASP	LYS	MET	THR	MET	ALA	GLN	PHE	VAL	LYS	TYR	LEU	ASP	TRP	HIS	GLU	LYS	LYS	LYS	GLY	GLY	LYS	VAL	SER	SER	GLY	GLY	LYS	VAL	LEU	GLY	LYS	ARG	SER	ALA	GLY	LYS	ALA	SER	ALA	THR	SER	GLY	LYS	ALA	SER	LYS	LYS	ALA	SER	LYS	LYS	THR	ALA	ALA	LYS	LYS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.33Å 93.33Å 423.78Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 1.91 50.00 – 1.91	Depositor EDS
% Data completeness (in resolution range)	98.7 (50.00-1.91) 98.9 (50.00-1.91)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.92 (at 1.91Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.243 , 0.264 0.233 , 0.251	Depositor DCC
R_{free} test set	8402 reflections (9.88%)	wwPDB-VP
Wilson B-factor (Å ²)	34.8	Xtriage
Anisotropy	0.337	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 32.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6019	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	D	0.31	0/259	0.68	1/401 (0.2%)
2	A	0.43	0/3826	0.92	16/5188 (0.3%)
3	B	0.37	0/1741	0.81	1/2366 (0.0%)
All	All	0.41	0/5826	0.87	18/7955 (0.2%)

There are no bond length outliers.

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	240	THR	N-CA-C	6.62	120.30	109.24
2	A	63	PHE	N-CA-C	-6.62	101.16	110.29
2	A	89	LYS	N-CA-C	-6.61	105.28	113.15
2	A	91	ALA	N-CA-C	6.57	122.80	113.02
2	A	435	THR	N-CA-C	6.47	120.10	112.72
2	A	406	LYS	N-CA-C	-6.46	104.97	112.92
3	B	74	LYS	N-CA-C	-6.40	106.43	114.56
2	A	417	VAL	N-CA-C	6.23	118.13	109.29
2	A	62	THR	N-CA-C	-6.03	102.76	110.53
2	A	41	LEU	N-CA-C	5.99	118.30	111.11
2	A	312	LYS	N-CA-C	5.62	117.48	111.36
2	A	432	LEU	N-CA-C	5.50	119.03	110.17
2	A	335	LEU	N-CA-C	5.32	119.03	112.54
2	A	93	ASP	N-CA-C	5.21	121.90	110.80
2	A	119	ILE	N-CA-C	5.21	116.08	108.53
1	D	6	DG	C2'-C3'-O3'	-5.07	103.89	111.50
2	A	428[A]	THR	N-CA-C	-5.04	101.53	109.50
2	A	428[B]	THR	N-CA-C	-5.04	101.53	109.50

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	231	0	127	4	0
2	A	3750	0	3636	29	0
3	B	1705	0	1615	28	0
4	A	4	0	0	2	0
5	A	16	0	24	5	0
5	B	4	0	6	1	0
6	A	243	0	0	3	0
6	B	47	0	0	0	0
6	D	19	0	0	0	0
All	All	6019	0	5408	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:10:GLN:HE21	3:B:12:SER:H	1.05	1.01
3:B:130:ASN:HD22	3:B:131:PRO:HD2	1.30	0.93
3:B:17[B]:LEU:HG	3:B:95:LEU:HD11	1.60	0.82
2:A:85:LEU:H	5:A:502:EDO:H21	1.50	0.77
4:A:497:CL:CL	3:B:182:LYS:HE3	2.25	0.73
3:B:71:PHE:HE1	3:B:123:PRO:HD3	1.59	0.68
3:B:62:TYR:HB3	3:B:212:THR:HG21	1.76	0.66
2:A:345:LEU:HD12	2:A:346:PRO:HD2	1.79	0.65
2:A:84:TYR:H	5:A:502:EDO:H21	1.62	0.63
3:B:71:PHE:CE1	3:B:123:PRO:HD3	2.33	0.63
2:A:88:GLN:HE22	2:A:92:GLY:C	2.06	0.62
3:B:130:ASN:ND2	3:B:131:PRO:HD2	2.11	0.60
2:A:390:TYR:HD2	2:A:392:ARG:HH21	1.47	0.59
1:D:6:DG:H3'	1:D:7:DG:H5''	1.86	0.58
2:A:84:TYR:H	5:A:502:EDO:C2	2.17	0.57
3:B:10:GLN:NE2	3:B:12:SER:H	1.88	0.57
3:B:21:LEU:HD22	3:B:95:LEU:HB3	1.85	0.57
2:A:281:THR:HB	3:B:10:GLN:HE22	1.71	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:98:ARG:NH1	3:B:98:ARG:HB3	2.22	0.55
2:A:124:ARG:NH1	2:A:143:SER:O	2.41	0.54
3:B:68:VAL:HG21	3:B:213:LEU:HD12	1.90	0.54
3:B:98:ARG:HB3	3:B:98:ARG:HH11	1.73	0.54
2:A:85:LEU:HD11	2:A:93:ASP:HB3	1.89	0.53
3:B:130:ASN:HD22	3:B:131:PRO:CD	2.09	0.53
3:B:167:ASP:HA	5:B:386:EDO:O2	2.08	0.53
3:B:132[B]:THR:OG1	3:B:133:ARG:N	2.41	0.53
3:B:217:ASP:O	3:B:220:VAL:HG22	2.09	0.53
2:A:388:LYS:HE2	6:A:578:HOH:O	2.09	0.52
3:B:17[A]:LEU:HD11	3:B:93:TRP:CH2	2.45	0.52
2:A:88:GLN:HE22	2:A:93:ASP:N	2.09	0.51
1:D:12:DT:H71	1:D:16:DG:H2'	1.92	0.51
2:A:83:LEU:HA	5:A:502:EDO:H22	1.94	0.50
4:A:497:CL:CL	3:B:182:LYS:CE	2.98	0.49
2:A:36:TYR:OH	2:A:124:ARG:NH2	2.47	0.47
2:A:88:GLN:OE1	2:A:93:ASP:HA	2.15	0.46
3:B:65:LYS:HA	3:B:213:LEU:HD11	1.97	0.46
2:A:51[B]:PRO:HB3	2:A:124:ARG:HG2	1.98	0.46
2:A:88:GLN:C	2:A:90:GLY:H	2.24	0.46
3:B:157:GLN:HA	3:B:160:ILE:HG22	1.97	0.45
3:B:128:ARG:HA	3:B:128:ARG:HD2	1.42	0.45
3:B:17[B]:LEU:HD21	3:B:93:TRP:CH2	2.52	0.45
3:B:129:LEU:HD12	3:B:129:LEU:HA	1.87	0.45
2:A:211:TYR:O	5:A:501:EDO:H12	2.18	0.44
1:D:8:DG:N3	1:D:8:DG:H2'	2.31	0.44
2:A:46:LEU:HD23	2:A:127:LEU:HG	2.00	0.43
2:A:403:ALA:HB2	6:A:652:HOH:O	2.19	0.42
2:A:392:ARG:H	2:A:392:ARG:HG2	1.56	0.42
2:A:88:GLN:NE2	2:A:92:GLY:C	2.76	0.42
1:D:6:DG:H3'	1:D:7:DG:C5'	2.49	0.42
2:A:423:GLN:HA	6:A:653:HOH:O	2.20	0.41
2:A:48[B]:SER:C	2:A:50[B]:GLN:N	2.77	0.41
3:B:92:ASN:HB3	3:B:119:HIS:HB2	2.03	0.41
2:A:66:LYS:HE2	2:A:68[B]:ASN:O	2.21	0.41
2:A:48[B]:SER:C	2:A:50[B]:GLN:H	2.29	0.40
2:A:180:THR:C	2:A:181:ILE:HD12	2.46	0.40
3:B:98:ARG:HG3	3:B:115:ARG:HD2	2.04	0.40
3:B:90:ILE:HG23	3:B:118:VAL:HG13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	472/495 (95%)	448 (95%)	21 (4%)	3 (1%)	21	10
3	B	217/385 (56%)	204 (94%)	13 (6%)	0	100	100
All	All	689/880 (78%)	652 (95%)	34 (5%)	3 (0%)	30	19

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	405	GLY
2	A	69	GLN
2	A	404	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	402/435 (92%)	391 (97%)	11 (3%)	39	24
3	B	177/308 (58%)	167 (94%)	10 (6%)	19	6
All	All	579/743 (78%)	558 (96%)	21 (4%)	36	15

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	35	LYS
2	A	129	LEU
2	A	168[A]	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	168[B]	ASP
2	A	240	THR
2	A	309	LEU
2	A	353	LEU
2	A	392	ARG
2	A	416	LEU
2	A	437	ASP
2	A	467	LEU
3	B	17[A]	LEU
3	B	17[B]	LEU
3	B	33[A]	SER
3	B	33[B]	SER
3	B	98	ARG
3	B	128	ARG
3	B	132[A]	THR
3	B	132[B]	THR
3	B	156	LEU
3	B	182	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	133	GLN
2	A	163	GLN
2	A	284	GLN
2	A	327	ASN
2	A	364	GLN
3	B	10	GLN
3	B	124	ASN
3	B	130	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	A	502	-	3,3,3	0.57	0	2,2,2	0.54	0
5	EDO	A	503	-	3,3,3	0.58	0	2,2,2	0.52	0
5	EDO	B	386	-	3,3,3	0.58	0	2,2,2	0.61	0
5	EDO	A	500	-	3,3,3	0.41	0	2,2,2	0.42	0
5	EDO	A	501	-	3,3,3	0.49	0	2,2,2	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	502	-	-	0/1/1/1	-
5	EDO	A	503	-	-	1/1/1/1	-
5	EDO	B	386	-	-	1/1/1/1	-
5	EDO	A	500	-	-	0/1/1/1	-
5	EDO	A	501	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	503	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
5	B	386	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	502	EDO	4	0
5	B	386	EDO	1	0
5	A	501	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	11/11 (100%)	-0.08	0 100 100	26, 33, 57, 61	0
2	A	461/495 (93%)	0.56	36 (7%) 19 22	11, 33, 58, 86	12 (2%)
3	B	216/385 (56%)	1.71	82 (37%) 1 0	24, 57, 82, 86	3 (1%)
All	All	688/891 (77%)	0.91	118 (17%) 4 5	11, 38, 78, 86	15 (2%)

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	91	ALA	7.2
3	B	132[A]	THR	7.0
2	A	68[A]	ASN	6.8
2	A	142[A]	TYR	6.8
3	B	86	LYS	6.0
2	A	405	GLY	5.9
3	B	131	PRO	5.6
3	B	125	LEU	5.5
3	B	129	LEU	5.5
2	A	49[A]	ALA	5.5
3	B	126	GLN	5.5
2	A	406	LYS	5.4
2	A	89	LYS	5.3
2	A	90	GLY	5.2
3	B	87	VAL	5.1
2	A	403	ALA	4.6
3	B	77	ASN	4.5
2	A	407	GLY	4.3
3	B	198	ASP	4.3
2	A	166	VAL	4.2
3	B	80	ILE	4.2
3	B	219	PHE	4.2
3	B	134	TYR	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	B	65	LYS	4.2
2	A	92	GLY	4.1
3	B	194	ALA	4.1
3	B	41	CYS	4.1
2	A	69	GLN	4.0
3	B	71	PHE	4.0
2	A	404	SER	4.0
2	A	50[A]	GLN	3.9
3	B	216	ALA	3.7
3	B	81	VAL	3.7
2	A	316	ASP	3.7
3	B	218	ILE	3.6
3	B	220	VAL	3.5
2	A	87	GLN	3.5
2	A	165	ALA	3.5
3	B	67	ALA	3.5
2	A	35	LYS	3.4
3	B	121	PHE	3.4
3	B	73	ALA	3.4
3	B	118	VAL	3.2
2	A	88	GLN	3.2
3	B	78	VAL	3.2
2	A	283	THR	3.1
3	B	83	LEU	3.1
3	B	63	PHE	3.1
3	B	123	PRO	3.1
3	B	90	ILE	3.1
3	B	215	ILE	3.1
3	B	75	PHE	3.0
3	B	224	GLY	3.0
2	A	392	ARG	3.0
3	B	213	LEU	2.9
3	B	197	GLY	2.9
3	B	169	LEU	2.9
3	B	164	VAL	2.9
3	B	88	ILE	2.9
3	B	76	PRO	2.8
3	B	211	ALA	2.8
3	B	122	LYS	2.8
3	B	39	LEU	2.8
3	B	120	SER	2.7
3	B	128	ARG	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	94	ALA	2.7
3	B	9	GLN	2.7
3	B	17[A]	LEU	2.7
3	B	69	ASN	2.7
2	A	93	ASP	2.6
2	A	315	ASP	2.6
3	B	221	GLN	2.6
3	B	205	PHE	2.6
3	B	79	ASN	2.6
3	B	89	VAL	2.6
3	B	35	LEU	2.6
3	B	40	LYS	2.5
2	A	311	ALA	2.5
3	B	72	HIS	2.5
3	B	11	GLN	2.5
3	B	177	ASP	2.5
3	B	74	LYS	2.5
3	B	223	LYS	2.5
3	B	119	HIS	2.5
3	B	175	VAL	2.5
3	B	210	THR	2.5
3	B	61	PRO	2.5
2	A	70	GLU	2.4
3	B	105	VAL	2.4
3	B	206	LYS	2.4
2	A	181	ILE	2.4
2	A	402	GLY	2.3
3	B	68	VAL	2.3
3	B	93	TRP	2.3
2	A	141	PHE	2.3
3	B	42	TYR	2.3
2	A	186	ILE	2.3
2	A	48[A]	SER	2.3
3	B	133	ARG	2.3
3	B	55	GLY	2.3
3	B	130	ASN	2.3
3	B	97	LEU	2.3
2	A	324	LEU	2.2
2	A	51[A]	PRO	2.2
3	B	168	ASN	2.2
3	B	99	ARG	2.2
3	B	167	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	85	LEU	2.2
3	B	47	TYR	2.1
3	B	28	PHE	2.1
3	B	212	THR	2.1
3	B	95	LEU	2.1
2	A	168[A]	ASP	2.1
3	B	117	ILE	2.0
3	B	91	ASN	2.0
3	B	106	PHE	2.0
3	B	10	GLN	2.0
3	B	38	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	502	4/4	0.70	0.20	49,50,52,54	0
5	EDO	A	503	4/4	0.79	0.21	59,60,60,61	0
5	EDO	B	386	4/4	0.80	0.16	63,63,64,64	0
5	EDO	A	501	4/4	0.86	0.16	52,52,53,53	0
4	CL	A	499	1/1	0.95	0.11	40,40,40,40	0
4	CL	A	497	1/1	0.96	0.10	36,36,36,36	0
5	EDO	A	500	4/4	0.96	0.08	32,33,34,35	0
4	CL	A	496	1/1	0.98	0.07	36,36,36,36	0
4	CL	A	498	1/1	0.98	0.09	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.