



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:29 AM UTC

PDB ID : 5I0I / pdb_00005i0i
Title : Crystal structure of myosin X motor domain with 2IQ motifs in pre-powerstroke state
Authors : Isabet, T.; Sweeney, H.L.; Houdusse, A.
Deposited on : 2016-02-04
Resolution : 3.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2361 (3.20-3.12)
Clashscore	190562	2486 (3.20-3.12)
Ramachandran outliers	187476	2405 (3.20-3.12)
Sidechain outliers	187428	2404 (3.20-3.12)
RSRZ outliers	180081	2361 (3.20-3.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	791	2% 86% 10% ...
1	B	791	4% 87% 10% ..
2	C	145	8% 68% 23% 8% .
3	E	145	2% 63% 28% 6% .
4	G	43	2% 60% 26% 14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	I	64	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	VO4	A	903	-	-	X	-
8	VO4	B	904	-	-	X	-

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 15350 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	784	Total	C	N	O	S	0	0	0
			6203	3935	1071	1169	28			
1	B	784	Total	C	N	O	S	0	1	0
			6159	3907	1060	1164	28			

- Molecule 2 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	145	Total	C	N	O	S	0	0	0
			1086	666	170	242	8			

- Molecule 3 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	145	Total	C	N	O	S	0	0	0
			1024	625	162	228	9			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	19	ALA	PHE	conflict	UNP P62158


- Molecule 4 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	G	43	Total	C	N	O	S	0	0	0
			306	188	45	71	2			

- Molecule 5 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	59	Total	C	N	O	S	0	0	0
			424	254	68	99	3			

- Molecule 6 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (CCD ID: MPO) (formula: $C_7H_{15}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
6	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Mg	0	0
			1	1		
7	B	1	Total	Mg	0	0
			1	1		

- Molecule 8 is VANADATE ION (CCD ID: VO4) (formula: O_4V).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total 5	O 4	V 1	0	0
8	B	1	Total 5	O 4	V 1	0	0

- Molecule 9 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
9	B	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 10 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	O	S	0	0
			5	4	1		
10	A	1	Total	O	S	0	0
			5	4	1		
10	A	1	Total	O	S	0	0
			5	4	1		
10	A	1	Total	O	S	0	0
			5	4	1		
10	A	1	Total	O	S	0	0
			5	4	1		
10	B	1	Total	O	S	0	0
			5	4	1		
10	C	1	Total	O	S	0	0
			5	4	1		
10	E	1	Total	O	S	0	0
			5	4	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	11	Total	O	0	0
			11	11		
11	B	5	Total	O	0	0
			5	5		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Unconventional myosin-X




• Molecule 1: Unconventional myosin-X




• Molecule 2: Calmodulin




• Molecule 3: Calmodulin



• Molecule 4: Calmodulin



• Molecule 5: Calmodulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	113.95Å 173.41Å 178.75Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.53 – 3.15 49.53 – 3.15	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.53-3.15) 99.9 (49.53-3.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 3.12Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.192 , 0.218 0.205 , 0.232	Depositor DCC
R_{free} test set	3082 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	103.7	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 100.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.006 for -h,l,k	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	15350	wwPDB-VP
Average B, all atoms (Å ²)	126.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPO, MG, VO4, SO4, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.90	4/6330 (0.1%)	1.34	48/8571 (0.6%)
1	B	0.82	1/6286 (0.0%)	1.34	32/8519 (0.4%)
2	C	0.89	0/1098	1.68	18/1486 (1.2%)
3	E	1.00	3/1032 (0.3%)	1.72	26/1401 (1.9%)
4	G	1.09	0/307	1.77	8/417 (1.9%)
5	I	0.98	0/425	1.62	2/576 (0.3%)
All	All	0.88	8/15478 (0.1%)	1.42	134/20970 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	22	ASP	CA-C	9.12	1.65	1.52
1	A	204	MET	SD-CE	7.73	1.98	1.79
1	B	747	MET	SD-CE	6.07	1.94	1.79
3	E	20	ASP	CA-C	5.89	1.60	1.52
1	A	662	MET	SD-CE	5.54	1.93	1.79

The worst 5 of 134 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	GLN	CA-C-N	12.63	144.44	121.70
1	A	179	GLN	C-N-CA	12.63	144.44	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	78	ASP	CA-C-N	9.67	139.10	121.70
2	C	78	ASP	C-N-CA	9.67	139.10	121.70
1	A	52	GLN	CA-C-N	9.30	138.44	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	52	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6203	0	5954	53	0
1	B	6159	0	5853	35	0
2	C	1086	0	955	19	0
3	E	1024	0	869	18	0
4	G	306	0	257	5	0
5	I	424	0	347	8	0
6	A	13	0	14	0	0
6	B	13	0	14	0	0
7	A	1	0	0	0	0
7	B	1	0	0	0	0
8	A	5	0	0	3	0
8	B	5	0	0	3	0
9	A	27	0	12	1	0
9	B	27	0	12	0	0
10	A	25	0	0	0	0
10	B	5	0	0	0	0
10	C	5	0	0	0	0
10	E	5	0	0	0	0
11	A	11	0	0	0	0
11	B	5	0	0	0	0
All	All	15350	0	14287	138	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 138 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:607:ARG:HH11	1:A:607:ARG:HG3	1.11	1.15
3:E:80:ASP:HA	3:E:81:SER:N	1.74	1.02
3:E:22:ASP:HB2	3:E:23:GLY:HA3	1.46	0.95
1:B:191:SER:HB3	1:B:195:ARG:HB2	1.52	0.89
1:A:65:VAL:HG11	1:A:70:SER:HB2	1.53	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	782/791 (99%)	738 (94%)	29 (4%)	15 (2%)	6	28
1	B	783/791 (99%)	735 (94%)	37 (5%)	11 (1%)	9	34
2	C	143/145 (99%)	124 (87%)	13 (9%)	6 (4%)	2	13
3	E	141/145 (97%)	122 (86%)	13 (9%)	6 (4%)	2	13
4	G	41/43 (95%)	32 (78%)	4 (10%)	5 (12%)	0	1
5	I	55/64 (86%)	50 (91%)	4 (7%)	1 (2%)	6	29
All	All	1945/1979 (98%)	1801 (93%)	100 (5%)	44 (2%)	5	24

5 of 44 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	THR
1	A	63	GLU
1	A	180	GLN
1	A	192	CYS
1	A	295	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	657/704 (93%)	634 (96%)	23 (4%)	32	59
1	B	644/704 (92%)	627 (97%)	17 (3%)	40	64
2	C	111/125 (89%)	96 (86%)	15 (14%)	4	16
3	E	97/124 (78%)	78 (80%)	19 (20%)	1	7
4	G	29/37 (78%)	22 (76%)	7 (24%)	1	3
5	I	40/55 (73%)	31 (78%)	9 (22%)	1	4
All	All	1578/1749 (90%)	1488 (94%)	90 (6%)	18	47

5 of 90 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	E	26	THR
3	E	133	ASP
3	E	42	ASN
3	E	91	VAL
4	G	87	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	248	ASN
4	G	97	ASN
1	B	507	ASN
5	I	137	ASN
2	C	41	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
8	VO4	B	904	9,7	0,4,4	-	-	-		
9	ADP	B	903	7,8	28,29,29	0.55	1 (3%)	43,45,45	0.76	2 (4%)
10	SO4	B	905	-	4,4,4	0.32	0	6,6,6	0.26	0
10	SO4	A	907	-	4,4,4	0.36	0	6,6,6	0.24	0
6	MPO	A	901	-	13,13,13	1.19	1 (7%)	17,17,17	0.38	0
6	MPO	B	901	-	13,13,13	1.04	1 (7%)	17,17,17	0.69	0
9	ADP	A	904	7,8	28,29,29	0.77	1 (3%)	43,45,45	0.86	4 (9%)
10	SO4	A	906	-	4,4,4	0.31	0	6,6,6	0.35	0
10	SO4	A	908	-	4,4,4	0.30	0	6,6,6	0.15	0
10	SO4	A	909	-	4,4,4	0.39	0	6,6,6	0.19	0
10	SO4	C	201	-	4,4,4	0.25	0	6,6,6	0.24	0
8	VO4	A	903	9,7	0,4,4	-	-	-		
10	SO4	A	905	-	4,4,4	0.50	0	6,6,6	0.42	0
10	SO4	E	201	-	4,4,4	0.29	0	6,6,6	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MPO	B	901	-	-	1/7/15/15	0/1/1/1
9	ADP	A	904	7,8	-	0/16/32/32	0/3/3/3
9	ADP	B	903	7,8	-	3/16/32/32	0/3/3/3
6	MPO	A	901	-	-	4/7/15/15	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	901	MPO	C1-S1	-4.26	1.71	1.77
6	B	901	MPO	C1-S1	-3.57	1.72	1.77
9	A	904	ADP	PB-O3B	-3.17	1.43	1.54
9	B	903	ADP	PB-O3B	-2.42	1.45	1.54

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	904	ADP	O3B-PB-O3A	-2.78	95.32	104.64
9	A	904	ADP	O2B-PB-O3A	2.61	113.40	104.64
9	A	904	ADP	O3'-C3'-C2'	2.50	119.83	111.82
9	A	904	ADP	O3'-C3'-C4'	2.41	118.00	111.08
9	B	903	ADP	O2B-PB-O3A	2.37	112.60	104.64

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	B	903	ADP	PA-O3A-PB-O2B
6	A	901	MPO	C2-C3-N1-C4
6	A	901	MPO	C2-C3-N1-C7
6	A	901	MPO	C1-C2-C3-N1
6	B	901	MPO	S1-C1-C2-C3


There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	B	904	VO4	3	0
9	A	904	ADP	1	0
8	A	903	VO4	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	E	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	E	80:ASP	C	81:SER	N	3.33

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	784/791 (99%)	-0.20	16 (2%) 65 44	41, 88, 196, 246	3 (0%)
1	B	784/791 (99%)	0.06	33 (4%) 40 23	48, 121, 232, 258	7 (0%)
2	C	145/145 (100%)	0.31	12 (8%) 17 10	109, 153, 217, 228	0
3	E	145/145 (100%)	0.13	3 (2%) 63 42	88, 144, 211, 234	0
4	G	43/43 (100%)	0.29	1 (2%) 61 40	191, 215, 230, 238	0
5	I	59/64 (92%)	0.18	3 (5%) 33 19	165, 214, 248, 253	0
All	All	1960/1979 (99%)	-0.01	68 (3%) 47 28	41, 116, 225, 258	10 (0%)

The worst 5 of 68 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	295	VAL	6.3
1	B	87	TYR	6.2
1	B	662	MET	5.9
1	B	602	CYS	5.9
1	B	400	SER	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	SO4	A	909	5/5	0.58	0.10	191,192,193,193	0
6	MPO	B	901	13/13	0.59	0.18	163,167,188,190	0
10	SO4	A	908	5/5	0.68	0.09	217,217,217,218	0
6	MPO	A	901	13/13	0.79	0.17	132,139,176,178	0
10	SO4	A	906	5/5	0.85	0.19	144,148,149,150	0
10	SO4	E	201	5/5	0.85	0.08	171,171,172,173	0
10	SO4	A	907	5/5	0.89	0.12	168,168,171,172	0
10	SO4	B	905	5/5	0.91	0.09	159,160,161,161	0
10	SO4	C	201	5/5	0.94	0.05	181,182,182,182	0
10	SO4	A	905	5/5	0.95	0.13	121,122,123,124	0
9	ADP	B	903	27/27	0.97	0.07	103,119,126,129	0
9	ADP	A	904	27/27	0.98	0.05	69,78,85,87	0
8	VO4	A	903	5/5	0.98	0.05	72,75,77,77	0
8	VO4	B	904	5/5	0.99	0.04	109,110,111,111	0
7	MG	A	902	1/1	1.00	0.01	69,69,69,69	0
7	MG	B	902	1/1	1.00	0.05	101,101,101,101	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ADP B 903:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ADP A 904:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.