



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:21 PM UTC

PDB ID : 4I1F / pdb_00004i1f
Title : Structure of Parkin-S223P E3 ligase
Authors : Loughheed, J.C.; Brecht, E.; Yao, N.H.
Deposited on : 2012-11-20
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ZN	A	502	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2679 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called E3 ubiquitin-protein ligase parkin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	306	Total	C	N	O	S	0	1	0
			2403	1493	436	435	39			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	223	PRO	SER	engineered mutation	UNP O60260

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	8	Total	Zn	0	0
			8	8		

- Molecule 3 is BARIUM ION (CCD ID: BA) (formula: Ba).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ba	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	267	Total	O	0	0
			267	267		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: E3 ubiquitin-protein ligase parkin



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	86.95Å 133.16Å 65.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.81 – 1.58 72.81 – 1.58	Depositor EDS
% Data completeness (in resolution range)	99.5 (72.81-1.58) 99.5 (72.81-1.58)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.78 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.205 , 0.245 0.201 , 0.238	Depositor DCC
R_{free} test set	2660 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2679	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, BA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.67	32/2468 (1.3%)	1.46	9/3341 (0.3%)

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	201	CYS	C-O	16.75	1.43	1.24
1	A	433	HIS	ND1-CE1	8.80	1.41	1.32
1	A	303	HIS	ND1-CE1	7.71	1.40	1.32
1	A	453	TRP	CD1-NE1	7.41	1.53	1.37
1	A	201	CYS	CA-C	7.07	1.60	1.52
1	A	248	VAL	N-CA	6.46	1.53	1.46
1	A	303	HIS	CG-CD2	6.30	1.42	1.35
1	A	211	LYS	N-CA	6.24	1.53	1.45
1	A	227	HIS	N-CA	6.13	1.54	1.46
1	A	205	SER	N-CA	6.03	1.53	1.46
1	A	463	PHE	C-O	5.87	1.31	1.23
1	A	257	HIS	ND1-CE1	5.87	1.38	1.32
1	A	151	LYS	CA-C	5.83	1.60	1.52
1	A	241	CYS	N-CA	5.80	1.53	1.46
1	A	238	CYS	CA-C	5.75	1.60	1.52
1	A	255	SER	CA-C	5.62	1.60	1.52
1	A	446	CYS	N-CA	5.57	1.53	1.46
1	A	433	HIS	CD2-NE2	5.56	1.44	1.37
1	A	212	CYS	CA-CB	5.56	1.62	1.53
1	A	410	THR	N-CA	5.55	1.52	1.45
1	A	264	PHE	N-CA	-5.54	1.39	1.46
1	A	252	GLN	N-CA	5.53	1.52	1.46
1	A	302	HIS	C-O	-5.50	1.17	1.24
1	A	407	SER	N-CA	5.46	1.52	1.45
1	A	266	LEU	N-CA	5.42	1.53	1.46
1	A	145	SER	CA-CB	5.22	1.62	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	401	ALA	CA-CB	5.21	1.59	1.52
1	A	297	LEU	N-CA	5.21	1.52	1.46
1	A	297	LEU	C-O	5.15	1.29	1.23
1	A	301	LEU	N-CA	5.15	1.53	1.46
1	A	273	ASN	N-CA	5.12	1.52	1.46
1	A	373	HIS	CG-CD2	5.04	1.41	1.35

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	CYS	CA-C-N	14.04	135.50	120.47
1	A	201	CYS	C-N-CA	14.04	135.50	120.47
1	A	312	TYR	N-CA-C	-5.63	105.22	111.36
1	A	344	GLU	CA-C-N	5.54	126.77	119.84
1	A	344	GLU	C-N-CA	5.54	126.77	119.84
1	A	201	CYS	N-CA-C	5.52	118.36	109.15
1	A	188	ILE	CA-C-N	-5.15	114.61	119.76
1	A	188	ILE	C-N-CA	-5.15	114.61	119.76
1	A	276	GLN	CB-CA-C	-5.04	103.31	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2403	0	2272	22	0
2	A	8	0	0	2	0
3	A	1	0	0	1	0
4	A	267	0	0	3	0
All	All	2679	0	2272	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:201:CYS:O	3:A:509:BA:BA	1.61	1.06
1:A:196:CYS:HG	2:A:502:ZN:ZN	0.93	0.82
1:A:415:THR:OG1	4:A:776:HOH:O	2.08	0.72
1:A:196:CYS:SG	1:A:201:CYS:HB3	2.31	0.70
1:A:432:MET:CE	1:A:448:ASN:HB2	2.23	0.69
1:A:420:ARG:HD2	1:A:441:CYS:SG	2.34	0.68
1:A:196:CYS:SG	2:A:502:ZN:ZN	1.82	0.67
1:A:265:HIS:HD2	4:A:629:HOH:O	1.78	0.67
1:A:440:GLN:CD	1:A:440:GLN:H	2.03	0.66
1:A:432:MET:HE1	1:A:448:ASN:HB2	1.79	0.64
1:A:302:HIS:CD2	1:A:305:ARG:HE	2.18	0.62
1:A:201:CYS:SG	1:A:202:PRO:HD2	2.47	0.55
1:A:349:LYS:NZ	4:A:861:HOH:O	2.33	0.55
1:A:343:PRO:HG2	1:A:366:ARG:NH1	2.30	0.47
1:A:196:CYS:SG	1:A:201:CYS:CB	3.04	0.45
1:A:432:MET:HE2	1:A:448:ASN:HB2	1.95	0.45
1:A:442:ARG:HA	1:A:442:ARG:HD2	1.79	0.45
1:A:458:MET:HE3	1:A:462:TRP:CH2	2.51	0.45
1:A:300:GLU:OE2	1:A:302:HIS:HB2	2.18	0.44
1:A:332:CYS:O	1:A:338:GLY:HA2	2.20	0.42
1:A:280:ASP:HB3	1:A:283:LEU:HB2	2.02	0.42
1:A:302:HIS:CD2	1:A:305:ARG:NE	2.88	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	299/325 (92%)	286 (96%)	12 (4%)	1 (0%)	36	20

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	151	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	268/281 (95%)	261 (97%)	7 (3%)	40 11

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	170	ARG
1	A	201	CYS
1	A	217	THR
1	A	271	ARG
1	A	306	ILE
1	A	419	PRO
1	A	440	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	265	HIS
1	A	279	HIS
1	A	302	HIS
1	A	313	ASN
1	A	316	GLN
1	A	438	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	306/325 (94%)	0.89	42 (13%) 6 11	13, 24, 47, 61	16 (5%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	217	THR	10.6
1	A	155	GLN	9.8
1	A	392	ARG	9.6
1	A	216	PRO	8.2
1	A	222	THR	8.1
1	A	200	HIS	7.4
1	A	282	GLN	7.4
1	A	376	GLU	7.1
1	A	347	GLN	7.0
1	A	367	GLU	6.5
1	A	299	LYS	5.8
1	A	390	ALA	4.5
1	A	201	CYS	4.3
1	A	214	ALA	3.8
1	A	354	GLY	3.7
1	A	346	ASP	3.5
1	A	198	SER	3.4
1	A	154	CYS	3.3
1	A	391	TYR	3.3
1	A	202	PRO	3.3
1	A	199	PRO	3.1
1	A	149	TYR	3.1
1	A	152	GLY	3.0
1	A	170	ARG	3.0
1	A	411	ILE	2.9
1	A	378	SER	2.9
1	A	150	CYS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	195	GLU	2.7
1	A	327	MET	2.6
1	A	213	GLY	2.5
1	A	224	VAL	2.5
1	A	211	LYS	2.4
1	A	153	PRO	2.4
1	A	375	GLY	2.4
1	A	203	GLY	2.4
1	A	328	GLY	2.4
1	A	309	GLU	2.4
1	A	281	PRO	2.3
1	A	393	VAL	2.1
1	A	283	LEU	2.1
1	A	306	ILE	2.1
1	A	317	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BA	A	509	1/1	0.92	0.09	38,38,38,38	1
2	ZN	A	501	1/1	0.94	0.06	23,23,23,23	1
2	ZN	A	502	1/1	0.95	0.08	33,33,33,33	0
2	ZN	A	507	1/1	0.96	0.05	28,28,28,28	0
2	ZN	A	505	1/1	0.98	0.06	26,26,26,26	0
2	ZN	A	506	1/1	0.99	0.07	38,38,38,38	0
2	ZN	A	503	1/1	1.00	0.02	17,17,17,17	0
2	ZN	A	508	1/1	1.00	0.02	15,15,15,15	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	504	1/1	1.00	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.