



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:53 PM UTC

PDB ID : 4I1Q / pdb_00004i1q
Title : Crystal Structure of hBRAP1 N-BAR domain
Authors : Sanchez-Barrena, M.J.
Deposited on : 2012-11-21
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	4% 64% 22% • • 10%
1	B	221	6% 56% 28% 5% 10%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bridging integrator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	200	Total	C	N	O	S	0	0	0
			1674	1054	288	327	5			
1	B	198	Total	C	N	O	S	0	0	0
			1657	1044	285	323	5			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	GLY	-	expression tag	UNP Q9UBW5
A	19	SER	-	expression tag	UNP Q9UBW5
A	48	ASN	SER	SEE REMARK 999	UNP Q9UBW5
B	18	GLY	-	expression tag	UNP Q9UBW5
B	19	SER	-	expression tag	UNP Q9UBW5
B	48	ASN	SER	SEE REMARK 999	UNP Q9UBW5


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	38	Total	O	0	0
			38	38		
2	B	24	Total	O	0	0
			24	24		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bridging integrator 2



• Molecule 1: Bridging integrator 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.15Å 81.28Å 81.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.41 – 2.53 57.41 – 2.54	Depositor EDS
% Data completeness (in resolution range)	87.6 (57.41-2.53) 94.1 (57.41-2.54)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 2.58Å)	Xtriage
Refinement program	REFMAC 5.7.0024	Depositor
R, R_{free}	0.196 , 0.252 0.195 , 0.251	Depositor DCC
R_{free} test set	802 reflections (4.65%)	wwPDB-VP
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 32.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.018 for -h,l,k 0.079 for -k,-h,l 0.036 for l,-k,h 0.016 for l,h,k 0.016 for k,l,h	Xtriage
Reported twinning fraction	0.888 for H, K, L 0.112 for K, H, -L	Depositor
Outliers	0 of 15848 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3393	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/1704	1.21	6/2288 (0.3%)
1	B	0.96	0/1687	1.21	5/2266 (0.2%)
All	All	0.95	0/3391	1.21	11/4554 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	161	ASN	N-CA-C	9.75	124.23	110.68
1	A	40	ASP	N-CA-C	-7.20	102.92	112.72
1	B	103	VAL	N-CA-C	-6.95	103.70	110.72
1	A	160	GLN	N-CA-C	6.15	119.98	112.23
1	B	135	ILE	N-CA-C	6.11	116.77	110.36
1	A	42	ARG	N-CA-C	-5.81	105.03	111.36
1	A	105	ASN	N-CA-C	5.41	117.18	111.28
1	B	194	GLU	N-CA-C	5.29	116.85	111.14
1	B	209	ILE	N-CA-C	5.26	115.88	110.36
1	B	132	PHE	N-CA-C	5.07	117.70	111.82
1	A	171	ALA	N-CA-C	-5.04	106.39	112.54

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	160	GLN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1674	0	1627	41	0
1	B	1657	0	1610	56	0
2	A	38	0	0	2	0
2	B	24	0	0	0	0
All	All	3393	0	3237	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:ALA:O	1:A:175:GLU:HG2	1.74	0.86
1:A:142:ARG:HB2	1:A:187:LEU:HD22	1.64	0.78
1:B:156:LEU:HD21	1:B:174:GLU:HA	1.68	0.75
1:B:156:LEU:HD11	1:B:173:ALA:O	1.88	0.74
1:A:217:ARG:NH1	1:B:228:ASN:OD1	2.22	0.73
1:A:125:MET:HG2	1:A:209:ILE:HD11	1.71	0.73
1:A:75:HIS:ND1	1:A:114:GLU:OE2	2.23	0.71
1:A:118:ALA:O	1:A:123:ARG:HG3	1.91	0.71
1:A:156:LEU:HB2	1:A:177:PHE:HD2	1.54	0.71
1:B:146:LEU:HB2	1:B:184:PHE:CE1	2.27	0.69
1:A:56:GLU:HG2	1:B:88:ILE:HG22	1.75	0.69
1:B:46:SER:O	1:B:49:ASN:HB2	1.92	0.69
1:B:94:ASP:OD2	1:B:237:LYS:NZ	2.27	0.67
1:A:156:LEU:HB2	1:A:177:PHE:CD2	2.31	0.66
1:B:43:PHE:HE1	1:B:192:LEU:HD13	1.61	0.65
1:A:40:ASP:O	1:A:44:GLU:HG2	1.98	0.63
1:A:149:TYR:HE1	1:A:181:GLN:HB3	1.62	0.62
1:A:125:MET:HG2	1:A:209:ILE:CD1	2.28	0.62
1:B:237:LYS:NZ	1:B:237:LYS:HB3	2.15	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:39:LYS:N	1:A:41:GLU:HG3	2.17	0.60
1:B:85:LEU:HD11	1:B:235:MET:HE1	1.85	0.59
1:A:52:GLN:NE2	1:A:199:TYR:OH	2.35	0.59
1:A:67:PHE:HA	1:B:77:SER:OG	2.04	0.57
1:B:45:GLN:HA	1:B:48:ASN:HB3	1.86	0.56
1:B:181:GLN:O	1:B:185:GLU:HG2	2.07	0.55
1:B:153:ARG:HG2	1:B:177:PHE:HE2	1.71	0.55
1:B:43:PHE:O	1:B:46:SER:N	2.40	0.54
1:A:181:GLN:HG3	1:A:182:THR:N	2.22	0.54
1:B:50:PHE:CZ	1:B:135:ILE:HG22	2.43	0.54
1:B:187:LEU:O	1:B:191:LEU:HB2	2.08	0.54
1:B:156:LEU:HG	1:B:173:ALA:HB1	1.90	0.53
1:B:136:LYS:O	1:B:139:ILE:HG13	2.08	0.53
2:A:322:HOH:O	1:B:228:ASN:HB3	2.08	0.52
1:A:145:LYS:HD2	1:A:183:VAL:CG1	2.40	0.52
1:A:149:TYR:CE1	1:A:181:GLN:CB	2.93	0.51
1:A:152:ALA:HB1	1:A:177:PHE:CD1	2.45	0.51
1:A:149:TYR:HE1	1:A:181:GLN:CB	2.24	0.51
1:B:142:ARG:NH2	1:B:188:ASN:OD1	2.44	0.50
1:B:186:ASP:O	1:B:190:GLU:HB2	2.11	0.50
1:A:149:TYR:CE1	1:A:181:GLN:HB3	2.43	0.50
1:B:43:PHE:CE1	1:B:192:LEU:HD13	2.43	0.50
1:A:141:LYS:HG2	1:A:144:ARG:NH2	2.26	0.50
1:B:237:LYS:NZ	1:B:237:LYS:CB	2.74	0.50
1:A:146:LEU:HB2	1:A:184:PHE:CE1	2.47	0.50
1:A:64:LEU:HD12	1:A:125:MET:HG3	1.93	0.49
1:B:45:GLN:O	1:B:48:ASN:HB3	2.12	0.49
1:B:50:PHE:CE1	1:B:135:ILE:HG21	2.48	0.48
1:A:82:SER:HB2	1:A:103:VAL:HG22	1.94	0.48
1:A:163:LYS:HD2	1:A:165:LYS:HG2	1.96	0.48
1:B:79:LYS:HD2	1:B:107:ASP:OD1	2.14	0.47
1:A:125:MET:O	1:A:129:VAL:HG22	2.15	0.47
1:A:110:TRP:NE1	1:A:224:MET:HE1	2.30	0.47
1:A:141:LYS:HG2	1:A:144:ARG:HH21	1.80	0.47
1:A:141:LYS:CG	1:A:144:ARG:NH2	2.77	0.47
1:B:62:LYS:HA	1:B:62:LYS:HD2	1.81	0.47
1:A:44:GLU:OE1	1:A:44:GLU:HA	2.14	0.46
1:B:237:LYS:HB3	1:B:237:LYS:HZ2	1.79	0.46
1:A:109:LEU:O	1:A:110:TRP:C	2.56	0.46
1:B:144:ARG:O	1:B:147:VAL:N	2.48	0.46
1:B:152:ALA:O	1:B:156:LEU:HD13	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:146:LEU:HB2	1:B:184:PHE:CD1	2.50	0.46
1:B:75:HIS:ND1	1:B:114:GLU:OE2	2.44	0.46
1:B:50:PHE:CZ	1:B:135:ILE:CG2	2.99	0.46
1:A:214:SER:O	1:B:228:ASN:ND2	2.45	0.46
1:B:54:GLN:HE22	1:B:136:LYS:HD2	1.81	0.45
1:B:212:ASN:O	1:B:216:LEU:HG	2.16	0.45
1:B:54:GLN:HA	1:B:132:PHE:CE2	2.52	0.44
1:A:156:LEU:HG	1:A:160:GLN:OE1	2.16	0.44
1:A:51:TYR:O	1:A:54:GLN:HG3	2.18	0.44
1:B:153:ARG:HG2	1:B:177:PHE:CE2	2.51	0.44
1:B:109:LEU:CB	1:B:224:MET:HE2	2.48	0.44
1:A:152:ALA:HB1	1:A:177:PHE:CG	2.52	0.43
1:B:188:ASN:O	1:B:192:LEU:HB2	2.19	0.43
1:B:185:GLU:O	1:B:189:GLN:CB	2.67	0.43
1:A:141:LYS:HD2	1:A:187:LEU:HD11	2.00	0.43
1:B:203:ILE:O	1:B:207:VAL:HG23	2.18	0.43
1:B:156:LEU:HD11	1:B:173:ALA:C	2.43	0.43
1:B:142:ARG:NE	1:B:188:ASN:OD1	2.51	0.42
1:A:126:GLU:OE2	2:A:338:HOH:O	2.21	0.42
1:B:193:GLU:O	1:B:197:ILE:HG13	2.20	0.42
1:B:82:SER:HB2	1:B:103:VAL:HG22	2.01	0.42
1:B:193:GLU:O	1:B:196:PRO:HD2	2.20	0.42
1:B:234:VAL:HG12	1:B:235:MET:HE2	2.01	0.42
1:B:75:HIS:CD2	1:B:75:HIS:C	2.97	0.41
1:B:234:VAL:HG12	1:B:235:MET:N	2.33	0.41
1:B:56:GLU:OE2	1:B:202:ARG:HD2	2.19	0.41
1:A:39:LYS:C	1:A:41:GLU:H	2.28	0.41
1:B:134:GLU:OE2	1:B:134:GLU:C	2.63	0.41
1:A:232:TYR:CD1	1:A:232:TYR:C	2.99	0.41
1:A:56:GLU:CG	1:B:88:ILE:HG22	2.48	0.40
1:B:85:LEU:HD11	1:B:235:MET:CE	2.49	0.40
1:B:64:LEU:HD23	1:B:64:LEU:HA	1.89	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/221 (90%)	184 (93%)	14 (7%)	0	100	100
1	B	196/221 (89%)	181 (92%)	13 (7%)	2 (1%)	12	23
All	All	394/442 (89%)	365 (93%)	27 (7%)	2 (0%)	24	41

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	162	ALA
1	B	236	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/198 (91%)	163 (90%)	18 (10%)	7	15
1	B	179/198 (90%)	156 (87%)	23 (13%)	4	8
All	All	360/396 (91%)	319 (89%)	41 (11%)	5	11

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LYS
1	A	40	ASP
1	A	41	GLU
1	A	52	GLN
1	A	83	GLU
1	A	147	VAL
1	A	155	HIS
1	A	157	GLU
1	A	163	LYS
1	A	166	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	169	LYS
1	A	179	LYS
1	A	183	VAL
1	A	209	ILE
1	A	217	ARG
1	A	225	SER
1	A	234	VAL
1	A	238	LEU
1	B	43	PHE
1	B	69	SER
1	B	81	VAL
1	B	82	SER
1	B	91	SER
1	B	107	ASP
1	B	111	GLU
1	B	134	GLU
1	B	135	ILE
1	B	139	ILE
1	B	146	LEU
1	B	147	VAL
1	B	151	SER
1	B	153	ARG
1	B	159	VAL
1	B	161	ASN
1	B	163	LYS
1	B	175	GLU
1	B	183	VAL
1	B	205	CYS
1	B	219	VAL
1	B	234	VAL
1	B	237	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	49	ASN
1	A	52	GLN
1	A	86	GLN
1	A	120	GLN
1	A	131	GLN
1	A	161	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	49	ASN
1	B	52	GLN
1	B	54	GLN
1	B	155	HIS
1	B	160	GLN
1	B	229	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	200/221 (90%)	0.23	8 (4%) 42 39	30, 47, 125, 156	0
1	B	198/221 (89%)	0.47	14 (7%) 22 19	31, 50, 119, 154	0
All	All	398/442 (90%)	0.35	22 (5%) 30 27	30, 49, 124, 156	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	168	ALA	3.9
1	B	183	VAL	3.7
1	B	102	ILE	3.6
1	A	39	LYS	3.4
1	B	192	LEU	3.3
1	A	162	ALA	3.2
1	A	159	VAL	3.1
1	B	170	THR	3.0
1	B	194	GLU	2.9
1	B	178	ASN	2.7
1	B	154	HIS	2.6
1	A	156	LEU	2.5
1	B	50	PHE	2.5
1	A	177	PHE	2.4
1	B	42	ARG	2.4
1	A	174	GLU	2.4
1	B	173	ALA	2.3
1	A	153	ARG	2.1
1	B	43	PHE	2.1
1	B	158	ALA	2.1
1	A	150	ASP	2.1
1	B	180	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.