



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:43 PM UTC

PDB ID : 6I2S / pdb_00006i2s
Title : Crystal structure of the SucA domain of Mycobacterium smegmatis KGD (R802A) in complex with GarA, following 2-oxoglutarate soak
Authors : Wagner, T.; Bellinzoni, M.; Alzari, P.M.
Deposited on : 2018-11-01
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	868	4% 88% 7% .
2	B	115	7% 69% 16% 16%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7309 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multifunctional 2-oxoglutarate metabolism enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	830	Total	C	N	O	S	0	1	0
			6371	4024	1107	1216	24			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	360	GLY	-	expression tag	UNP A0R2B1
A	802	ALA	ARG	engineered mutation	UNP A0R2B1

- Molecule 2 is a protein called Glycogen accumulation regulator GarA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	97	Total	C	N	O	0	1	0
			739	463	134	142			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	44	GLY	-	expression tag	UNP A0QYG2

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		

- Molecule 5 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 6 is (4S)-4-{3-[(4-amino-2-methylpyrimidin-5-yl)methyl]-5-(2-{[(S)-hydroxy(phosphonooxy)phosphoryl]oxy}ethyl)-4-methyl-1,3lambda 5 -thiazol-2-yl}-4-hydroxybutanoic acid (CCD ID: TD6) (formula: $C_{16}H_{25}N_4O_{10}P_2S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
6	A	1	Total	C	N	O	P	S	0	0
			33	16	4	10	2	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	151	Total 151	O 151	0	0
7	B	8	Total 8	O 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme



- Molecule 2: Glycogen accumulation regulator GarA



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	141.27Å 168.06Å 101.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.20 – 2.40 35.20 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.0 (35.20-2.40) 97.9 (35.20-2.40)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 2.40Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.192 , 0.219 0.197 , 0.224	Depositor DCC
R_{free} test set	2342 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	55.6	Xtriage
Anisotropy	0.395	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 60.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7309	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG, CA, TD6

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.87	1/6502 (0.0%)	1.31	13/8830 (0.1%)
2	B	0.78	0/755	1.07	1/1023 (0.1%)
All	All	0.86	1/7257 (0.0%)	1.29	14/9853 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	589	MET	SD-CE	-6.67	1.62	1.79

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	700	LYS	CA-C-N	6.14	128.79	120.38
1	A	700	LYS	C-N-CA	6.14	128.79	120.38
1	A	1058	ASP	CA-CB-CG	5.97	118.57	112.60
1	A	627	GLY	CA-C-N	5.65	128.12	120.38
1	A	627	GLY	C-N-CA	5.65	128.12	120.38
1	A	791	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	1025	ILE	CA-C-N	5.43	127.56	120.28
1	A	1025	ILE	C-N-CA	5.43	127.56	120.28
1	A	660	LEU	CA-C-N	5.21	127.52	120.38
1	A	660	LEU	C-N-CA	5.21	127.52	120.38
2	B	126	ASP	N-CA-C	-5.19	106.63	113.12
1	A	877	ASP	CA-CB-CG	5.14	117.74	112.60
1	A	368	ALA	CA-C-N	5.02	127.00	120.28
1	A	368	ALA	C-N-CA	5.02	127.00	120.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6371	0	6103	23	0
2	B	739	0	721	10	0
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	5	0	0	0	0
6	A	33	0	21	0	0
7	A	151	0	0	1	0
7	B	8	0	0	0	0
All	All	7309	0	6845	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:58:VAL:HG22	2:B:68:ARG:HG2	1.75	0.68
1:A:1044:THR:HG22	1:A:1047:ASN:H	1.57	0.68
2:B:61:ARG:HB3	2:B:142:ARG:HB2	1.77	0.66
1:A:504:LYS:HE2	1:A:745:ARG:HB3	1.81	0.62
1:A:642:LEU:HD22	1:A:674:ILE:HB	1.82	0.61
1:A:603:PRO:HD3	1:A:991:PHE:CZ	2.40	0.56
1:A:538:PRO:HB2	7:A:1506:HOH:O	2.09	0.51
1:A:632:ASP:OD1	1:A:634:ARG:HB2	2.10	0.51
2:B:92:THR:HG21	2:B:140:LYS:HD2	1.93	0.51
1:A:1171:PRO:HB3	1:A:1201:ARG:O	2.12	0.50
1:A:692:SER:HB2	1:A:697:ASP:OD2	2.12	0.50
1:A:981:VAL:HG22	1:A:988:ILE:HD11	1.94	0.49
1:A:589:MET:HE2	2:B:140:LYS:HG3	1.95	0.49
2:B:97:HIS:CD2	2:B:138:ILE:HG23	2.48	0.49
1:A:875:ARG:HA	1:A:1079:VAL:HG13	1.96	0.47
1:A:1198:ILE:HG22	1:A:1221:ILE:HG23	1.96	0.47
1:A:908:PHE:CZ	1:A:1070:LYS:HG2	2.50	0.46
1:A:1041:MET:HE2	1:A:1117:GLY:HA3	1.98	0.46
2:B:119:TYR:HB2	2:B:137:GLN:HB3	1.96	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1183:THR:HB	1:A:1187:ILE:HD12	1.98	0.46
1:A:1022:SER:HB3	1:A:1206:PRO:HB3	1.98	0.46
2:B:55:ALA:HB3	2:B:71:LEU:HB2	1.99	0.45
1:A:794:ARG:HG2	2:B:95[B]:ARG:HH21	1.81	0.45
1:A:628:GLU:HG2	1:A:664:ARG:O	2.17	0.44
2:B:75:THR:HG22	2:B:101:ARG:HG2	2.01	0.43
1:A:617:VAL:HG11	1:A:639:PRO:HG3	2.01	0.42
1:A:705:PRO:HG2	1:A:735:VAL:HG13	2.01	0.42
1:A:510:GLY:O	1:A:741:CYS:HB2	2.20	0.41
2:B:60:LYS:HG3	2:B:144:VAL:CG2	2.52	0.41
1:A:1177:TRP:CD1	1:A:1197:ARG:HD3	2.56	0.40
1:A:1174:GLN:OE1	1:A:1206:PRO:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	821/868 (95%)	798 (97%)	22 (3%)	1 (0%)	48	64
2	B	96/115 (84%)	94 (98%)	2 (2%)	0	100	100
All	All	917/983 (93%)	892 (97%)	24 (3%)	1 (0%)	48	64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	682	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	648/725 (89%)	633 (98%)	15 (2%)	44	66
2	B	79/93 (85%)	79 (100%)	0	100	100
All	All	727/818 (89%)	712 (98%)	15 (2%)	47	69

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	412	LEU
1	A	421	VAL
1	A	467	VAL
1	A	496	LEU
1	A	642	LEU
1	A	748	ASN
1	A	801	GLU
1	A	811	GLU
1	A	909	THR
1	A	915	ILE
1	A	950	LEU
1	A	1079	VAL
1	A	1112	LEU
1	A	1135	VAL
1	A	1137	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	394	ASN
1	A	406	ASN
1	A	633	ASN
1	A	679	GLN
1	A	757	GLN
1	A	933	ASN
2	B	73	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PO4	A	1303	-	4,4,4	1.97	2 (50%)	6,6,6	0.44	0
6	TD6	A	1304	3	31,34,34	0.45	0	38,50,50	0.62	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	TD6	A	1304	3	-	6/25/26/26	0/2/2/2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	1303	PO4	P-O3	2.01	1.60	1.54
5	A	1303	PO4	P-O4	2.01	1.60	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	1304	TD6	C4-C5-S1	2.45	113.11	110.56

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	1304	TD6	OL1-C11-C2-N3
6	A	1304	TD6	OL1-C11-C2-S1
6	A	1304	TD6	C7-O7-PA-O1A
6	A	1304	TD6	PA-O3A-PB-O2B
6	A	1304	TD6	CLB-C13-CLC-OL2
6	A	1304	TD6	CLB-C13-CLC-OL3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	830/868 (95%)	0.29	39 (4%) 36 32	34, 67, 113, 150	1 (0%)
2	B	97/115 (84%)	0.53	8 (8%) 17 14	47, 79, 121, 137	1 (1%)
All	All	927/983 (94%)	0.32	47 (5%) 33 30	34, 70, 116, 150	2 (0%)

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	405	VAL	5.1
1	A	1227	GLY	4.5
2	B	54	SER	4.1
1	A	747	HIS	3.8
1	A	404	ASP	3.7
1	A	827	ILE	3.5
1	A	830	LYS	3.4
1	A	363	ILE	3.4
1	A	615	GLY	3.3
1	A	1193	THR	3.0
1	A	472	ASP	3.0
1	A	561	PHE	2.9
1	A	427	VAL	2.8
1	A	422	ASP	2.8
1	A	829	SER	2.8
2	B	147	THR	2.8
2	B	104	GLY	2.8
1	A	406	ASN	2.8
1	A	562	GLU	2.7
1	A	1080	SER	2.7
1	A	828	PRO	2.7
1	A	364	GLU	2.6
1	A	871	ALA	2.6
1	A	888	SER	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	811	GLU	2.5
1	A	560	GLU	2.5
1	A	413	TRP	2.5
1	A	835	VAL	2.5
1	A	1100	THR	2.4
2	B	102	LEU	2.4
2	B	95[A]	ARG	2.3
1	A	749	GLU	2.3
1	A	793	LEU	2.3
1	A	412	LEU	2.2
1	A	626	THR	2.2
1	A	876	ILE	2.2
1	A	407	SER	2.2
1	A	559	SER	2.2
1	A	575	ASP	2.2
2	B	107	PHE	2.1
1	A	1161	VAL	2.1
1	A	1103	GLU	2.1
1	A	1081	ASP	2.1
1	A	397	PHE	2.1
2	B	69	PHE	2.1
1	A	500	TYR	2.0
2	B	74	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PO4	A	1303	5/5	0.92	0.11	113,115,116,116	0
6	TD6	A	1304	33/33	0.98	0.07	39,52,83,87	0
4	CA	A	1302	1/1	0.99	0.03	64,64,64,64	0
3	MG	A	1301	1/1	1.00	0.05	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.