



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:35 PM UTC

PDB ID : 7I2X / pdb_00007i2x
Title : Group deposition for crystallographic fragment screening of the NS5 RNA-dependent RNA polymerase from Dengue virus serotype 2 – PanDDA analysis group deposition of ground-state model
Authors : Aschenbrenner, J.C.; Saini, M.; Chopra, A.; Marples, P.G.; Balcomb, B.H.; Lithgo, R.M.; Fearon, D.; von Delft, F.; Ruiz, F.X.; Arnold, E.
Deposited on : 2025-03-06
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	637	19% 80% 10% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PO4	A	1009	-	X	X	-

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 5346 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NS5 RNA-dependent RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	0	8	0
			4819	3035	864	886	34			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	GLY	-	expression tag	UNP Q91H74
A	265	PRO	-	expression tag	UNP Q91H74


- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	1
			24	12	2	8	2		

- Molecule 5 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		
6	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 7 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		
7	A	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	457	Total	O	0	0
			457	457		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NS5 RNA-dependent RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	82.48Å 116.86Å 148.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.39 – 1.50 74.39 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.4 (74.39-1.50) 98.5 (74.39-1.50)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.193 , 0.220 (Not available) , 0.208	Depositor DCC
R_{free} test set	5803 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	24.9	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5346	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PEG, DMS, MES, PO4, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.09	2/4929 (0.0%)	1.29	4/6645 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	768	HIS	CE1-NE2	5.90	1.38	1.32
1	A	510	GLU	CD-OE2	5.55	1.35	1.25

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	510	GLU	CB-CG-CD	6.25	123.23	112.60
1	A	844	ASP	CA-CB-CG	5.12	117.72	112.60
1	A	479	LEU	CA-C-N	5.05	125.55	119.94
1	A	479	LEU	C-N-CA	5.05	125.55	119.94

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4819	0	4721	51	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	2	0	0	0	0
4	A	24	0	26	1	0
5	A	12	0	18	4	0
6	A	21	0	30	0	0
7	A	10	0	0	4	0
8	A	457	0	0	15	0
All	All	5346	0	4795	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:664:ASP:OD1	7:A:1009:PO4:O4	1.93	0.86
1:A:664:ASP:OD1	7:A:1009:PO4:P	2.38	0.81
1:A:318:SER:O	8:A:1101:HOH:O	2.03	0.76
5:A:1005:DMS:H11	8:A:1280:HOH:O	1.87	0.75
1:A:510:GLU:O	8:A:1102:HOH:O	2.09	0.70
1:A:534:ASP:OD1	7:A:1009:PO4:O4	2.12	0.67
1:A:403:VAL:HG21	1:A:426:VAL:HG21	1.77	0.66
5:A:1005:DMS:H12	8:A:1324:HOH:O	1.97	0.65
1:A:705:GLN:OE1	8:A:1103:HOH:O	2.15	0.64
1:A:534:ASP:OD1	7:A:1009:PO4:P	2.59	0.61
1:A:447:CYS:SG	1:A:450:CYS:HB2	2.42	0.60
1:A:886:MET:HE1	8:A:1233:HOH:O	2.02	0.59
1:A:886:MET:CE	8:A:1233:HOH:O	2.51	0.58
1:A:764:LEU:HG	1:A:765:MET:CE	2.33	0.57
1:A:305:TYR:OH	1:A:308:SER:OG	2.17	0.56
1:A:834:GLU:HG3	8:A:1139:HOH:O	2.06	0.55
1:A:510:GLU:O	1:A:514:LYS:HG3	2.09	0.53
1:A:733:GLU:O	1:A:737:ARG:HG3	2.09	0.53
1:A:805:THR:CG2	1:A:809:MET:HE1	2.40	0.52
1:A:356:LYS:HD3	8:A:1520:HOH:O	2.09	0.52
1:A:800:THR:HG21	8:A:1102:HOH:O	2.09	0.52
1:A:570:LYS:HG2	8:A:1142:HOH:O	2.09	0.51
1:A:764:LEU:HG	1:A:765:MET:HE2	1.93	0.51
1:A:312:LYS:O	1:A:313:GLN:HG3	2.12	0.50
1:A:528:GLY:O	1:A:668:LYS:HE3	2.12	0.50
1:A:748:LEU:HD13	4:A:1004[B]:MES:H61	1.94	0.49
1:A:790:THR:HG22	8:A:1379:HOH:O	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:411:ALA:HA	1:A:477:MET:O	2.13	0.48
1:A:834:GLU:OE2	1:A:890:ARG:NE	2.41	0.48
1:A:312:LYS:O	1:A:313:GLN:CG	2.62	0.48
1:A:409:LEU:O	1:A:482:ARG:HG2	2.14	0.47
1:A:453:ASN:ND2	1:A:579:ARG:HD2	2.30	0.47
1:A:285:LYS:HE3	1:A:292:TRP:CD1	2.50	0.45
1:A:562:LYS:HE3	1:A:566:GLU:OE2	2.15	0.45
1:A:638:THR:HA	1:A:641:ILE:HG22	1.98	0.45
1:A:805:THR:HG23	1:A:809:MET:HE1	1.99	0.45
1:A:475:TRP:CD1	1:A:475:TRP:N	2.84	0.44
1:A:693:GLN:HG3	8:A:1503:HOH:O	2.18	0.44
1:A:638:THR:HA	1:A:641:ILE:CG2	2.47	0.44
1:A:400:THR:HG23	1:A:426:VAL:HG11	1.99	0.44
1:A:562:LYS:NZ	8:A:1117:HOH:O	2.48	0.44
1:A:638:THR:O	1:A:641:ILE:HG23	2.18	0.44
1:A:452:TYR:HB2	1:A:578:VAL:HG22	2.00	0.44
1:A:357:GLU:N	1:A:357:GLU:CD	2.76	0.43
1:A:801[B]:HIS:H	1:A:801[B]:HIS:CD2	2.38	0.42
1:A:400:THR:HG23	1:A:426:VAL:CG1	2.50	0.42
1:A:428:ASP:OD1	1:A:430:GLY:N	2.51	0.41
1:A:758:TYR:CD1	5:A:1007:DMS:H21	2.56	0.41
1:A:579:ARG:HA	1:A:591:ASP:O	2.20	0.41
1:A:303:TRP:CE3	1:A:593:ILE:HD12	2.56	0.41
1:A:394:CYS:HB3	1:A:486:PHE:CE2	2.56	0.41
1:A:406:ASN:OD1	1:A:423:ARG:NH1	2.53	0.40
1:A:303:TRP:CD2	1:A:593:ILE:HD12	2.57	0.40
5:A:1005:DMS:C1	8:A:1280:HOH:O	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	465/637 (73%)	450 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	518/554 (94%)	505 (98%)	13 (2%)	42	14

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	274	LEU
1	A	290	THR
1	A	356	LYS
1	A	375	LYS
1	A	412	ILE
1	A	423	ARG
1	A	429	SER
1	A	482	ARG
1	A	589	VAL
1	A	635	LEU
1	A	641	ILE
1	A	693	GLN
1	A	887	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	306	HIS
1	A	603	GLN
1	A	618	GLN
1	A	705	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 3 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	PO4	A	1011	-	4,4,4	0.98	0	6,6,6	0.58	0
7	PO4	A	1009	-	4,4,4	4.62	3 (75%)	6,6,6	1.14	1 (16%)
5	DMS	A	1007	-	3,3,3	0.21	0	3,3,3	0.35	0
5	DMS	A	1005	-	3,3,3	0.83	0	3,3,3	0.45	0
6	PEG	A	1012	-	6,6,6	0.17	0	5,5,5	0.09	0
6	PEG	A	1010	-	6,6,6	0.18	0	5,5,5	0.10	0
6	PEG	A	1008	-	6,6,6	0.27	0	5,5,5	0.17	0
4	MES	A	1004[A]	-	12,12,12	0.88	0	15,16,16	0.96	1 (6%)
4	MES	A	1004[B]	-	12,12,12	0.76	0	15,16,16	1.03	1 (6%)
5	DMS	A	1006	-	3,3,3	0.36	0	3,3,3	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PEG	A	1012	-	-	2/4/4/4	-
6	PEG	A	1010	-	-	3/4/4/4	-
6	PEG	A	1008	-	-	1/4/4/4	-
4	MES	A	1004[A]	-	-	3/6/14/14	0/1/1/1
4	MES	A	1004[B]	-	-	2/6/14/14	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	1009	PO4	P-O1	7.22	1.67	1.50
7	A	1009	PO4	P-O2	4.97	1.69	1.54
7	A	1009	PO4	P-O3	2.15	1.60	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1004[A]	MES	O1S-S-C8	-2.26	103.31	106.73
7	A	1009	PO4	O4-P-O3	2.07	114.36	107.91
4	A	1004[B]	MES	O1S-S-C8	-2.05	103.62	106.73

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1004[A]	MES	C7-C8-S-O1S
4	A	1004[A]	MES	C7-C8-S-O3S
4	A	1004[B]	MES	C8-C7-N4-C3
6	A	1008	PEG	O2-C3-C4-O4
6	A	1012	PEG	O2-C3-C4-O4
4	A	1004[B]	MES	C8-C7-N4-C5
4	A	1004[A]	MES	C7-C8-S-O2S
6	A	1010	PEG	O1-C1-C2-O2
6	A	1010	PEG	C4-C3-O2-C2
6	A	1012	PEG	C4-C3-O2-C2
6	A	1010	PEG	O2-C3-C4-O4

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1009	PO4	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1007	DMS	1	0
5	A	1005	DMS	3	0
4	A	1004[B]	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	582/637 (91%)	1.05	121 (20%) 2 2	12, 33, 94, 156	8 (1%)

All (121) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	799	ALA	7.0
1	A	355	PHE	6.5
1	A	413	PHE	6.4
1	A	589	VAL	6.1
1	A	294	TYR	5.6
1	A	412	ILE	5.2
1	A	409	LEU	4.9
1	A	419	TRP	4.7
1	A	475	TRP	4.7
1	A	359	VAL	4.6
1	A	794	THR	4.6
1	A	889	PHE	4.6
1	A	317	ALA	4.5
1	A	314	THR	4.4
1	A	292	TRP	4.4
1	A	284	ILE	4.4
1	A	407	ALA	4.3
1	A	411	ALA	4.3
1	A	426	VAL	4.3
1	A	281	ILE	4.2
1	A	274	LEU	4.2
1	A	473	ALA	4.2
1	A	584	THR	4.0
1	A	800	THR	4.0
1	A	793	THR	3.9
1	A	592	ILE	3.8
1	A	290	THR	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	410	GLY	3.7
1	A	600	GLY	3.7
1	A	431	PHE	3.7
1	A	295	ASP	3.7
1	A	304	ALA	3.6
1	A	593	ILE	3.5
1	A	344	THR	3.5
1	A	635	LEU	3.5
1	A	435	VAL	3.5
1	A	309	TYR	3.4
1	A	299	PRO	3.4
1	A	311	THR	3.4
1	A	358	LYS	3.2
1	A	293	HIS	3.2
1	A	577	VAL	3.2
1	A	356	LYS	3.2
1	A	443	LEU	3.2
1	A	637	VAL	3.2
1	A	583	PRO	3.1
1	A	298	HIS	3.1
1	A	408	ALA	3.1
1	A	363	THR	3.1
1	A	801[A]	HIS	3.1
1	A	441	LEU	3.1
1	A	571	LEU	3.1
1	A	399	PHE	3.1
1	A	428	ASP	3.0
1	A	474	ILE	3.0
1	A	300	TYR	3.0
1	A	454	MET	3.0
1	A	641	ILE	2.9
1	A	305	TYR	2.9
1	A	434	LEU	2.9
1	A	580	VAL	2.9
1	A	512[A]	LEU	2.9
1	A	588	THR	2.9
1	A	746	TRP	2.8
1	A	361	THR	2.8
1	A	744	ALA	2.8
1	A	303	TRP	2.8
1	A	324	VAL	2.8
1	A	455	MET	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	476	TYR	2.8
1	A	417	ASN	2.8
1	A	634	HIS	2.8
1	A	717	ILE	2.7
1	A	418	LYS	2.7
1	A	748	LEU	2.7
1	A	422	ALA	2.6
1	A	578	VAL	2.6
1	A	719[A]	LYS	2.6
1	A	310	GLU	2.6
1	A	447	CYS	2.6
1	A	582	ARG	2.6
1	A	403	VAL	2.6
1	A	891	ARG	2.6
1	A	432	TRP	2.5
1	A	723	VAL	2.5
1	A	884	PRO	2.5
1	A	445	GLY	2.5
1	A	885	SER	2.5
1	A	638	THR	2.5
1	A	478	TRP	2.5
1	A	288	HIS	2.4
1	A	790	THR	2.4
1	A	448	GLU	2.4
1	A	564	LEU	2.4
1	A	308	SER	2.4
1	A	451	VAL	2.4
1	A	472	ARG	2.4
1	A	596	ARG	2.3
1	A	406	ASN	2.3
1	A	427	GLU	2.3
1	A	430	GLY	2.3
1	A	277	ILE	2.3
1	A	791	SER	2.3
1	A	694	TRP	2.2
1	A	544	LEU	2.2
1	A	632	ILE	2.2
1	A	745	GLY	2.2
1	A	286	GLN	2.2
1	A	276	ILE	2.2
1	A	559	GLY	2.2
1	A	319	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	479	LEU	2.1
1	A	484	LEU	2.1
1	A	273	ASN	2.1
1	A	886	MET	2.1
1	A	275	ASP	2.1
1	A	880	THR	2.1
1	A	620	ILE	2.1
1	A	442	HIS	2.1
1	A	645	ASN	2.0
1	A	453	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PO4	A	1011	5/5	0.65	0.18	43,44,48,52	5
6	PEG	A	1008	7/7	0.70	0.17	66,74,78,78	0
5	DMS	A	1006	4/4	0.74	0.27	62,84,86,99	0
7	PO4	A	1009	5/5	0.76	0.18	19,20,24,35	5
6	PEG	A	1012	7/7	0.82	0.17	60,65,71,72	0
6	PEG	A	1010	7/7	0.85	0.15	54,60,65,66	7
5	DMS	A	1005	4/4	0.94	0.13	27,30,30,33	4
5	DMS	A	1007	4/4	0.95	0.11	46,48,50,54	0
2	CL	A	1001	1/1	0.98	0.08	36,36,36,36	0
3	ZN	A	1003	1/1	0.98	0.06	45,45,45,45	0
4	MES	A	1004[A]	12/12	0.98	0.06	17,18,20,21	12
4	MES	A	1004[B]	12/12	0.98	0.06	17,24,31,33	12
3	ZN	A	1002	1/1	1.00	0.01	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.