



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 28, 2026 – 06:50 PM UTC

PDB ID : 8I2E / pdb\_00008i2e  
Title : Crystal structure of Bacillus subtilis LytE in complex with IseA  
Authors : Tandukar, S.; Kwon, E.; Kim, D.Y.  
Deposited on : 2023-01-14  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	 74% 21% .
1	C	160	 81% 15% .
2	B	311	 31% 7% 62%
2	D	311	 30% 8% 62%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein YoeB.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	153	1211	781	204	222	4	0	0	0
1	C	153	1211	781	204	222	4	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLY	-	expression tag	UNP O34841
A	23	SER	-	expression tag	UNP O34841
C	22	GLY	-	expression tag	UNP O34841
C	23	SER	-	expression tag	UNP O34841

- Molecule 2 is a protein called Probable peptidoglycan endopeptidase LytE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	118	906	582	151	170	3	0	0	0
2	D	118	906	582	151	170	3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	24	GLY	-	expression tag	UNP P54421
B	25	SER	-	expression tag	UNP P54421
D	24	GLY	-	expression tag	UNP P54421
D	25	SER	-	expression tag	UNP P54421



SER	SER	SER	THR	TYR	LYS	VAL	LYS	GLY	ASP	SER	LEU	TRP	LYS	ILE	LYS	LYS	TYR	GLY	MET	THR	ILE	ASN	GLU	LEU	LYS	LYS	LEU	ASN	GLY	LEU	LYS	GLY	ASP	LEU	LEU	ARG	VAL	ILE	GLN	TYR	VAL	ASN	LEU	LYS	LEU	LYS	GLY	THR	SER	THR	SER	SER	SER	SER	SER	PRO	LYS	VAL
SER	SER	SER	THR	SER	SER	THR	TYR	VAL	LYS	SER	GLY	ASP	SER	LEU	ILE	LYS	ALA	SER	LYS	TYR	THR	THR	THR	VAL	LEU	SER	LYS	LEU	LYS	SER	LEU	ASN	GLY	ASP	VAL	ILE	TYR	VAL	ASN	GLN	VAL	LEU	LYS	LEU	LYS	VAL	LYS	THR	SER	THR	SER	SER	SER	SER	SER	PRO	LYS	ALA
SER	SER	SER	SER	SER	SER	SER	SER	THR	THR	LYS	THR	THR	S217	L218	V231	K237	G238	G239	F245	D246	C247	S248	I251	I255	Q258	R263	T264	S265	I276	T288	M297	Y300	I301	G302	H308	D312	S313	V314	I329	R333	F334																	

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.01Å 87.01Å 217.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.89 – 3.20 46.89 – 3.20	Depositor EDS
% Data completeness (in resolution range)	97.9 (46.89-3.20) 97.9 (46.89-3.20)	Depositor EDS
$R_{merge}$	0.20	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 3.19Å)	Xtrriage
Refinement program	PHENIX 1.18	Depositor
R, $R_{free}$	0.234 , 0.282 0.241 , 0.290	Depositor DCC
$R_{free}$ test set	663 reflections (4.57%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	80.3	Xtrriage
Anisotropy	0.415	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4234	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/1239	0.85	0/1670
1	C	0.47	0/1239	0.77	0/1670
2	B	0.50	0/933	0.80	0/1263
2	D	0.46	0/933	0.68	0/1263
All	All	0.49	0/4344	0.78	0/5866

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1211	0	1235	24	0
1	C	1211	0	1235	17	0
2	B	906	0	875	16	1
2	D	906	0	875	19	1
All	All	4234	0	4220	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 75 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:LEU:HD11	1:A:154:ASP:HB2	1.54	0.88
1:C:126:LEU:HD11	1:C:154:ASP:HB2	1.64	0.80
2:B:248:SER:HB2	2:B:263:ARG:HH21	1.49	0.76
1:A:168:GLU:HB2	1:A:173:LYS:HE2	1.67	0.74
1:C:145:THR:HG22	1:C:164:THR:HG23	1.74	0.67

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:325:LYS:NZ	2:D:312:ASP:OD2[3_555]	2.19	0.01

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/160 (94%)	146 (97%)	5 (3%)	0	100	100
1	C	151/160 (94%)	149 (99%)	2 (1%)	0	100	100
2	B	116/311 (37%)	113 (97%)	3 (3%)	0	100	100
2	D	116/311 (37%)	112 (97%)	4 (3%)	0	100	100
All	All	534/942 (57%)	520 (97%)	14 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/138 (96%)	133 (100%)	0	100	100
1	C	133/138 (96%)	133 (100%)	0	100	100
2	B	98/276 (36%)	98 (100%)	0	100	100
2	D	98/276 (36%)	98 (100%)	0	100	100
All	All	462/828 (56%)	462 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	89	ASN
2	B	256	ASN
1	C	34	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/160 (95%)	0.30	1 (0%) 84 69	52, 72, 95, 117	0
1	C	153/160 (95%)	0.31	4 (2%) 57 37	52, 72, 104, 148	0
2	B	118/311 (37%)	0.29	0 100 100	54, 68, 92, 123	0
2	D	118/311 (37%)	0.36	1 (0%) 82 67	58, 77, 103, 110	0
All	All	542/942 (57%)	0.32	6 (1%) 78 61	52, 72, 99, 148	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	312	ASP	2.3
1	C	141	GLY	2.3
1	C	124	ASP	2.2
1	C	123	GLY	2.1
1	C	30	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.