



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:04 AM UTC

PDB ID : 2I37 / pdb_00002i37
Title : Crystal structure of a photoactivated rhodopsin
Authors : Lodowski, D.T.; Stenkamp, R.E.; Salom, D.; Le Trong, I.; Palczewski, K.
Deposited on : 2006-08-17
Resolution : 4.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 4.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1077 (4.50-3.82)
Clashscore	190562	1019 (4.48-3.84)
Ramachandran outliers	187476	1035 (4.50-3.82)
Sidechain outliers	187428	1022 (4.50-3.82)
RSRZ outliers	180081	1074 (4.50-3.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	349	7% 76% 14% • 9%
1	B	349	6% 76% 14% • 8%
1	C	349	6% 75% 14% • 9%
2	D	3	67% 33%
3	E	2	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	2	 100%
3	I	2	 100%
4	F	5	 80% 20%
5	H	4	 25% 50% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAG	G	1	X	-	-	-
3	NAG	I	1	X	-	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 7814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Rhodopsin.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2518	1680	385	427	26			
1	B	320	Total	C	N	O	S	0	0	0
			2534	1689	387	432	26			
1	C	319	Total	C	N	O	S	0	0	0
			2525	1682	387	430	26			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
3	I	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	F	5	Total	C	N	O	0	0	0
			64	36	3	25			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	H	4	Total	C	N	O	0	0	0
			50	28	2	20			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Rhodopsin




• Molecule 1: Rhodopsin



• Molecule 1: Rhodopsin





- Molecule 2: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 67% 33%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G: 100%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I: 100%




- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-beta-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F: 80% 20%



- Molecule 5: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 1 2	Depositor
Cell constants a, b, c, α , β , γ	161.25Å 161.25Å 143.44Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 4.15 30.00 – 4.15	Depositor EDS
% Data completeness (in resolution range)	76.8 (30.00-4.15) 76.5 (30.00-4.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.37 (at 4.11Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.377 , 0.382 0.376 , 0.379	Depositor DCC
R_{free} test set	645 reflections (3.98%)	wwPDB-VP
Wilson B-factor (Å ²)	233.0	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.21 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.049 for -h,-k,l	Xtriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	7814	wwPDB-VP
Average B, all atoms (Å ²)	183.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, ACE, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/2596	0.75	1/3538 (0.0%)
1	B	0.56	0/2611	0.78	2/3558 (0.1%)
1	C	0.54	0/2601	0.76	1/3541 (0.0%)
All	All	0.54	0/7808	0.76	4/10637 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	170	PRO	N-CA-C	5.79	117.76	110.70
1	B	170	PRO	N-CA-C	5.64	117.58	110.70
1	A	170	PRO	N-CA-C	5.62	117.56	110.70
1	B	290	ILE	CB-CA-C	-5.04	108.91	113.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2518	0	2491	23	0
1	B	2534	0	2513	24	0
1	C	2525	0	2494	24	0
2	D	39	0	34	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	28	0	25	0	0
3	G	28	0	25	0	0
3	I	28	0	25	0	0
4	F	64	0	55	2	0
5	H	50	0	43	4	0
All	All	7814	0	7705	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:16:LYS:O	1:C:7:PRO:HA	1.94	0.67
1:B:322:CYS:O	1:B:323:CYS:SG	2.56	0.64
1:C:4:THR:CG2	5:H:1:NAG:H83	2.32	0.60
1:A:20:VAL:O	2:D:1:NAG:O5	2.23	0.55
1:A:219:ILE:O	1:A:219:ILE:HG22	2.08	0.54
1:C:219:ILE:HG22	1:C:219:ILE:O	2.07	0.54
1:B:219:ILE:HG22	1:B:219:ILE:O	2.07	0.54
1:A:179:ILE:HG13	1:A:180:PRO:HD2	1.90	0.54
1:A:96:TYR:OH	1:B:100:HIS:NE2	2.34	0.53
1:C:179:ILE:HG13	1:C:180:PRO:HD2	1.91	0.53
1:B:179:ILE:HG13	1:B:180:PRO:HD2	1.92	0.52
1:C:4:THR:HG21	5:H:1:NAG:H83	1.91	0.52
1:B:169:ALA:N	1:B:170:PRO:HD2	2.26	0.51
1:C:169:ALA:N	1:C:170:PRO:HD2	2.25	0.51
1:A:169:ALA:N	1:A:170:PRO:HD2	2.26	0.50
1:B:137:VAL:O	1:B:137:VAL:HG12	2.12	0.49
1:A:52:PHE:HB2	1:A:87:VAL:HG11	1.94	0.49
1:B:303:PRO:HA	1:B:306:TYR:HB3	1.95	0.49
1:A:303:PRO:HA	1:A:306:TYR:HB3	1.95	0.49
1:B:52:PHE:HB2	1:B:87:VAL:HG11	1.93	0.49
1:C:52:PHE:HB2	1:C:87:VAL:HG11	1.95	0.49
1:C:303:PRO:HA	1:C:306:TYR:HB3	1.95	0.49
1:A:137:VAL:HG12	1:A:137:VAL:O	2.13	0.48
1:C:137:VAL:HG12	1:C:137:VAL:O	2.13	0.48
1:B:68:LEU:HD11	1:B:316:CYS:SG	2.54	0.47
1:B:20:VAL:HA	1:B:30:TYR:CZ	2.50	0.46
4:F:1:NAG:H82	5:H:1:NAG:H82	1.98	0.46
1:C:68:LEU:HD11	1:C:316:CYS:SG	2.55	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:ILE:HB	1:B:215:PRO:HD3	1.98	0.45
1:C:318:VAL:O	1:C:322:CYS:N	2.49	0.45
1:C:44:MET:HE1	1:C:91:PHE:CD1	2.51	0.45
1:A:214:ILE:HB	1:A:215:PRO:HD3	1.99	0.45
1:B:44:MET:HE1	1:B:91:PHE:CD1	2.51	0.45
1:C:214:ILE:HB	1:C:215:PRO:HD3	1.99	0.45
1:A:44:MET:HE1	1:A:91:PHE:CD1	2.51	0.45
1:A:27:PRO:HA	1:A:102:TYR:HB3	1.98	0.45
1:A:49:MET:SD	1:B:52:PHE:HD2	2.39	0.45
1:A:68:LEU:HD11	1:A:316:CYS:SG	2.57	0.45
1:B:318:VAL:O	1:B:322:CYS:N	2.50	0.44
1:C:20:VAL:HA	1:C:30:TYR:CZ	2.53	0.44
1:A:20:VAL:HA	1:A:30:TYR:CZ	2.53	0.44
1:A:105:PHE:HB3	1:A:109:GLY:HA3	2.00	0.43
1:C:105:PHE:HB3	1:C:109:GLY:HA3	1.99	0.43
1:B:105:PHE:HB3	1:B:109:GLY:HA3	2.00	0.43
1:A:17:THR:O	1:A:18:GLY:C	2.61	0.43
1:B:23:PRO:HA	1:B:28:GLN:HE21	1.83	0.43
1:B:72:LEU:HD21	1:B:310:ASN:ND2	2.33	0.43
4:F:1:NAG:O7	5:H:1:NAG:O3	2.26	0.43
1:B:27:PRO:HA	1:B:102:TYR:HB3	2.00	0.43
1:C:17:THR:O	1:C:18:GLY:C	2.62	0.42
1:C:27:PRO:HA	1:C:102:TYR:HB3	1.99	0.42
1:C:322:CYS:O	1:C:323:CYS:SG	2.77	0.42
1:A:194:PRO:O	1:A:195:HIS:C	2.63	0.42
1:A:219:ILE:O	1:A:219:ILE:CG2	2.68	0.42
1:A:72:LEU:HD21	1:A:310:ASN:ND2	2.35	0.41
1:B:219:ILE:O	1:B:219:ILE:CG2	2.68	0.41
1:A:23:PRO:HA	1:A:28:GLN:HE21	1.86	0.41
1:B:97:THR:HG21	1:B:185:CYS:HA	2.03	0.41
1:C:97:THR:HG21	1:C:185:CYS:HA	2.02	0.41
1:C:72:LEU:HD21	1:C:310:ASN:ND2	2.36	0.41
1:B:16:LYS:HD3	1:C:195:HIS:CD2	2.55	0.41
1:C:271:VAL:HG13	1:C:287:PHE:CZ	2.56	0.41
1:C:219:ILE:O	1:C:219:ILE:CG2	2.68	0.41
1:B:17:THR:O	1:B:18:GLY:C	2.64	0.41
1:A:102:TYR:CE2	1:A:104:VAL:HG12	2.57	0.40
1:A:271:VAL:HG13	1:A:287:PHE:CZ	2.56	0.40
1:A:290:ILE:N	1:A:291:PRO:HD2	2.36	0.40
1:B:241:ALA:HB1	1:B:244:GLN:HB2	2.04	0.40
1:C:145:ASN:O	1:C:145:ASN:ND2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/349 (89%)	272 (87%)	35 (11%)	5 (2%)	7	36
1	B	314/349 (90%)	271 (86%)	38 (12%)	5 (2%)	7	36
1	C	312/349 (89%)	269 (86%)	37 (12%)	6 (2%)	6	32
All	All	938/1047 (90%)	812 (87%)	110 (12%)	16 (2%)	7	35

All (16) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	322	CYS
1	A	18	GLY
1	A	323	CYS
1	B	18	GLY
1	C	18	GLY
1	C	324	GLY
1	B	322	CYS
1	B	324	GLY
1	C	330	ASP
1	A	194	PRO
1	A	285	PRO
1	B	194	PRO
1	B	285	PRO
1	C	194	PRO
1	C	285	PRO
1	C	329	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/296 (92%)	254 (94%)	17 (6%)	16	39
1	B	273/296 (92%)	256 (94%)	17 (6%)	16	39
1	C	272/296 (92%)	254 (93%)	18 (7%)	15	38
All	All	816/888 (92%)	764 (94%)	52 (6%)	16	39

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	ARG
1	A	97	THR
1	A	102	TYR
1	A	103	PHE
1	A	104	VAL
1	A	119	LEU
1	A	122	GLU
1	A	135	ARG
1	A	157	VAL
1	A	165	LEU
1	A	172	LEU
1	A	175	TRP
1	A	179	ILE
1	A	216	LEU
1	A	221	PHE
1	A	263	ILE
1	A	286	ILE
1	B	97	THR
1	B	102	TYR
1	B	103	PHE
1	B	104	VAL
1	B	119	LEU
1	B	123	ILE
1	B	135	ARG
1	B	157	VAL
1	B	165	LEU
1	B	172	LEU
1	B	175	TRP
1	B	179	ILE
1	B	216	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	221	PHE
1	B	263	ILE
1	B	286	ILE
1	B	322	CYS
1	C	21	ARG
1	C	97	THR
1	C	102	TYR
1	C	103	PHE
1	C	104	VAL
1	C	119	LEU
1	C	135	ARG
1	C	157	VAL
1	C	165	LEU
1	C	172	LEU
1	C	175	TRP
1	C	179	ILE
1	C	216	LEU
1	C	221	PHE
1	C	263	ILE
1	C	286	ILE
1	C	322	CYS
1	C	332	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	73	ASN
1	A	151	ASN
1	A	225	GLN
1	A	310	ASN
1	A	326	ASN
1	B	36	GLN
1	B	73	ASN
1	B	151	ASN
1	B	225	GLN
1	B	310	ASN
1	B	315	ASN
1	C	36	GLN
1	C	73	ASN
1	C	145	ASN
1	C	151	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	225	GLN
1	C	310	ASN
1	C	312	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

18 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	D	1	2,1	14,14,15	0.59	0	17,19,21	1.48	3 (17%)
2	NAG	D	2	2	14,14,15	0.52	0	17,19,21	1.49	3 (17%)
2	MAN	D	3	2	11,11,12	0.54	0	15,15,17	1.42	1 (6%)
3	NAG	E	1	3,1	14,14,15	0.57	0	17,19,21	0.96	1 (5%)
3	NAG	E	2	3	14,14,15	0.83	1 (7%)	17,19,21	1.58	3 (17%)
4	NAG	F	1	4,1	14,14,15	0.67	0	17,19,21	1.18	1 (5%)
4	NAG	F	2	4	14,14,15	0.45	0	17,19,21	1.17	2 (11%)
4	MAN	F	3	4	11,11,12	0.79	0	15,15,17	1.63	3 (20%)
4	BMA	F	4	4	11,11,12	0.63	0	15,15,17	1.12	1 (6%)
4	NAG	F	5	4	14,14,15	0.65	0	17,19,21	1.11	1 (5%)
3	NAG	G	1	3,1	14,14,15	0.59	0	17,19,21	1.44	3 (17%)
3	NAG	G	2	3	14,14,15	0.60	0	17,19,21	0.90	1 (5%)
5	NAG	H	1	5,1	14,14,15	0.47	0	17,19,21	0.93	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	H	2	5	14,14,15	0.63	0	17,19,21	0.90	0
5	MAN	H	3	5	11,11,12	0.53	0	15,15,17	2.48	4 (26%)
5	MAN	H	4	5	11,11,12	0.51	0	15,15,17	1.32	1 (6%)
3	NAG	I	1	3,1	14,14,15	0.51	0	17,19,21	1.35	3 (17%)
3	NAG	I	2	3	14,14,15	0.73	0	17,19,21	1.35	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	D	1	2,1	-	1/6/23/26	0/1/1/1
2	NAG	D	2	2	-	4/6/23/26	0/1/1/1
2	MAN	D	3	2	-	2/2/19/22	1/1/1/1
3	NAG	E	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	E	2	3	-	5/6/23/26	0/1/1/1
4	NAG	F	1	4,1	-	5/6/23/26	0/1/1/1
4	NAG	F	2	4	-	4/6/23/26	0/1/1/1
4	MAN	F	3	4	-	1/2/19/22	0/1/1/1
4	BMA	F	4	4	-	1/2/19/22	0/1/1/1
4	NAG	F	5	4	-	4/6/23/26	0/1/1/1
3	NAG	G	1	3,1	1/1/5/7	4/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
5	NAG	H	1	5,1	-	4/6/23/26	0/1/1/1
5	NAG	H	2	5	-	5/6/23/26	0/1/1/1
5	MAN	H	3	5	-	2/2/19/22	0/1/1/1
5	MAN	H	4	5	-	0/2/19/22	0/1/1/1
3	NAG	I	1	3,1	1/1/5/7	4/6/23/26	0/1/1/1
3	NAG	I	2	3	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	2	NAG	C1-C2	2.36	1.55	1.52

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	H	3	MAN	C1-O5-C5	8.04	122.97	112.19
2	D	3	MAN	C1-O5-C5	4.52	118.24	112.19
4	F	3	MAN	C1-C2-C3	4.20	115.75	109.64
5	H	4	MAN	C1-O5-C5	4.17	117.78	112.19
2	D	1	NAG	C1-O5-C5	3.85	117.35	112.19
3	E	2	NAG	C1-O5-C5	3.68	117.11	112.19
3	I	1	NAG	C1-O5-C5	3.65	117.08	112.19
2	D	2	NAG	C2-N2-C7	-3.60	118.07	122.90
2	D	2	NAG	O5-C1-C2	-3.60	105.72	111.29
4	F	2	NAG	C1-O5-C5	3.39	116.73	112.19
4	F	5	NAG	C4-C3-C2	3.04	115.47	111.02
3	G	1	NAG	O5-C1-C2	-3.00	106.64	111.29
3	G	1	NAG	C4-C3-C2	3.00	115.41	111.02
5	H	3	MAN	O3-C3-C2	2.91	115.98	110.05
3	I	2	NAG	C2-N2-C7	2.76	126.59	122.90
4	F	4	BMA	C1-O5-C5	2.73	115.84	112.19
3	I	2	NAG	C4-C3-C2	2.69	114.96	111.02
3	E	1	NAG	C4-C3-C2	2.66	114.92	111.02
3	G	1	NAG	C1-O5-C5	2.60	115.68	112.19
4	F	3	MAN	O3-C3-C4	2.59	116.49	110.38
3	G	2	NAG	C1-O5-C5	2.54	115.59	112.19
5	H	3	MAN	C1-C2-C3	2.28	112.96	109.64
3	I	2	NAG	C8-C7-N2	2.23	119.82	116.12
3	I	1	NAG	O5-C1-C2	2.18	114.66	111.29
4	F	2	NAG	C3-C4-C5	-2.17	106.29	110.23
4	F	1	NAG	C1-C2-N2	-2.17	107.02	110.43
3	E	2	NAG	C2-N2-C7	2.14	125.77	122.90
4	F	3	MAN	C1-O5-C5	2.11	115.02	112.19
3	E	2	NAG	O5-C1-C2	2.11	114.56	111.29
2	D	1	NAG	O5-C1-C2	-2.05	108.11	111.29
5	H	3	MAN	O5-C1-C2	2.05	115.67	110.79
2	D	2	NAG	C1-O5-C5	2.02	114.89	112.19
3	I	1	NAG	C3-C4-C5	-2.02	106.58	110.23
5	H	1	NAG	O5-C5-C6	2.01	111.58	107.66
2	D	1	NAG	O4-C4-C3	2.01	115.10	110.38

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	G	1	NAG	C1
3	I	1	NAG	C1

All (54) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2	NAG	C8-C7-N2-C2
2	D	2	NAG	O7-C7-N2-C2
3	E	1	NAG	C8-C7-N2-C2
3	E	1	NAG	O7-C7-N2-C2
3	E	2	NAG	C1-C2-N2-C7
3	G	1	NAG	C8-C7-N2-C2
3	G	1	NAG	O7-C7-N2-C2
3	G	2	NAG	C8-C7-N2-C2
3	G	2	NAG	O7-C7-N2-C2
3	I	1	NAG	O7-C7-N2-C2
4	F	2	NAG	C8-C7-N2-C2
4	F	2	NAG	O7-C7-N2-C2
4	F	5	NAG	C8-C7-N2-C2
4	F	5	NAG	O7-C7-N2-C2
5	H	1	NAG	C8-C7-N2-C2
5	H	1	NAG	O7-C7-N2-C2
5	H	2	NAG	C8-C7-N2-C2
5	H	2	NAG	O7-C7-N2-C2
3	I	1	NAG	C8-C7-N2-C2
4	F	2	NAG	O5-C5-C6-O6
4	F	1	NAG	O5-C5-C6-O6
4	F	5	NAG	O5-C5-C6-O6
4	F	2	NAG	C4-C5-C6-O6
4	F	1	NAG	C4-C5-C6-O6
4	F	5	NAG	C4-C5-C6-O6
3	I	2	NAG	C8-C7-N2-C2
4	F	1	NAG	C8-C7-N2-C2
4	F	1	NAG	O7-C7-N2-C2
2	D	3	MAN	O5-C5-C6-O6
3	G	1	NAG	O5-C5-C6-O6
3	E	1	NAG	O5-C5-C6-O6
3	G	1	NAG	C4-C5-C6-O6
2	D	3	MAN	C4-C5-C6-O6
3	E	1	NAG	C4-C5-C6-O6
3	I	2	NAG	O7-C7-N2-C2
3	I	1	NAG	O5-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
5	H	2	NAG	C4-C5-C6-O6
5	H	3	MAN	C4-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
5	H	2	NAG	O5-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	E	2	NAG	C8-C7-N2-C2
5	H	3	MAN	O5-C5-C6-O6
4	F	3	MAN	O5-C5-C6-O6
5	H	1	NAG	O5-C5-C6-O6
4	F	4	BMA	O5-C5-C6-O6
3	E	2	NAG	O7-C7-N2-C2
3	I	1	NAG	C4-C5-C6-O6
4	F	1	NAG	C3-C2-N2-C7
5	H	1	NAG	C3-C2-N2-C7
5	H	2	NAG	C3-C2-N2-C7
2	D	2	NAG	C4-C5-C6-O6


All (1) ring outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	3	MAN	C1-C2-C3-C4-C5-O5


3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	H	1	NAG	4	0
4	F	1	NAG	2	0
2	D	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/349 (90%)	0.33	23 (7%) 21 22	183, 183, 183, 183	0
1	B	319/349 (91%)	0.43	20 (6%) 26 25	183, 183, 183, 183	0
1	C	318/349 (91%)	0.16	21 (6%) 24 24	183, 183, 183, 183	0
All	All	954/1047 (91%)	0.31	64 (6%) 24 23	183, 183, 183, 183	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	121	GLY	13.4
1	B	120	GLY	9.8
1	B	124	ALA	5.9
1	A	183	MET	5.6
1	B	223	TYR	5.3
1	B	125	LEU	5.2
1	C	5	GLU	5.1
1	A	191	TYR	5.1
1	B	128	LEU	4.9
1	C	118	THR	4.8
1	A	10	TYR	4.7
1	B	182	GLY	4.6
1	C	121	GLY	4.6
1	A	4	THR	4.5
1	B	122	GLU	4.4
1	A	317	MET	4.3
1	B	296	LYS	4.3
1	C	182	GLY	4.2
1	A	276	PHE	4.2
1	A	296	LYS	4.0
1	B	304	VAL	3.9
1	C	251	THR	3.8
1	C	191	TYR	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	117	ALA	3.7
1	A	121	GLY	3.7
1	B	79	LEU	3.5
1	A	124	ALA	3.5
1	C	63	VAL	3.5
1	B	76	LEU	3.4
1	B	212	PHE	3.4
1	C	296	LYS	3.4
1	A	86	MET	3.4
1	A	286	ILE	3.3
1	B	72	LEU	3.3
1	A	321	LEU	3.2
1	C	12	PRO	3.2
1	A	11	VAL	3.2
1	A	189	ILE	2.8
1	C	173	VAL	2.8
1	B	70	THR	2.7
1	B	187	CYS	2.7
1	A	193	THR	2.6
1	A	197	GLU	2.6
1	C	286	ILE	2.6
1	C	298	SER	2.5
1	A	92	THR	2.5
1	A	5	GLU	2.5
1	C	190	ASP	2.4
1	C	302	ASN	2.4
1	C	183	MET	2.4
1	C	9	PHE	2.3
1	B	269	ALA	2.3
1	C	48	ILE	2.3
1	B	243	THR	2.3
1	C	247	GLU	2.3
1	A	225	GLN	2.3
1	C	31	LEU	2.2
1	A	31	LEU	2.2
1	C	184	GLN	2.2
1	A	192	TYR	2.1
1	A	272	ALA	2.1
1	B	118	THR	2.1
1	C	292	ALA	2.1
1	A	85	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	G	2	14/15	-0.28	0.13	183,183,183,183	0
3	NAG	I	2	14/15	0.19	0.10	183,183,183,183	0
3	NAG	G	1	14/15	0.21	0.18	183,183,183,183	0
5	MAN	H	4	11/12	0.31	0.12	183,183,183,183	0
3	NAG	E	2	14/15	0.37	0.15	183,183,183,183	0
5	MAN	H	3	11/12	0.51	0.10	183,183,183,183	0
3	NAG	I	1	14/15	0.52	0.14	183,183,183,183	0
2	MAN	D	3	11/12	0.56	0.16	183,183,183,183	0
4	NAG	F	5	14/15	0.57	0.10	183,183,183,183	1
4	MAN	F	3	11/12	0.64	0.11	183,183,183,183	0
4	BMA	F	4	11/12	0.84	0.04	183,183,183,183	0
2	NAG	D	2	14/15	0.88	0.11	183,183,183,183	0
3	NAG	E	1	14/15	0.89	0.09	183,183,183,183	0
4	NAG	F	2	14/15	0.90	0.08	183,183,183,183	0
5	NAG	H	2	14/15	0.90	0.09	183,183,183,183	0
5	NAG	H	1	14/15	0.92	0.21	183,183,183,183	0
2	NAG	D	1	14/15	0.93	0.21	183,183,183,183	0
4	NAG	F	1	14/15	0.93	0.13	183,183,183,183	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain I:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.