



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:46 AM UTC

PDB ID : 4I3G / pdb_00004i3g
Title : Crystal Structure of DesR, a beta-glucosidase from *Streptomyces venezuelae* in complex with D-glucose.
Authors : Zmudka, M.W.; Holden, H.M.
Deposited on : 2012-11-26
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	829	4% 88% 5% 6%
1	B	829	3% 90% • 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	B	907	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 13233 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Beta-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	778	Total	C	N	O	S	0	6	0
			5785	3621	1019	1128	17			
1	B	780	Total	C	N	O	S	0	1	0
			5781	3618	1020	1127	16			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	VAL	-	expression tag	UNP C5IXN7
A	560	LYS	GLU	conflict	UNP C5IXN7
A	726	THR	ALA	conflict	UNP C5IXN7
B	1	VAL	-	expression tag	UNP C5IXN7
B	560	LYS	GLU	conflict	UNP C5IXN7
B	726	THR	ALA	conflict	UNP C5IXN7

- Molecule 2 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			12	6	6		
2	B	1	Total	C	O	0	0
			12	6	6		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 3[N-MORPHOLINO]PROPANE SULFONIC ACID (CCD ID: MPO) (formula: $C_7H_{15}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			13	7	1	4	1		
4	B	1	Total	C	N	O	S	0	0
			13	7	1	4	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	784	Total	O	0	0
			784	784		
6	B	757	Total	O	0	0
			757	757		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-glucosidase



• Molecule 1: Beta-glucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	131.27Å 198.31Å 67.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.40 30.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	94.7 (30.00-1.40) 94.7 (30.00-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.172 , 0.199 0.172 , 0.200	Depositor DCC
R_{free} test set	17334 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.022	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 39.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13233	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MPO, SO4, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/5922	1.14	7/8068 (0.1%)
1	B	0.69	0/5904	1.15	4/8044 (0.0%)
All	All	0.70	0/11826	1.14	11/16112 (0.1%)

There are no bond length outliers.

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	219	ARG	NE-CZ-NH2	-6.73	113.15	119.20
1	B	537	THR	CA-C-N	-5.81	113.98	120.14
1	B	537	THR	C-N-CA	-5.81	113.98	120.14
1	B	334	GLU	CA-CB-CG	5.62	125.34	114.10
1	A	127	MET	CG-SD-CE	5.55	113.10	100.90
1	A	219	ARG	NH1-CZ-NH2	5.50	126.45	119.30
1	A	187	GLY	N-CA-C	-5.37	107.66	114.16
1	A	74	ASP	O-C-N	5.29	126.00	121.30
1	A	131	TYR	CA-CB-CG	5.07	123.02	113.90
1	B	131	TYR	CA-CB-CG	5.05	122.98	113.90
1	A	330	THR	O-C-N	5.04	127.26	122.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5785	0	5738	29	0
1	B	5781	0	5721	26	0
2	A	12	0	12	1	0
2	B	12	0	12	1	0
3	A	40	0	0	0	0
3	B	20	0	0	1	0
4	A	13	0	15	0	0
4	B	13	0	15	1	0
5	A	8	0	12	0	0
5	B	8	0	12	4	0
6	A	784	0	0	4	0
6	B	757	0	0	4	0
All	All	13233	0	11537	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:475:THR:CG2	1:B:524:THR:HG23	1.64	1.25
1:B:475:THR:HG21	1:B:524:THR:CG2	1.81	1.11
1:B:475:THR:HG21	1:B:524:THR:HG23	1.13	1.10
1:A:680[B]:GLN:OE1	6:A:1760:HOH:O	1.84	0.95
1:B:475:THR:CG2	1:B:524:THR:CG2	2.45	0.91
1:A:820:ARG:HG2	1:A:821:GLY:N	1.91	0.85
1:B:511:GLY:HA2	5:B:907:EDO:H22	1.57	0.84
1:B:391:LYS:HE3	6:B:1442:HOH:O	1.86	0.76
1:B:412:HIS:NE2	5:B:907:EDO:H21	2.01	0.76
1:B:133:LYS:HE3	6:B:1626:HOH:O	1.86	0.73
1:A:464:GLU:HG3	1:A:465:PRO:HD2	1.73	0.70
1:A:820:ARG:HG2	1:A:821:GLY:H	1.59	0.66
5:B:907:EDO:H12	6:B:1677:HOH:O	1.95	0.65
1:B:204:ASN:C	1:B:204:ASN:HD22	2.04	0.65
1:B:807:ASN:ND2	1:B:826:ASN:OD1	2.32	0.62
1:B:309:LYS:HE2	1:B:310:PHE:CZ	2.34	0.61
1:A:560:LYS:HE3	1:A:564:LYS:HE3	1.83	0.61
1:B:207:PHE:HB3	4:B:906:MPO:H52	1.83	0.60
1:B:412:HIS:NE2	5:B:907:EDO:C2	2.67	0.58
1:A:273:ASP:OD2	2:A:901:BGC:H1	2.03	0.58
1:A:464:GLU:O	1:A:465:PRO:C	2.47	0.57
1:A:798:THR:HG22	1:A:798:THR:O	2.06	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:818:ASP:O	1:A:820:ARG:NH1	2.40	0.54
1:B:273:ASP:OD2	2:B:901:BGC:H1	2.08	0.53
1:A:464:GLU:HG3	1:A:465:PRO:CD	2.39	0.52
1:A:798:THR:O	1:A:798:THR:CG2	2.57	0.52
1:A:204:ASN:C	1:A:204:ASN:HD22	2.18	0.52
1:A:465:PRO:HB3	1:A:535:SER:O	2.10	0.52
1:A:133:LYS:HE3	6:A:1269:HOH:O	2.11	0.49
1:A:820:ARG:NH2	6:A:1776:HOH:O	2.45	0.49
1:A:820:ARG:CG	1:A:821:GLY:N	2.72	0.49
3:B:902:SO4:O3	6:B:1302:HOH:O	2.20	0.49
1:B:181:GLN:HE21	1:B:185:ILE:HD11	1.77	0.48
1:A:127:MET:HA	1:A:127:MET:CE	2.44	0.48
1:A:448:ILE:HD12	1:A:543:LEU:HG	1.96	0.48
1:A:74:ASP:HB2	1:A:79:ASN:OD1	2.14	0.48
1:B:577:THR:HG22	1:B:578:GLU:O	2.13	0.47
1:B:489:ARG:HD3	1:B:513:VAL:HG22	1.96	0.47
1:A:688:HIS:O	1:A:693:TRP:HB2	2.15	0.47
1:A:309:LYS:HE2	1:A:310:PHE:CZ	2.51	0.46
1:B:74:ASP:HB2	1:B:79:ASN:OD1	2.16	0.46
1:A:488:VAL:CG1	1:A:541:LEU:HD11	2.47	0.45
1:B:154:ASN:HD21	1:B:169:GLU:C	2.25	0.45
1:A:395:VAL:C	1:A:396:ILE:HG13	2.43	0.44
1:B:404:LYS:O	1:B:632:PRO:HD2	2.18	0.44
1:A:154:ASN:HD21	1:A:169:GLU:C	2.26	0.43
1:B:204:ASN:C	1:B:204:ASN:ND2	2.75	0.43
1:A:404:LYS:O	1:A:632:PRO:HD2	2.19	0.42
1:B:309:LYS:HE2	1:B:310:PHE:CE2	2.54	0.42
1:B:79:ASN:OD1	1:B:79:ASN:C	2.63	0.42
1:A:610:ASN:HA	1:A:630:TRP:O	2.20	0.42
1:B:239:CYS:SG	1:B:272:SER:HA	2.60	0.42
1:A:215:GLU:HG2	6:A:1325:HOH:O	2.20	0.41
1:A:181:GLN:HE21	1:A:185:ILE:HD11	1.85	0.41
1:A:494:TYR:CD2	1:A:507:GLY:HA3	2.56	0.41
1:B:367:LYS:HB3	1:B:367:LYS:HE2	1.89	0.41
1:B:475:THR:HG23	1:B:524:THR:CG2	2.42	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	782/829 (94%)	765 (98%)	16 (2%)	1 (0%)	48	21
1	B	779/829 (94%)	767 (98%)	12 (2%)	0	100	100
All	All	1561/1658 (94%)	1532 (98%)	28 (2%)	1 (0%)	48	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	297	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	602/632 (95%)	596 (99%)	6 (1%)	68	40
1	B	599/632 (95%)	595 (99%)	4 (1%)	76	52
All	All	1201/1264 (95%)	1191 (99%)	10 (1%)	73	48

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	ASN
1	A	127	MET
1	A	204	ASN
1	A	467	LYS
1	A	815	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	820	ARG
1	B	101	ASN
1	B	204	ASN
1	B	807	ASN
1	B	815	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (26) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	154	ASN
1	A	181	GLN
1	A	200	ASN
1	A	204	ASN
1	A	267	GLN
1	A	371	ASN
1	A	458	ASN
1	A	656	GLN
1	A	663	ASN
1	A	664	GLN
1	A	681	GLN
1	A	744	GLN
1	A	790	GLN
1	A	811	GLN
1	B	154	ASN
1	B	181	GLN
1	B	200	ASN
1	B	204	ASN
1	B	371	ASN
1	B	656	GLN
1	B	663	ASN
1	B	664	GLN
1	B	681	GLN
1	B	744	GLN
1	B	792	GLN
1	B	807	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	B	907	-	3,3,3	0.35	0	2,2,2	0.12	0
4	MPO	B	906	-	13,13,13	2.15	1 (7%)	17,17,17	2.27	6 (35%)
3	SO4	B	902	-	4,4,4	0.51	0	6,6,6	0.39	0
2	BGC	B	901	-	12,12,12	0.52	0	17,17,17	0.97	1 (5%)
3	SO4	B	903	-	4,4,4	0.26	0	6,6,6	0.39	0
3	SO4	A	905	-	4,4,4	0.61	0	6,6,6	0.39	0
3	SO4	B	905	-	4,4,4	0.26	0	6,6,6	0.27	0
2	BGC	A	901	-	12,12,12	0.80	0	17,17,17	0.95	1 (5%)
3	SO4	A	903	-	4,4,4	0.27	0	6,6,6	0.32	0
5	EDO	A	911	-	3,3,3	0.46	0	2,2,2	0.30	0
3	SO4	A	904	-	4,4,4	0.25	0	6,6,6	0.52	0
4	MPO	A	910	-	13,13,13	2.18	1 (7%)	17,17,17	2.64	6 (35%)
5	EDO	B	908	-	3,3,3	0.44	0	2,2,2	0.38	0
3	SO4	A	909	-	4,4,4	0.26	0	6,6,6	0.12	0
3	SO4	A	907	-	4,4,4	0.24	0	6,6,6	0.16	0
3	SO4	A	908	-	4,4,4	0.32	0	6,6,6	0.38	0
3	SO4	A	906	-	4,4,4	0.32	0	6,6,6	0.76	0
3	SO4	A	902	-	4,4,4	0.22	0	6,6,6	0.54	0
3	SO4	B	904	-	4,4,4	0.30	0	6,6,6	0.27	0
5	EDO	A	912	-	3,3,3	0.32	0	2,2,2	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	907	-	-	1/1/1/1	-
4	MPO	B	906	-	-	2/7/15/15	0/1/1/1
2	BGC	B	901	-	-	0/2/22/22	0/1/1/1
2	BGC	A	901	-	-	0/2/22/22	0/1/1/1
5	EDO	A	911	-	-	0/1/1/1	-
5	EDO	A	912	-	-	1/1/1/1	-
5	EDO	B	908	-	-	1/1/1/1	-
4	MPO	A	910	-	-	1/7/15/15	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	910	MPO	C1-S1	-7.44	1.67	1.77
4	B	906	MPO	C1-S1	-7.34	1.67	1.77

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	910	MPO	C7-N1-C4	7.33	124.63	108.84
4	B	906	MPO	C7-N1-C4	6.09	121.96	108.84
4	A	910	MPO	O1-S1-C1	4.26	113.16	106.73
4	A	910	MPO	C3-N1-C4	3.59	120.80	111.24
4	B	906	MPO	C3-N1-C4	3.53	120.66	111.24
4	A	910	MPO	C6-C7-N1	3.48	115.41	110.12
4	B	906	MPO	O1-S1-C1	3.22	111.59	106.73
4	B	906	MPO	C3-C2-C1	-2.72	103.83	113.02
2	A	901	BGC	C3-C4-C5	-2.63	105.46	110.23
4	B	906	MPO	O2-S1-O1	-2.37	106.12	113.82
2	B	901	BGC	C3-C4-C5	-2.33	106.01	110.23
4	B	906	MPO	C3-N1-C7	2.30	117.38	111.24
4	A	910	MPO	O3-S1-C1	2.12	110.15	106.00
4	A	910	MPO	O2-S1-C1	-2.10	103.55	106.73

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	910	MPO	C2-C3-N1-C4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	B	906	MPO	C2-C3-N1-C4
5	B	908	EDO	O1-C1-C2-O2
5	A	912	EDO	O1-C1-C2-O2
5	B	907	EDO	O1-C1-C2-O2
4	B	906	MPO	C2-C3-N1-C7

There are no ring outliers.

5 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	907	EDO	4	0
4	B	906	MPO	1	0
3	B	902	SO4	1	0
2	B	901	BGC	1	0
2	A	901	BGC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	778/829 (93%)	0.07	33 (4%)	40 42	9, 17, 35, 59	6 (0%)
1	B	780/829 (94%)	0.16	21 (2%)	56 58	10, 19, 33, 47	1 (0%)
All	All	1558/1658 (93%)	0.11	54 (3%)	47 49	9, 18, 34, 59	7 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	797	ALA	5.7
1	B	828	TRP	5.1
1	A	536	ALA	4.7
1	B	536	ALA	4.2
1	B	797	ALA	3.8
1	A	470	ALA	3.6
1	A	539	LEU	3.6
1	A	451	GLY	3.4
1	B	296	PRO	3.3
1	A	296	PRO	3.2
1	A	820	ARG	3.2
1	A	798	THR	3.1
1	A	532	PHE	3.1
1	A	402	ASP	3.1
1	A	726	THR	3.0
1	B	49	TYR	3.0
1	B	348	ALA	3.0
1	A	461	HIS	2.7
1	A	537	THR	2.6
1	A	465	PRO	2.5
1	B	818	ASP	2.5
1	A	453	LEU	2.5
1	B	349	THR	2.5
1	A	468	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	466	GLY	2.4
1	A	534	MET	2.4
1	B	465	PRO	2.4
1	A	535	SER	2.4
1	A	531	GLY	2.4
1	A	533	ALA	2.4
1	A	796	ALA	2.4
1	B	601	ALA	2.3
1	B	575	ASP	2.3
1	B	553	ALA	2.3
1	B	468	ALA	2.3
1	A	521	THR	2.3
1	B	492	GLY	2.2
1	A	51	SER	2.2
1	A	297	GLY	2.2
1	B	819	LEU	2.2
1	B	796	ALA	2.2
1	A	455	PRO	2.2
1	A	469	GLY	2.1
1	A	281	ASP	2.1
1	A	457	PHE	2.1
1	A	299	VAL	2.1
1	A	466	GLY	2.1
1	B	471	LEU	2.1
1	A	467	LYS	2.1
1	A	528	THR	2.0
1	B	563	ARG	2.0
1	B	755	PRO	2.0
1	A	295	LEU	2.0
1	B	543	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	A	907	5/5	0.74	0.14	87,89,90,91	0
5	EDO	B	908	4/4	0.78	0.18	45,50,51,51	0
3	SO4	A	909	5/5	0.81	0.14	94,95,96,96	0
3	SO4	B	903	5/5	0.83	0.10	64,66,68,69	0
3	SO4	A	906	5/5	0.84	0.10	26,40,45,46	0
3	SO4	A	904	5/5	0.84	0.12	27,35,40,41	0
3	SO4	A	908	5/5	0.84	0.13	57,58,60,60	0
4	MPO	B	906	13/13	0.85	0.15	34,43,48,48	0
3	SO4	B	904	5/5	0.86	0.10	43,46,49,53	0
5	EDO	A	911	4/4	0.87	0.13	41,41,44,45	0
3	SO4	B	902	5/5	0.87	0.13	20,32,34,43	0
4	MPO	A	910	13/13	0.88	0.13	29,38,49,53	0
5	EDO	B	907	4/4	0.88	0.12	34,34,37,39	0
3	SO4	A	905	5/5	0.88	0.13	17,26,27,35	0
3	SO4	A	902	5/5	0.89	0.10	40,43,49,51	0
5	EDO	A	912	4/4	0.91	0.15	16,21,38,45	0
3	SO4	A	903	5/5	0.93	0.08	51,53,57,57	0
3	SO4	B	905	5/5	0.93	0.07	49,53,55,56	0
2	BGC	A	901	12/12	0.97	0.06	12,13,16,19	0
2	BGC	B	901	12/12	0.97	0.05	13,15,17,20	0

6.5 Other polymers ⓘ

There are no such residues in this entry.