



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:26 PM UTC

PDB ID : 2I4Q / pdb_00002i4q
Title : Human renin/PF02342674 complex
Authors : Holsworth, D.D.; Jalaie, M.; Zhang, E.; McConnell, P.; Mochalkin, I.; Finzel, B.C.
Deposited on : 2006-08-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	8% 92% 8%
1	B	336	10% 88% 11%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Renin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	336	Total	C	N	O	S	0	0	0
			2592	1654	420	504	14			
1	B	336	Total	C	N	O	S	0	0	0
			2592	1654	420	504	14			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ASN	-	cloning artifact	UNP P00797
A	1	THR	-	cloning artifact	UNP P00797
B	0	ASN	-	cloning artifact	UNP P00797
B	1	THR	-	cloning artifact	UNP P00797

- Molecule 2 is (2S)-6-(2,4-DIAMINO-6-ETHYLPYRIMIDIN-5-YL)-2-(3,5-DIFLUOROPHENYL)-4-(3-METHOXYPROPYL)-2-METHYL-2H-1,4-BENZOXAZIN-3(4H)-ONE (CCD ID: UA4) (formula: C₂₅H₂₇F₂N₅O₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	0	0
			35	25	2	5	3		
2	B	1	Total	C	F	N	O	0	0
			35	25	2	5	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	109	Total	O	0	0
			109	109		
3	B	81	Total	O	0	0
			81	81		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Renin



• Molecule 1: Renin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, α , β , γ	141.90Å 141.90Å 141.90Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.00 – 2.30 35.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	88.8 (35.00-2.30) 80.2 (35.00-2.30)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.41 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.229 , 0.261 0.221 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	39.7	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 25.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.039 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5444	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: UA4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2652	0.73	0/3598
1	B	0.46	0/2652	0.76	0/3598
All	All	0.46	0/5304	0.75	0/7196

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2592	0	2524	11	0
1	B	2592	0	2524	20	0
2	A	35	0	27	0	0
2	B	35	0	27	0	0
3	A	109	0	0	1	0
3	B	81	0	0	1	0
All	All	5444	0	5102	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:217:LEU:HB2	1:B:307:THR:HG22	1.60	0.83
1:B:197:GLN:NE2	1:B:307:THR:HG21	1.93	0.83
1:B:0:ASN:N	3:B:417:HOH:O	2.22	0.71
1:B:70:ASN:O	1:B:87:LEU:O	2.11	0.69
1:B:197:GLN:HE21	1:B:307:THR:HG21	1.65	0.61
1:A:0:ASN:N	3:A:445:HOH:O	2.36	0.58
1:B:133:GLY:O	1:B:134:ARG:HG3	2.06	0.55
1:B:193:THR:HG23	1:B:327:ASN:HD21	1.73	0.53
1:A:276:SER:O	1:A:280:VAL:HG13	2.12	0.50
1:A:155:PHE:CZ	1:A:171:ILE:HD12	2.49	0.47
1:B:217:LEU:HB2	1:B:307:THR:CG2	2.38	0.46
1:B:112:LEU:HA	1:B:113:PRO:HA	1.79	0.45
1:B:305:GLY:HA2	1:B:307:THR:HG23	1.99	0.45
1:B:280:VAL:HG22	1:B:293:LEU:HD22	1.98	0.45
1:B:193:THR:HG23	1:B:327:ASN:ND2	2.33	0.44
1:B:155:PHE:CZ	1:B:171:ILE:HD12	2.53	0.43
1:A:254:CYS:HB2	1:A:286:SER:O	2.19	0.43
1:B:217:LEU:HD12	1:B:307:THR:HG22	2.02	0.42
1:A:101:GLN:OE1	1:A:138:ILE:HA	2.18	0.42
1:B:9:ASN:OD1	1:B:9:ASN:C	2.63	0.41
1:B:225:SER:O	1:B:312:ALA:HB3	2.21	0.41
1:B:281:PHE:CZ	1:B:294:ALA:HB2	2.56	0.41
1:A:221:ASP:O	1:A:311:GLY:HA2	2.20	0.41
1:A:40:TRP:CD1	1:A:76:LEU:HD23	2.55	0.41
1:B:40:TRP:CE3	1:B:106:VAL:HG21	2.56	0.41
1:B:257:GLY:HA3	1:B:282:GLN:OE1	2.21	0.41
1:A:273:THR:O	1:A:317:LYS:NZ	2.50	0.40
1:A:87:LEU:HD13	1:A:102:MET:HE3	2.04	0.40
1:A:155:PHE:CE1	1:A:171:ILE:HD12	2.56	0.40
1:B:74:LEU:HG	1:B:87:LEU:HD13	2.02	0.40
1:A:43:SER:HB2	1:A:105:GLU:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/336 (99%)	327 (98%)	7 (2%)	0	100	100
1	B	334/336 (99%)	323 (97%)	11 (3%)	0	100	100
All	All	668/672 (99%)	650 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/287 (100%)	276 (96%)	11 (4%)	29	44
1	B	287/287 (100%)	268 (93%)	19 (7%)	15	21
All	All	574/574 (100%)	544 (95%)	30 (5%)	21	31

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LEU
1	A	84	SER
1	A	163	ASN
1	A	195	VAL
1	A	210	LEU
1	A	217	LEU
1	A	237	LEU
1	A	246	ARG
1	A	270	LYS
1	A	280	VAL
1	A	333	LEU
1	B	2	THR
1	B	51	THR
1	B	74	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	87	LEU
1	B	112	LEU
1	B	134	ARG
1	B	149	LYS
1	B	162	GLU
1	B	164	SER
1	B	179	GLN
1	B	219	LEU
1	B	241	LEU
1	B	246	ARG
1	B	256	GLU
1	B	274	LEU
1	B	288	LYS
1	B	290	LEU
1	B	325	ARG
1	B	335	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	0	ASN
1	A	89	GLN
1	A	163	ASN
1	A	170	GLN
1	A	327	ASN
1	B	130	GLN
1	B	141	ASN
1	B	165	GLN
1	B	189	ASN
1	B	197	GLN
1	B	327	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UA4	A	336	-	37,38,38	3.43	14 (37%)	49,56,56	2.58	18 (36%)
2	UA4	B	336	-	37,38,38	3.52	13 (35%)	49,56,56	2.69	18 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UA4	A	336	-	-	3/17/36/36	0/4/4/4
2	UA4	B	336	-	-	3/17/36/36	0/4/4/4

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	336	UA4	C12-C13	-10.83	1.21	1.38
2	A	336	UA4	C12-C13	-10.52	1.21	1.38
2	B	336	UA4	C28-C27	-7.91	1.27	1.39
2	A	336	UA4	C28-C27	-7.16	1.28	1.39
2	B	336	UA4	C12-C11	-6.24	1.27	1.39
2	A	336	UA4	C12-C11	-6.19	1.27	1.39
2	B	336	UA4	C28-C29	-6.03	1.27	1.37
2	B	336	UA4	C32-C27	5.99	1.48	1.39
2	A	336	UA4	C32-C27	5.93	1.48	1.39
2	B	336	UA4	C30-C31	5.87	1.47	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	336	UA4	C30-C31	5.80	1.47	1.37
2	A	336	UA4	O17-C14	5.71	1.46	1.37
2	B	336	UA4	C13-C14	-5.30	1.28	1.39
2	A	336	UA4	C13-C14	-5.16	1.28	1.39
2	A	336	UA4	C28-C29	-5.05	1.29	1.37
2	B	336	UA4	O17-C14	4.57	1.44	1.37
2	B	336	UA4	C16-C11	3.93	1.46	1.39
2	A	336	UA4	C16-C11	3.88	1.46	1.39
2	B	336	UA4	C16-C15	-3.81	1.33	1.39
2	A	336	UA4	C15-C14	-3.62	1.34	1.40
2	B	336	UA4	C15-C14	-3.59	1.34	1.40
2	A	336	UA4	C16-C15	-3.46	1.34	1.39
2	A	336	UA4	C2-C3	3.36	1.50	1.43
2	B	336	UA4	C2-C3	3.10	1.49	1.43
2	A	336	UA4	C15-N20	-2.94	1.36	1.41
2	B	336	UA4	C15-N20	-2.91	1.36	1.41
2	A	336	UA4	C32-C31	-2.01	1.34	1.37

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	336	UA4	C5-N6-C1	8.63	122.55	116.26
2	A	336	UA4	C5-N6-C1	7.99	122.09	116.26
2	B	336	UA4	C2-C3-N4	-6.93	118.53	122.55
2	A	336	UA4	C2-C3-N4	-6.24	118.93	122.55
2	A	336	UA4	C27-C28-C29	5.90	124.69	118.35
2	B	336	UA4	C27-C28-C29	5.38	124.13	118.35
2	B	336	UA4	C30-C31-C32	-5.17	117.21	123.50
2	A	336	UA4	C30-C31-C32	-4.68	117.82	123.50
2	B	336	UA4	C16-C11-C2	-4.66	112.68	120.55
2	B	336	UA4	C14-C15-N20	4.62	124.09	118.58
2	A	336	UA4	C12-C11-C2	4.45	128.43	120.80
2	A	336	UA4	C16-C11-C2	-4.44	113.06	120.55
2	B	336	UA4	C12-C11-C2	4.35	128.25	120.80
2	B	336	UA4	C2-C1-N6	-4.03	118.86	123.47
2	A	336	UA4	C14-C15-N20	3.97	123.31	118.58
2	A	336	UA4	C2-C1-N6	-3.62	119.32	123.47
2	B	336	UA4	C15-N20-C19	-2.96	119.99	122.80
2	B	336	UA4	C3-C2-C1	2.84	118.01	115.97
2	A	336	UA4	C12-C13-C14	2.81	125.38	120.06
2	A	336	UA4	O17-C14-C13	2.79	120.82	116.60
2	B	336	UA4	C5-N4-C3	2.77	120.60	117.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	336	UA4	C16-C15-C14	-2.70	115.80	119.78
2	A	336	UA4	F34-C31-C32	2.69	122.10	118.28
2	B	336	UA4	N6-C5-N4	-2.66	121.41	125.48
2	A	336	UA4	C5-N4-C3	2.63	120.44	117.28
2	B	336	UA4	C12-C13-C14	2.58	124.95	120.06
2	A	336	UA4	F33-C29-C28	2.58	121.94	118.28
2	A	336	UA4	N6-C5-N4	-2.44	121.75	125.48
2	A	336	UA4	C16-C15-C14	-2.42	116.22	119.78
2	B	336	UA4	F34-C31-C32	2.35	121.61	118.28
2	B	336	UA4	C23-C22-N20	-2.18	107.72	112.45
2	A	336	UA4	C15-N20-C19	-2.12	120.79	122.80
2	A	336	UA4	C23-C22-N20	-2.12	107.85	112.45
2	A	336	UA4	C3-C2-C1	2.06	117.45	115.97
2	B	336	UA4	F34-C31-C30	2.04	121.17	118.28
2	B	336	UA4	O17-C14-C13	2.02	119.66	116.60

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	336	UA4	C35-C18-C27-C28
2	A	336	UA4	C35-C18-C27-C32
2	B	336	UA4	C35-C18-C27-C32
2	B	336	UA4	C22-C23-C24-O25
2	A	336	UA4	C22-C23-C24-O25
2	B	336	UA4	C35-C18-C27-C28

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	336/336 (100%)	0.60	28 (8%)	17 19	27, 41, 65, 77	0
1	B	336/336 (100%)	0.75	34 (10%)	12 14	29, 46, 64, 74	0
All	All	672/672 (100%)	0.67	62 (9%)	14 16	27, 44, 64, 77	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	209	THR	5.3
1	A	248	PHE	4.9
1	B	210	LEU	4.8
1	A	247	LEU	4.5
1	B	247	LEU	4.3
1	A	210	LEU	4.1
1	A	208	SER	4.1
1	B	208	SER	4.0
1	A	165	GLN	3.7
1	B	52	ALA	3.6
1	A	207	SER	3.6
1	A	240	ALA	3.4
1	A	286	SER	3.4
1	B	163	ASN	3.4
1	A	296	HIS	3.4
1	B	209	THR	3.4
1	A	146	GLY	3.3
1	B	51	THR	3.3
1	A	269	GLY	3.1
1	A	162	GLU	2.9
1	A	214	ASP	2.9
1	B	55	TYR	2.8
1	A	164	SER	2.8
1	A	285	TYR	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	134	ARG	2.8
1	B	213	GLU	2.7
1	B	50	TYR	2.7
1	B	165	GLN	2.7
1	A	205	VAL	2.6
1	B	193	THR	2.6
1	A	243	ALA	2.5
1	B	115	MET	2.5
1	B	162	GLU	2.5
1	B	164	SER	2.4
1	B	207	SER	2.4
1	A	239	GLU	2.4
1	B	71	GLY	2.4
1	B	150	GLU	2.4
1	B	304	THR	2.4
1	B	0	ASN	2.4
1	B	335	ARG	2.4
1	B	305	GLY	2.4
1	B	49	LEU	2.4
1	A	245	LYS	2.3
1	A	217	LEU	2.2
1	B	270	LYS	2.2
1	B	70	ASN	2.2
1	B	96	GLY	2.2
1	B	24	PRO	2.2
1	B	298	MET	2.1
1	B	179	GLN	2.1
1	B	74	LEU	2.1
1	A	179	GLN	2.1
1	A	211	LEU	2.1
1	A	226	TYR	2.1
1	B	1	THR	2.1
1	A	270	LYS	2.1
1	A	160	ASP	2.1
1	A	303	PRO	2.1
1	A	246	ARG	2.0
1	B	205	VAL	2.0
1	B	102	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UA4	A	336	35/35	0.92	0.09	31,33,39,40	0
2	UA4	B	336	35/35	0.93	0.09	34,37,42,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around UA4 B 336:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.