



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 09:15 PM UTC

PDB ID : 4I4J / pdb_00004i4j
Title : The structure of SgcE10, the ACP-polyene thioesterase involved in C-1027 biosynthesis
Authors : Kim, Y.; Bigelow, L.; Bearden, J.; Babnigg, J.; Bingman, C.A.; Yennamalli, R.; Lohman, J.; Ma, M.; Shen, B.; Phillips Jr., G.N.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG); Enzyme Discovery for Natural Product Biosynthesis (NatPro)
Deposited on : 2012-11-27
Resolution : 2.78 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5248 (2.80-2.76)
Clashscore	190562	5693 (2.80-2.76)
Ramachandran outliers	187476	5590 (2.80-2.76)
Sidechain outliers	187428	5592 (2.80-2.76)
RSRZ outliers	180081	5251 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	159	72%16%•12%
1	B	159	% 66%21%•12%
1	C	159	79%9%•12%
1	D	159	67%18%•14%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	159	3% 67% 19% • 13%
1	F	159	% 69% 21% 11%
1	G	159	75% 13% • 11%
1	H	159	77% 11% • 11%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9254 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACP-polyene thioesterase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	140	Total	C	N	O	S	Se	0	2	0
			1155	736	197	217	3	2			
1	B	140	Total	C	N	O	S	Se	0	0	0
			1135	725	192	213	3	2			
1	C	140	Total	C	N	O	S	Se	0	1	0
			1146	731	196	214	3	2			
1	D	136	Total	C	N	O	S	Se	0	0	0
			1109	710	187	207	3	2			
1	E	138	Total	C	N	O	S	Se	0	0	0
			1120	717	189	209	3	2			
1	F	142	Total	C	N	O	S	Se	0	0	0
			1154	735	197	217	3	2			
1	G	142	Total	C	N	O	S	Se	0	0	0
			1151	733	195	218	3	2			
1	H	141	Total	C	N	O	S	Se	0	0	0
			1143	729	194	215	3	2			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q8GME8
A	-1	ASN	-	expression tag	UNP Q8GME8
A	0	ALA	-	expression tag	UNP Q8GME8
B	-2	SER	-	expression tag	UNP Q8GME8
B	-1	ASN	-	expression tag	UNP Q8GME8
B	0	ALA	-	expression tag	UNP Q8GME8
C	-2	SER	-	expression tag	UNP Q8GME8
C	-1	ASN	-	expression tag	UNP Q8GME8
C	0	ALA	-	expression tag	UNP Q8GME8
D	-2	SER	-	expression tag	UNP Q8GME8
D	-1	ASN	-	expression tag	UNP Q8GME8
D	0	ALA	-	expression tag	UNP Q8GME8
E	-2	SER	-	expression tag	UNP Q8GME8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	ASN	-	expression tag	UNP Q8GME8
E	0	ALA	-	expression tag	UNP Q8GME8
F	-2	SER	-	expression tag	UNP Q8GME8
F	-1	ASN	-	expression tag	UNP Q8GME8
F	0	ALA	-	expression tag	UNP Q8GME8
G	-2	SER	-	expression tag	UNP Q8GME8
G	-1	ASN	-	expression tag	UNP Q8GME8
G	0	ALA	-	expression tag	UNP Q8GME8
H	-2	SER	-	expression tag	UNP Q8GME8
H	-1	ASN	-	expression tag	UNP Q8GME8
H	0	ALA	-	expression tag	UNP Q8GME8

- Molecule 2 is D(-)-TARTARIC ACID (CCD ID: TAR) (formula: $C_4H_6O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			10	4	6		
2	B	1	Total	C	O	0	0
			10	4	6		
2	C	1	Total	C	O	0	0
			10	4	6		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	F	1	Total	C	O	0	0
			6	3	3		
4	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	6	Total 6	O 6	0	0
5	B	19	Total 19	O 19	0	0
5	C	12	Total 12	O 12	0	0
5	D	18	Total 18	O 18	0	0
5	E	6	Total 6	O 6	0	0
5	F	7	Total 7	O 7	0	0
5	G	10	Total 10	O 10	0	0
5	H	17	Total 17	O 17	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ACP-polyene thioesterase

Chain A: 



- Molecule 1: ACP-polyene thioesterase

Chain B: 




- Molecule 1: ACP-polyene thioesterase

Chain C: 



- Molecule 1: ACP-polyene thioesterase

Chain D: 



- Molecule 1: ACP-polyene thioesterase

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	57.42Å 184.53Å 59.40Å 90.00° 95.60° 90.00°	Depositor
Resolution (Å)	39.16 – 2.78 39.16 – 2.78	Depositor EDS
% Data completeness (in resolution range)	95.5 (39.16-2.78) 95.3 (39.16-2.78)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.77Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1161)	Depositor
R, R_{free}	0.211 , 0.270 0.214 , 0.270	Depositor DCC
R_{free} test set	2015 reflections (6.70%)	wwPDB-VP
Wilson B-factor (Å ²)	69.1	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 15.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.035 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9254	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, TAR, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/1176	0.69	0/1587
1	B	0.26	0/1156	0.69	0/1561
1	C	0.26	0/1167	0.68	0/1575
1	D	0.25	0/1128	0.67	0/1520
1	E	0.27	0/1140	0.69	0/1537
1	F	0.27	0/1175	0.70	0/1586
1	G	0.25	0/1172	0.68	0/1584
1	H	0.26	0/1164	0.74	2/1573 (0.1%)
All	All	0.26	0/9278	0.69	2/12523 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	142	ALA	CA-C-N	-6.09	115.50	121.65
1	H	142	ALA	C-N-CA	-6.09	115.50	121.65

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1155	0	1137	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1135	0	1120	19	0
1	C	1146	0	1132	9	0
1	D	1109	0	1096	18	0
1	E	1120	0	1106	16	0
1	F	1154	0	1137	13	0
1	G	1151	0	1129	11	0
1	H	1143	0	1125	8	0
2	A	10	0	4	1	0
2	B	10	0	4	0	0
2	C	10	0	4	0	0
3	C	4	0	6	0	0
4	F	12	0	16	2	0
5	A	6	0	0	0	0
5	B	19	0	0	0	0
5	C	12	0	0	0	0
5	D	18	0	0	2	0
5	E	6	0	0	0	0
5	F	7	0	0	0	0
5	G	10	0	0	0	0
5	H	17	0	0	2	0
All	All	9254	0	9016	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:122:ARG:HB2	1:F:131:THR:HB	1.71	0.72
1:D:46:ARG:NH1	1:D:144:TYR:OH	2.28	0.66
1:A:32:LEU:HD13	1:B:32:LEU:HD13	1.81	0.63
1:B:84:ARG:NH2	1:B:98:ASP:OD2	2.33	0.61
1:A:34:TRP:HB3	1:A:81:ILE:HD13	1.82	0.61
1:C:74:THR:OG1	1:C:75:ALA:N	2.34	0.60
1:C:88:LEU:HD12	1:C:93:LEU:HB3	1.82	0.60
1:H:59:LYS:HD2	1:H:128:THR:HG21	1.84	0.59
1:E:54:VAL:HG23	1:E:59:LYS:HA	1.85	0.59
1:H:25:ASN:ND2	5:H:205:HOH:O	2.36	0.58
1:F:46:ARG:NH1	1:F:144:TYR:OH	2.36	0.58
1:B:93:LEU:HD23	1:B:133:ILE:HD13	1.86	0.57
1:G:91:THR:HG23	1:G:92:GLN:HG3	1.86	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:7:ASP:HB3	1:B:84:ARG:HH11	1.71	0.55
1:E:91:THR:HG23	1:E:92:GLN:HG3	1.89	0.55
1:E:80:SER:HB3	1:E:100:ILE:HB	1.89	0.55
1:F:63:LEU:HD11	1:F:119:ALA:HB2	1.88	0.54
1:H:86:SER:HB3	1:H:94:GLU:HG3	1.89	0.54
1:G:52:ALA:HA	1:G:55:GLN:HE21	1.71	0.54
1:H:11:LEU:HD13	1:H:41:LEU:HD22	1.90	0.54
1:A:85:LEU:HB3	1:A:145:SER:HA	1.89	0.54
1:F:82:ARG:HD2	1:F:100:ILE:HD13	1.89	0.54
1:A:84:ARG:NH2	1:A:98:ASP:OD2	2.34	0.53
1:G:85:LEU:HD22	1:G:141:LEU:HB3	1.90	0.53
1:A:39:ARG:NH1	1:A:40[B]:GLU:OE1	2.42	0.53
1:G:50:VAL:HG11	1:G:137:LEU:HB2	1.90	0.52
1:E:63:LEU:HB2	1:E:117:ARG:HG2	1.91	0.51
1:G:80:SER:HB3	1:G:100:ILE:HB	1.92	0.51
1:F:37:ARG:HH11	4:F:201:GOL:H32	1.76	0.51
1:D:74:THR:OG1	1:D:75:ALA:N	2.41	0.51
1:E:121:MSE:HB3	1:E:128:THR:HB	1.94	0.50
1:H:12:ARG:NH2	5:H:204:HOH:O	2.42	0.50
1:H:124:PRO:HG2	1:H:127:ALA:HB3	1.93	0.50
1:F:93:LEU:HD23	1:F:133:ILE:HD13	1.94	0.50
1:A:122:ARG:HB3	1:A:129:VAL:HG12	1.93	0.50
1:E:21:ASN:HD21	1:E:25:ASN:HB2	1.76	0.50
1:A:85:LEU:HD22	1:A:141:LEU:HB3	1.94	0.49
1:D:121:MSE:HB3	1:D:128:THR:HB	1.94	0.49
1:E:11:LEU:HD23	1:E:81:ILE:HD12	1.95	0.48
1:F:87:GLU:HB3	1:F:94:GLU:HB3	1.96	0.48
1:C:22:LEU:H	1:C:22:LEU:HD23	1.79	0.48
1:A:35:GLN:HG3	1:A:99:TYR:OH	2.14	0.48
1:D:54:VAL:HA	1:D:58:LEU:HB3	1.95	0.47
1:B:71:ALA:HB3	1:B:111:VAL:HG23	1.95	0.47
1:F:74:THR:OG1	1:F:75:ALA:N	2.47	0.47
1:D:82:ARG:HE	1:D:100:ILE:HG13	1.79	0.47
1:A:61:PHE:CE1	1:B:72:GLU:HG3	2.49	0.47
1:C:83:MSE:HE2	1:C:95:PHE:CG	2.49	0.46
1:B:73:ILE:HG23	1:B:111:VAL:HG21	1.96	0.46
1:F:123:GLY:HA3	1:F:127:ALA:HB3	1.97	0.46
1:E:122:ARG:HG2	1:E:131:THR:OG1	2.16	0.46
1:B:74:THR:HG22	1:B:75:ALA:H	1.80	0.46
1:D:104:ASP:OD1	1:D:104:ASP:N	2.49	0.46
1:H:95:PHE:HE1	1:H:141:LEU:HD21	1.81	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:41:LEU:HD12	1:C:44:LYS:HD3	1.98	0.45
1:D:46:ARG:HB3	1:D:140:ALA:HB1	1.97	0.45
1:G:73:ILE:HG22	1:G:101:LYS:HE2	1.98	0.45
1:C:68:GLU:HG2	1:D:64:LYS:HD2	1.98	0.45
1:E:74:THR:OG1	1:E:75:ALA:N	2.49	0.45
1:B:80:SER:HB3	1:B:100:ILE:HB	1.98	0.45
1:E:29:VAL:HG12	1:E:33:ARG:HD2	1.99	0.45
1:A:7:ASP:HB2	1:A:84:ARG:HH11	1.82	0.44
1:D:133:ILE:HG23	1:D:137:LEU:HD23	1.98	0.44
1:E:30:ASN:HA	1:E:33:ARG:HD3	1.99	0.44
1:G:5:ASN:HA	1:G:6:PRO:HA	1.77	0.44
1:A:122:ARG:N	1:A:129:VAL:O	2.50	0.44
1:B:57:ASP:OD1	1:B:122:ARG:NH1	2.48	0.44
1:D:35:GLN:HG3	1:D:99:TYR:OH	2.18	0.44
1:A:74:THR:OG1	1:A:75:ALA:N	2.51	0.44
1:D:82:ARG:H	1:D:82:ARG:HG2	1.65	0.43
1:A:61:PHE:HE1	1:B:72:GLU:HG3	1.83	0.43
1:G:87:GLU:OE2	1:G:117:ARG:NH2	2.52	0.43
1:F:57:ASP:N	1:F:57:ASP:OD1	2.53	0.42
2:A:201:TAR:O41	2:A:201:TAR:O2	2.36	0.42
1:D:42:PHE:CG	1:D:83:MSE:HE1	2.54	0.42
1:E:40:GLU:O	1:E:44:LYS:HG2	2.20	0.42
1:D:8:TYR:CG	1:D:82:ARG:HB3	2.54	0.42
1:A:59:LYS:HE3	1:A:59:LYS:HB2	1.87	0.42
1:D:8:TYR:CD1	1:D:82:ARG:HB3	2.54	0.42
1:D:55:GLN:HG2	5:D:204:HOH:O	2.20	0.42
1:E:39:ARG:HH11	1:E:116:GLN:HE22	1.68	0.42
1:D:63:LEU:HD11	1:D:119:ALA:HB2	2.01	0.42
1:E:11:LEU:HB3	1:E:81:ILE:HB	2.00	0.42
1:B:66:ASP:OD1	1:B:115:ARG:HB2	2.20	0.41
1:C:39:ARG:HG3	1:C:83:MSE:HE1	2.01	0.41
1:H:74:THR:HG22	1:H:75:ALA:H	1.85	0.41
1:E:20:THR:O	1:G:21:ASN:HA	2.20	0.41
1:F:59:LYS:HB3	1:F:121:MSE:HB2	2.02	0.41
1:B:8:TYR:CE1	1:B:82:ARG:HD2	2.55	0.41
1:F:28:TYR:HB2	1:F:69:PHE:CZ	2.56	0.41
1:B:61:PHE:CD2	1:B:121:MSE:HG3	2.55	0.41
1:B:76:PHE:CZ	1:C:51:LEU:HB3	2.56	0.41
1:B:87:GLU:HB3	1:B:94:GLU:HB3	2.02	0.41
1:C:29:VAL:O	1:C:33:ARG:HG3	2.20	0.41
1:F:35:GLN:HG3	1:F:99:TYR:OH	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:THR:OG1	1:B:78:GLU:OE1	2.36	0.40
1:D:36:GLY:HA2	1:D:39:ARG:NH1	2.36	0.40
1:D:56:GLU:HG2	5:D:204:HOH:O	2.21	0.40
1:A:23:VAL:HG21	1:B:59:LYS:HD2	2.03	0.40
4:F:202:GOL:H11	1:G:37:ARG:HH21	1.87	0.40
1:B:42:PHE:CG	1:B:83:MSE:HE1	2.56	0.40
1:E:85:LEU:HD11	1:E:93:LEU:HD22	2.04	0.40
1:G:121:MSE:HB3	1:G:128:THR:OG1	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	140/159 (88%)	134 (96%)	6 (4%)	0	100	100
1	B	138/159 (87%)	133 (96%)	5 (4%)	0	100	100
1	C	139/159 (87%)	131 (94%)	8 (6%)	0	100	100
1	D	132/159 (83%)	126 (96%)	5 (4%)	1 (1%)	16	40
1	E	134/159 (84%)	130 (97%)	4 (3%)	0	100	100
1	F	140/159 (88%)	130 (93%)	9 (6%)	1 (1%)	18	44
1	G	140/159 (88%)	136 (97%)	4 (3%)	0	100	100
1	H	139/159 (87%)	132 (95%)	6 (4%)	1 (1%)	18	44
All	All	1102/1272 (87%)	1052 (96%)	47 (4%)	3 (0%)	36	63

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	124	PRO
1	H	124	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	130	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	123/129 (95%)	116 (94%)	7 (6%)	18	46
1	B	121/129 (94%)	114 (94%)	7 (6%)	18	45
1	C	122/129 (95%)	120 (98%)	2 (2%)	55	81
1	D	118/129 (92%)	115 (98%)	3 (2%)	42	73
1	E	119/129 (92%)	113 (95%)	6 (5%)	22	51
1	F	123/129 (95%)	115 (94%)	8 (6%)	15	40
1	G	123/129 (95%)	120 (98%)	3 (2%)	43	74
1	H	122/129 (95%)	118 (97%)	4 (3%)	33	66
All	All	971/1032 (94%)	931 (96%)	40 (4%)	27	59

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	LEU
1	A	100	ILE
1	A	109	THR
1	A	113	ARG
1	A	125	ASN
1	A	126	THR
1	A	129	VAL
1	B	39	ARG
1	B	74	THR
1	B	82	ARG
1	B	96	THR
1	B	105	ASP
1	B	125	ASN
1	B	128	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	74	THR
1	C	133	ILE
1	D	82	ARG
1	D	90	GLN
1	D	139	GLU
1	E	40	GLU
1	E	55	GLN
1	E	58	LEU
1	E	65	VAL
1	E	131	THR
1	E	137	LEU
1	F	14	THR
1	F	25	ASN
1	F	90	GLN
1	F	91	THR
1	F	104	ASP
1	F	117	ARG
1	F	126	THR
1	F	129	VAL
1	G	50	VAL
1	G	74	THR
1	G	90	GLN
1	H	32	LEU
1	H	54	VAL
1	H	93	LEU
1	H	126	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	125	ASN
1	B	25	ASN
1	B	35	GLN
1	B	90	GLN
1	C	13	HIS
1	C	116	GLN
1	C	125	ASN
1	D	21	ASN
1	D	25	ASN
1	D	55	GLN
1	D	90	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	116	GLN
1	E	35	GLN
1	E	90	GLN
1	E	116	GLN
1	F	21	ASN
1	F	25	ASN
1	F	55	GLN
1	F	90	GLN
1	G	35	GLN
1	G	55	GLN
1	G	90	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	F	202	-	5,5,5	0.36	0	5,5,5	0.30	0
2	TAR	C	201	-	9,9,9	1.25	0	12,12,12	0.92	0
2	TAR	A	201	-	9,9,9	1.24	0	12,12,12	0.94	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	F	201	-	5,5,5	0.37	0	5,5,5	0.31	0
2	TAR	B	201	-	9,9,9	1.25	0	12,12,12	0.98	0
3	EDO	C	202	-	3,3,3	0.43	0	2,2,2	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	F	202	-	-	2/4/4/4	-
2	TAR	C	201	-	-	8/12/12/12	-
2	TAR	A	201	-	-	6/12/12/12	-
4	GOL	F	201	-	-	2/4/4/4	-
2	TAR	B	201	-	-	10/12/12/12	-
3	EDO	C	202	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (28) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	TAR	O3-C3-C4-O4
2	A	201	TAR	O3-C3-C4-O41
2	B	201	TAR	O3-C3-C4-O4
2	B	201	TAR	O3-C3-C4-O41
4	F	201	GOL	O1-C1-C2-C3
4	F	202	GOL	O1-C1-C2-C3
2	C	201	TAR	C1-C2-C3-C4
2	A	201	TAR	O11-C1-C2-C3
2	B	201	TAR	O1-C1-C2-C3
2	B	201	TAR	O11-C1-C2-C3
2	C	201	TAR	O2-C2-C3-O3
2	C	201	TAR	C1-C2-C3-O3
2	C	201	TAR	O1-C1-C2-O2
2	C	201	TAR	O11-C1-C2-O2
2	C	201	TAR	O3-C3-C4-O4
2	C	201	TAR	O3-C3-C4-O41
2	A	201	TAR	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	201	TAR	O2-C2-C3-C4
2	B	201	TAR	O2-C2-C3-O3
2	B	201	TAR	O1-C1-C2-O2
2	B	201	TAR	O11-C1-C2-O2
2	B	201	TAR	C1-C2-C3-O3
2	B	201	TAR	O2-C2-C3-C4
2	A	201	TAR	O1-C1-C2-O2
2	A	201	TAR	O11-C1-C2-O2
4	F	202	GOL	O1-C1-C2-O2
4	F	201	GOL	O1-C1-C2-O2
2	B	201	TAR	C1-C2-C3-C4

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	F	202	GOL	1	0
2	A	201	TAR	1	0
4	F	201	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	138/159 (86%)	-0.47	0 100 100	12, 32, 67, 77	2 (1%)
1	B	138/159 (86%)	-0.56	1 (0%) 84 79	13, 30, 59, 68	0
1	C	138/159 (86%)	-0.57	0 100 100	7, 25, 55, 78	1 (0%)
1	D	134/159 (84%)	-0.54	0 100 100	7, 34, 67, 84	0
1	E	136/159 (85%)	-0.18	4 (2%) 53 46	16, 45, 86, 96	0
1	F	140/159 (88%)	-0.40	2 (1%) 73 67	16, 39, 74, 84	0
1	G	140/159 (88%)	-0.57	0 100 100	15, 31, 59, 79	0
1	H	139/159 (87%)	-0.58	0 100 100	17, 31, 59, 76	0
All	All	1103/1272 (86%)	-0.48	7 (0%) 85 81	7, 32, 70, 96	3 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	93	LEU	3.0
1	F	128	THR	2.5
1	F	127	ALA	2.2
1	E	133	ILE	2.2
1	B	6	PRO	2.2
1	E	145	SER	2.2
1	E	124	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	GOL	F	202	6/6	0.66	0.20	50,51,51,52	0
2	TAR	B	201	10/10	0.73	0.11	58,59,61,62	0
3	EDO	C	202	4/4	0.76	0.13	36,37,37,37	0
2	TAR	A	201	10/10	0.78	0.11	69,70,71,72	0
2	TAR	C	201	10/10	0.80	0.12	53,55,57,58	0
4	GOL	F	201	6/6	0.82	0.13	46,48,49,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.