



Full wwPDB EM Validation Report ⓘ

Mar 9, 2026 – 12:10 PM UTC

PDB ID : 6I52 / pdb_00006i52
EMDB ID : EMD-4410
Title : Yeast RPA bound to ssDNA
Authors : Yates, L.A.; Aramayo, R.J.; Zhang, X.
Deposited on : 2018-11-12
Resolution : 4.70 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 4.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	1989 (4.20 - 5.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	
2	B	132	
3	C	178	
4	D	20	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replication factor A protein 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	122	Total	C	N	O	S	0	0
			966	610	160	190	6		

- Molecule 2 is a protein called Replication factor A protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	132	Total	C	N	O	S	0	0
			1041	661	181	195	4		

There are 19 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	ASN	deletion	UNP P26754
B	?	-	ASP	deletion	UNP P26754
B	?	-	LEU	deletion	UNP P26754
B	?	-	ALA	deletion	UNP P26754
B	?	-	ALA	deletion	UNP P26754
B	?	-	GLY	deletion	UNP P26754
B	?	-	ASN	deletion	UNP P26754
B	?	-	ASP	deletion	UNP P26754
B	?	-	ASP	deletion	UNP P26754
B	?	-	SER	deletion	UNP P26754
B	?	-	SER	deletion	UNP P26754
B	?	-	GLY	deletion	UNP P26754
B	?	-	LYS	deletion	UNP P26754
B	?	-	GLY	deletion	UNP P26754
B	?	-	TYR	deletion	UNP P26754
B	?	-	GLY	deletion	UNP P26754
B	?	-	SER	deletion	UNP P26754
B	?	-	GLN	deletion	UNP P26754
B	?	-	VAL	deletion	UNP P26754

- Molecule 3 is a protein called Replication factor A protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	178	Total	C	N	O	S	0	0
			1449	906	254	283	6		

- Molecule 4 is a DNA chain called DNA (5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	20	Total 400	C 200	N 40	O 140	P 20	0	0



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	340864	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	2.1	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.162	Depositor
Minimum map value	-0.123	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.012	Depositor
Recommended contour level	0.045	Depositor
Map size (Å)	127.2, 127.2, 127.2	wwPDB
Map dimensions	120, 120, 120	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.06, 1.06, 1.06	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/981	0.28	0/1329
2	B	0.12	0/1060	0.33	0/1431
3	C	0.12	0/1474	0.39	0/1993
4	D	0.17	0/439	0.48	0/676
All	All	0.12	0/3954	0.36	0/5429

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	966	0	971	29	0
2	B	1041	0	1037	30	0
3	C	1449	0	1409	17	0
4	D	400	0	241	5	0
All	All	3856	0	3658	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:71:PHE:O	2:B:136:VAL:HB	1.59	0.99
1:A:22:ARG:HA	1:A:75:PHE:O	1.67	0.92
2:B:41:PRO:HA	2:B:70:CYS:O	1.71	0.89
1:A:42:THR:HG22	1:A:43:ILE:HG12	1.71	0.70
1:A:108:VAL:O	1:A:111:LEU:HB3	1.96	0.66
3:C:519:TYR:HB2	3:C:538:ASP:HA	1.78	0.64
1:A:72:TRP:HB2	1:A:95:CYS:O	1.96	0.64
1:A:74:GLU:HB3	1:A:93:VAL:HB	1.79	0.63
1:A:42:THR:HG23	1:A:109:VAL:HG11	1.80	0.62
1:A:114:LEU:HD23	1:A:117:LYS:HZ3	1.64	0.62
1:A:41:PRO:HA	1:A:51:VAL:HA	1.81	0.62
3:C:500:ASP:OD1	3:C:501:GLY:N	2.33	0.62
2:B:50:SER:HB2	2:B:61:SER:HB2	1.82	0.61
2:B:68:HIS:ND1	2:B:138:GLY:O	2.33	0.61
2:B:79:THR:HB	2:B:86:PHE:HB2	1.83	0.61
2:B:80:ASP:HB2	2:B:85:ILE:HA	1.84	0.59
1:A:18:ALA:HA	1:A:79:ASN:HB3	1.85	0.58
2:B:163:VAL:O	2:B:166:HIS:ND1	2.36	0.58
1:A:76:VAL:HB	1:A:91:ASP:HB3	1.86	0.57
2:B:78:ILE:HA	2:B:87:LEU:HD12	1.85	0.57
2:B:134:VAL:HG12	2:B:156:PRO:HA	1.86	0.57
3:C:472:ALA:HA	3:C:574:GLU:HA	1.87	0.56
2:B:168:LEU:O	2:B:172:LYS:CB	2.54	0.56
1:A:27:ILE:HG21	1:A:67:PHE:HB2	1.90	0.54
1:A:26:GLN:HB3	1:A:39:GLN:HB2	1.89	0.54
2:B:134:VAL:HA	2:B:155:LYS:O	2.08	0.54
2:B:99:ARG:NH1	4:D:25:DT:OP1	2.41	0.54
1:A:38:LEU:O	1:A:53:MET:HA	2.08	0.53
1:A:121:ILE:HG13	2:B:171:ILE:HG12	1.91	0.53
1:A:56:LEU:HD22	1:A:87:PHE:H	1.73	0.53
2:B:77:ASN:HB2	2:B:88:THR:HB	1.91	0.52
1:A:39:GLN:HA	1:A:53:MET:HG2	1.92	0.51
2:B:172:LYS:HE3	3:C:611:LEU:HB3	1.92	0.51
3:C:538:ASP:O	3:C:542:LYS:HB2	2.11	0.51
3:C:458:LEU:HB2	3:C:459:GLY:HA2	1.94	0.50
3:C:492:ASN:HA	4:D:13:DT:H5'	1.93	0.50
2:B:168:LEU:O	2:B:172:LYS:HB2	2.13	0.49
1:A:75:PHE:HZ	1:A:89:ILE:HG12	1.76	0.49
2:B:41:PRO:HB2	2:B:166:HIS:CD2	2.47	0.48
2:B:173:CYS:O	2:B:177:ALA:CB	2.61	0.48
1:A:34:SER:O	1:A:58:ASN:N	2.45	0.47
3:C:450:ILE:HB	3:C:530:ASN:HB2	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:79:THR:HB	2:B:86:PHE:CB	2.45	0.47
1:A:80:ASN:ND2	1:A:86:GLY:O	2.43	0.46
2:B:168:LEU:O	2:B:172:LYS:HB3	2.16	0.46
1:A:41:PRO:HB3	1:A:51:VAL:HG22	1.98	0.46
3:C:558:GLU:HG2	3:C:560:PRO:HD2	1.97	0.46
2:B:36:VAL:HG13	2:B:37:ASN:H	1.81	0.45
1:A:19:PRO:HD2	1:A:21:PHE:HE2	1.81	0.45
3:C:476:LEU:HD11	3:C:570:ILE:HG22	1.98	0.45
1:A:27:ILE:HG23	1:A:36:LEU:HD11	1.98	0.45
2:B:161:ASN:OD1	2:B:161:ASN:N	2.48	0.45
3:C:481:PHE:HA	3:C:521:LEU:HB2	1.98	0.45
4:D:11:DT:H2''	4:D:12:DT:O5'	2.16	0.45
3:C:518:ARG:NH2	4:D:13:DT:O2	2.41	0.44
2:B:79:THR:O	2:B:86:PHE:HB2	2.17	0.44
2:B:149:ILE:HG22	2:B:151:TYR:H	1.82	0.44
2:B:79:THR:HB	2:B:86:PHE:C	2.42	0.44
3:C:538:ASP:O	3:C:542:LYS:CB	2.66	0.44
1:A:22:ARG:HG2	1:A:76:VAL:HG22	2.00	0.44
1:A:97:PHE:HA	2:B:159:SER:HA	1.99	0.43
3:C:585:THR:HG23	3:C:590:SER:HB2	2.00	0.43
3:C:532:LEU:HG	3:C:533:TRP:H	1.84	0.43
1:A:75:PHE:CZ	1:A:89:ILE:HG12	2.53	0.43
2:B:85:ILE:HB	2:B:100:LYS:O	2.19	0.42
2:B:173:CYS:O	2:B:177:ALA:HB2	2.19	0.42
3:C:483:TYR:CE1	3:C:518:ARG:HB2	2.54	0.42
1:A:103:LEU:HD22	2:B:161:ASN:HB3	2.01	0.41
1:A:62:SER:HB3	1:A:65:LYS:HB2	2.03	0.41
1:A:16:VAL:HB	1:A:19:PRO:HG3	2.01	0.41
2:B:42:VAL:HG12	2:B:43:THR:H	1.85	0.41
3:C:526:ILE:HG22	3:C:531:GLN:HB3	2.03	0.40
4:D:15:DT:H6	4:D:15:DT:H2'	1.68	0.40
1:A:73:TYR:HA	1:A:93:VAL:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/122 (98%)	109 (91%)	9 (8%)	2 (2%)	7	35
2	B	128/132 (97%)	105 (82%)	23 (18%)	0	100	100
3	C	176/178 (99%)	157 (89%)	17 (10%)	2 (1%)	11	45
All	All	424/432 (98%)	371 (88%)	49 (12%)	4 (1%)	16	49

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	ILE
1	A	43	ILE
3	C	461	SER
3	C	466	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/113 (100%)	107 (95%)	6 (5%)	20	42
2	B	116/116 (100%)	106 (91%)	10 (9%)	10	30
3	C	157/157 (100%)	151 (96%)	6 (4%)	29	51
All	All	386/386 (100%)	364 (94%)	22 (6%)	20	41

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	GLU
1	A	20	VAL
1	A	28	LYS
1	A	30	GLN
1	A	97	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	121	ILE
2	B	36	VAL
2	B	38	THR
2	B	42	VAL
2	B	67	HIS
2	B	79	THR
2	B	98	VAL
2	B	101	TRP
2	B	126	GLN
2	B	147	LYS
2	B	180	MET
3	C	443	PHE
3	C	478	VAL
3	C	545	LEU
3	C	575	TYR
3	C	577	PHE
3	C	579	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	GLN
2	B	52	GLN
2	B	126	GLN
3	C	454	GLN
3	C	498	GLN
3	C	531	GLN
3	C	543	GLN
3	C	600	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	105:ALA	C	125:ALA	N	11.94


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-4410. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 60



Y Index: 60




Z Index: 60


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 62



Y Index: 55




Z Index: 50


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.045. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 43 nm^3 ; this corresponds to an approximate mass of 39 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.213 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-4410 and PDB model 6I52. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.045 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.045).

9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 57% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.045) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.5670	0.1010
A	0.5630	0.1000
B	0.5510	0.0970
C	0.5980	0.1140
D	0.5070	0.0630

