



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:31 AM UTC

PDB ID : 6I58 / pdb_00006i58
Title : Allosteric activation of human prekallikrein by apple domain disc rotation
Authors : Li, C.; Pathak, M.; McCrae, K.; Dreveny, I.; Emsley, J.
Deposited on : 2018-11-13
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	607	5% 78% 14% • •
2	B	2	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	706	-	-	X	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4753 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Coagulation factor XI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	585	Total	C	N	O	S	36	1	0
			4602	2903	796	864	39			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	Cl	0	0
			5	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	83	Total 83	O 83	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Coagulation factor XI



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	80.76Å 80.76Å 251.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.71 – 2.60 37.71 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.9 (37.71-2.60) 98.9 (37.71-2.60)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.42 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.208 , 0.280 0.217 , 0.281	Depositor DCC
R_{free} test set	2641 reflections (9.96%)	wwPDB-VP
Wilson B-factor (Å ²)	70.1	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4753	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PEG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/4712	1.06	13/6380 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	14	GLY	N-CA-C	8.73	125.08	114.69
1	A	531	TYR	CB-CA-C	-6.89	100.69	110.34
1	A	17	ILE	CB-CA-C	-6.41	105.35	111.44
1	A	406	GLN	CB-CA-C	-6.28	100.84	111.02
1	A	195	SER	N-CA-C	6.02	118.44	109.59
1	A	418	VAL	N-CA-C	-5.76	99.34	107.75
1	A	92	CYS	N-CA-C	5.54	117.68	108.20
1	A	249	THR	N-CA-C	5.49	117.74	109.23
1	A	315	THR	CB-CA-C	5.40	115.52	110.17
1	A	322	ASN	CB-CA-C	-5.35	110.39	116.54
1	A	229	PRO	N-CA-C	5.26	123.31	112.47
1	A	8	LYS	CA-CB-CG	-5.20	103.70	114.10
1	A	371	VAL	N-CA-C	-5.08	98.77	109.34

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	391	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4602	0	4479	62	0
2	B	28	0	25	0	0
3	A	5	0	0	0	0
4	A	7	0	10	4	0
5	A	28	0	26	1	0
6	A	83	0	0	0	0
All	All	4753	0	4540	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:367:LYS:HD2	1:A:368:PRO:HD2	1.19	1.13
1:A:406:GLN:HG3	1:A:471:THR:HG22	1.31	1.06
1:A:390:THR:HG23	1:A:423:ILE:O	1.59	1.01
1:A:509:LYS:HA	1:A:509:LYS:HE2	1.57	0.85
1:A:40:LEU:HD13	1:A:61:LYS:HB2	1.65	0.77
5:A:708:NAG:O7	5:A:708:NAG:O3	2.05	0.73
1:A:371:VAL:HG12	1:A:516:LYS:NZ	2.03	0.73
1:A:370:ILE:HD12	1:A:370:ILE:C	2.14	0.70
1:A:40:LEU:H	1:A:40:LEU:HD12	1.56	0.70
1:A:371:VAL:HG13	1:A:380:GLU:CD	2.18	0.69
1:A:479:ARG:HB3	4:A:706:PEG:H21	1.76	0.66
1:A:28:CYS:SG	1:A:41:PHE:CD2	2.89	0.66
1:A:371:VAL:HG12	1:A:516:LYS:HZ1	1.63	0.62
1:A:355:LEU:HA	1:A:358:MET:HE3	1.82	0.61
1:A:479:ARG:CB	4:A:706:PEG:H21	2.29	0.61
1:A:459:SER:OG	1:A:460:GLY:N	2.31	0.61
1:A:406:GLN:HG3	1:A:471:THR:CG2	2.18	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:317:ALA:CB	1:A:323:GLU:O	2.49	0.60
1:A:90:SER:O	1:A:91:ALA:HB3	2.03	0.57
1:A:323:GLU:HA	1:A:323:GLU:OE2	2.05	0.55
1:A:40:LEU:HD12	1:A:40:LEU:N	2.16	0.55
1:A:40:LEU:CD1	1:A:61:LYS:HB2	2.35	0.54
1:A:40:LEU:HD22	1:A:68:LEU:HD12	1.89	0.54
1:A:364:THR:HG23	1:A:365:LYS:HE3	1.90	0.54
1:A:367:LYS:HB3	1:A:369:ARG:NH1	2.23	0.53
1:A:29:GLN:HA	1:A:41:PHE:CE1	2.43	0.52
1:A:198:ASP:HB3	1:A:239:LEU:HD12	1.92	0.52
1:A:39:LEU:HD11	1:A:308:ARG:HG2	1.91	0.52
1:A:317:ALA:HB1	1:A:323:GLU:O	2.11	0.51
1:A:191:VAL:HG22	1:A:251:ILE:HB	1.93	0.50
1:A:474:TYR:CZ	1:A:480:PRO:HG3	2.46	0.49
1:A:367:LYS:CD	1:A:368:PRO:HD2	2.14	0.47
1:A:20:VAL:HB	1:A:156:THR:HG21	1.96	0.47
1:A:126:VAL:HA	1:A:180:LEU:HD11	1.96	0.46
1:A:367:LYS:HD2	1:A:368:PRO:CD	2.14	0.46
1:A:87:HIS:C	1:A:87:HIS:HD1	2.22	0.46
1:A:272:PHE:HB3	1:A:355:LEU:HB3	1.97	0.46
1:A:129:HIS:ND1	1:A:155:GLY:HA2	2.31	0.46
1:A:40:LEU:HD13	1:A:61:LYS:CB	2.42	0.45
1:A:532:ARG:NE	1:A:533:GLY:H	2.15	0.45
1:A:456:MET:HB2	1:A:456:MET:HE3	1.48	0.44
1:A:503:TYR:C	1:A:556:ASP:HA	2.43	0.44
1:A:252:LYS:O	1:A:253:LYS:CG	2.65	0.44
1:A:325:LYS:HD3	1:A:325:LYS:HA	1.65	0.44
1:A:479:ARG:CB	4:A:706:PEG:C2	2.97	0.43
1:A:371:VAL:HG13	1:A:380:GLU:OE2	2.17	0.43
1:A:87:HIS:C	1:A:87:HIS:ND1	2.77	0.43
1:A:390:THR:HG23	1:A:423:ILE:C	2.39	0.42
1:A:479:ARG:HB3	4:A:706:PEG:C2	2.46	0.42
1:A:415:PHE:CZ	1:A:466:LEU:HD21	2.53	0.42
1:A:263:GLN:H	1:A:263:GLN:HG3	1.35	0.42
1:A:370:ILE:O	1:A:370:ILE:HG23	2.19	0.42
1:A:530:ARG:HD3	1:A:586:ARG:HA	2.01	0.42
1:A:186:ILE:CG2	1:A:257:LEU:HD21	2.49	0.41
1:A:298:ALA:O	1:A:302:LEU:HD23	2.21	0.41
1:A:355:LEU:HD23	1:A:358:MET:HE1	2.03	0.41
1:A:186:ILE:HG23	1:A:257:LEU:HD21	2.02	0.41
1:A:532:ARG:HD2	1:A:532:ARG:HA	1.27	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:LYS:O	1:A:253:LYS:HG2	2.21	0.41
1:A:254:SER:O	1:A:255:LYS:HB2	2.20	0.41
1:A:357:LYS:HA	1:A:357:LYS:HD3	1.92	0.40
1:A:409:LEU:HD12	1:A:409:LEU:HA	1.86	0.40
1:A:564:LYS:HB2	1:A:569:TRP:CZ3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	576/607 (95%)	549 (95%)	27 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	520/535 (97%)	472 (91%)	48 (9%)	8	19

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	40	LEU
1	A	57	THR
1	A	84	GLN
1	A	86	SER
1	A	87	HIS
1	A	90	SER
1	A	142	GLU
1	A	156	THR
1	A	166	VAL
1	A	173	LYS
1	A	175	CYS
1	A	186	ILE
1	A	191	VAL
1	A	200	VAL
1	A	231	GLU
1	A	246	LEU
1	A	250	ARG
1	A	252	LYS
1	A	253	LYS
1	A	257	LEU
1	A	261	SER
1	A	263	GLN
1	A	294	LYS
1	A	323	GLU
1	A	325	LYS
1	A	337	SER
1	A	365	LYS
1	A	366	ILE
1	A	370	ILE
1	A	387	LEU
1	A	446	GLN
1	A	450	ILE
1	A	456	MET
1	A	459	SER
1	A	499	THR
1	A	508	ASP
1	A	509	LYS
1	A	510	ILE
1	A	515	GLN
1	A	532	ARG
1	A	534	HIS
1	A	541	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	545	TYR
1	A	554	LYS
1	A	579	GLU
1	A	594	VAL
1	A	604	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	106	ASN
1	A	116	GLN
1	A	233	GLN
1	A	235	ASN
1	A	274	HIS
1	A	343	HIS
1	A	384	GLN
1	A	566	ASN
1	A	570	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.78	0	17,19,21	1.50	3 (17%)
2	NAG	B	2	2	14,14,15	0.66	0	17,19,21	1.04	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	6/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	C1-O5-C5	3.46	116.82	112.19
2	B	1	NAG	O4-C4-C3	-3.18	102.89	110.38
2	B	2	NAG	O5-C1-C2	-2.92	106.78	111.29
2	B	1	NAG	O5-C5-C6	2.05	111.65	107.66

There are no chirality outliers.


All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C8-C7-N2-C2
2	B	1	NAG	O7-C7-N2-C2
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	1	NAG	C3-C2-N2-C7
2	B	2	NAG	O5-C5-C6-O6
2	B	2	NAG	C4-C5-C6-O6
2	B	1	NAG	C1-C2-N2-C7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 5 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	A	707	1	14,14,15	0.71	0	17,19,21	2.59	4 (23%)
5	NAG	A	708	1	14,14,15	0.30	0	17,19,21	0.56	0
4	PEG	A	706	-	6,6,6	0.44	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	A	707	1	-	3/6/23/26	0/1/1/1
5	NAG	A	708	1	-	6/6/23/26	0/1/1/1
4	PEG	A	706	-	-	1/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	707	NAG	C2-N2-C7	7.43	132.86	122.90
5	A	707	NAG	C8-C7-N2	5.25	124.82	116.12
5	A	707	NAG	C1-O5-C5	3.84	117.34	112.19
5	A	707	NAG	O7-C7-C8	-3.05	116.63	122.05

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	707	NAG	C3-C2-N2-C7
5	A	708	NAG	C8-C7-N2-C2
5	A	707	NAG	C8-C7-N2-C2
5	A	707	NAG	O7-C7-N2-C2
5	A	708	NAG	O7-C7-N2-C2
5	A	708	NAG	C4-C5-C6-O6
4	A	706	PEG	O1-C1-C2-O2
5	A	708	NAG	O5-C5-C6-O6
5	A	708	NAG	C1-C2-N2-C7
5	A	708	NAG	C3-C2-N2-C7

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	708	NAG	1	0
4	A	706	PEG	4	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	585/607 (96%)	0.29	29 (4%) 34 28	30, 70, 113, 177	20 (3%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	370	ILE	4.4
1	A	391	SER	4.1
1	A	371	VAL	3.6
1	A	554	LYS	3.3
1	A	536	ILE	2.9
1	A	367	LYS	2.8
1	A	165	LYS	2.7
1	A	510	ILE	2.7
1	A	452	ASP	2.6
1	A	366	ILE	2.6
1	A	580	GLY	2.6
1	A	41	PHE	2.6
1	A	317	ALA	2.6
1	A	531	TYR	2.5
1	A	456	MET	2.5
1	A	193	ALA	2.4
1	A	532	ARG	2.4
1	A	87	HIS	2.3
1	A	355	LEU	2.3
1	A	89	ILE	2.2
1	A	364	THR	2.2
1	A	186	ILE	2.2
1	A	527	CYS	2.1
1	A	509	LYS	2.1
1	A	587	PRO	2.1
1	A	431	LEU	2.1
1	A	429	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	92	CYS	2.1
1	A	323	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	B	1	14/15	0.75	0.14	54,59,73,73	14
2	NAG	B	2	14/15	0.78	0.13	50,57,59,61	14

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	NAG	A	708	14/15	0.44	0.14	117,136,148,150	0
5	NAG	A	707	14/15	0.53	0.20	101,113,123,123	14
4	PEG	A	706	7/7	0.87	0.14	54,55,59,60	1
3	CL	A	701	1/1	0.93	0.14	89,89,89,89	0
3	CL	A	704	1/1	0.93	0.09	88,88,88,88	0
3	CL	A	705	1/1	0.93	0.15	95,95,95,95	0
3	CL	A	703	1/1	0.94	0.17	90,90,90,90	0
3	CL	A	702	1/1	0.94	0.13	80,80,80,80	0

6.5 Other polymers [i](#)

There are no such residues in this entry.