



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:41 AM UTC

PDB ID : 6I5E / pdb_00006i5e
Title : X-ray structure of apo human soluble Epoxide Hydrolase C-terminal Domain (hsEH CTD)
Authors : Abis, G.; Kopec, J.; Yue, W.W.; Conte, M.R.
Deposited on : 2018-11-13
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	
1	B	344	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional epoxide hydrolase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2543	1635	426	459	23			
1	B	316	Total	C	N	O	S	0	0	0
			2543	1635	426	459	23			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	212	MET	-	initiating methionine	UNP P34913
A	213	HIS	-	expression tag	UNP P34913
A	214	HIS	-	expression tag	UNP P34913
A	215	HIS	-	expression tag	UNP P34913
A	216	HIS	-	expression tag	UNP P34913
A	217	HIS	-	expression tag	UNP P34913
A	218	HIS	-	expression tag	UNP P34913
A	219	SER	-	expression tag	UNP P34913
A	220	THR	-	expression tag	UNP P34913
A	221	GLU	-	expression tag	UNP P34913
A	222	ASN	-	expression tag	UNP P34913
A	223	LEU	-	expression tag	UNP P34913
A	224	TYR	-	expression tag	UNP P34913
A	225	PHE	-	expression tag	UNP P34913
A	226	GLN	-	expression tag	UNP P34913
A	227	GLY	-	expression tag	UNP P34913
A	228	SER	-	expression tag	UNP P34913
A	229	SER	-	expression tag	UNP P34913
B	212	MET	-	initiating methionine	UNP P34913
B	213	HIS	-	expression tag	UNP P34913
B	214	HIS	-	expression tag	UNP P34913
B	215	HIS	-	expression tag	UNP P34913
B	216	HIS	-	expression tag	UNP P34913
B	217	HIS	-	expression tag	UNP P34913
B	218	HIS	-	expression tag	UNP P34913

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	219	SER	-	expression tag	UNP P34913
B	220	THR	-	expression tag	UNP P34913
B	221	GLU	-	expression tag	UNP P34913
B	222	ASN	-	expression tag	UNP P34913
B	223	LEU	-	expression tag	UNP P34913
B	224	TYR	-	expression tag	UNP P34913
B	225	PHE	-	expression tag	UNP P34913
B	226	GLN	-	expression tag	UNP P34913
B	227	GLY	-	expression tag	UNP P34913
B	228	SER	-	expression tag	UNP P34913
B	229	SER	-	expression tag	UNP P34913


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	9	Total O 9 9	0	0
2	B	4	Total O 4 4	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional epoxide hydrolase 2



- Molecule 1: Bifunctional epoxide hydrolase 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.22Å 80.14Å 104.70Å 90.00° 95.39° 90.00°	Depositor
Resolution (Å)	70.51 – 2.60 70.51 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.9 (70.51-2.60) 99.9 (70.51-2.60)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.246 , 0.277 0.247 , 0.276	Depositor DCC
R_{free} test set	1101 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	34.0	Xtriage
Anisotropy	0.818	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 25.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	5099	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2619	0.91	0/3553
1	B	0.72	0/2619	0.91	1/3553 (0.0%)
All	All	0.73	0/5238	0.91	1/7106 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	234	PRO	N-CA-C	5.02	121.23	113.75

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	291	MET	Peptide
1	B	291	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2543	0	2465	19	0
1	B	2543	0	2465	10	0
2	A	9	0	0	0	0
2	B	4	0	0	0	0
All	All	5099	0	4930	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:369:MET:SD	1:A:374:SER:OG	2.27	0.93
1:A:375:ILE:O	1:A:377:ALA:N	2.06	0.89
1:A:375:ILE:C	1:A:377:ALA:H	1.95	0.74
1:A:375:ILE:O	1:A:378:ASN:N	2.23	0.65
1:A:375:ILE:C	1:A:377:ALA:N	2.54	0.64
1:B:500:VAL:CG1	1:B:502:GLN:HG3	2.29	0.62
1:A:266:GLY:HA3	1:A:335:ASP:HB3	1.82	0.60
1:B:266:GLY:HA3	1:B:335:ASP:HB3	1.91	0.53
1:B:310:MET:HG2	1:B:343:TYR:CD2	2.45	0.52
1:A:254:GLU:HG3	1:A:289:LEU:HD23	1.93	0.51
1:A:545:ASP:N	1:A:545:ASP:OD1	2.42	0.51
1:A:369:MET:SD	1:A:374:SER:CA	2.99	0.50
1:B:339:MET:HE3	1:B:339:MET:HA	1.94	0.50
1:A:369:MET:SD	1:A:374:SER:CB	3.01	0.48
1:A:375:ILE:O	1:A:376:LYS:C	2.58	0.47
1:B:364:PRO:HG2	1:B:510:TRP:CE2	2.50	0.46
1:A:306:GLU:H	1:A:306:GLU:CD	2.25	0.44
1:A:332:ILE:HG12	1:A:356:ALA:HB3	2.00	0.44
1:A:245:LYS:HB2	1:A:246:PRO:HD2	2.01	0.43
1:A:310:MET:HG3	1:A:343:TYR:CE2	2.53	0.42
1:B:234:PRO:HG2	1:B:449:PHE:CE1	2.54	0.42
1:A:369:MET:SD	1:A:374:SER:N	2.93	0.42
1:A:269:GLU:HB3	1:A:273:SER:OG	2.20	0.42
1:A:277:GLN:HG3	1:A:535:ASN:ND2	2.35	0.42
1:B:500:VAL:HG12	1:B:502:GLN:HG3	2.02	0.41
1:B:245:LYS:O	1:B:246:PRO:C	2.63	0.41
1:B:355:VAL:O	1:B:488:ALA:HA	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:369:MET:O	1:A:369:MET:HG3	2.21	0.41
1:B:378:ASN:HD22	1:B:379:PRO:HD2	1.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/344 (91%)	302 (96%)	10 (3%)	2 (1%)	21	42
1	B	314/344 (91%)	306 (98%)	7 (2%)	1 (0%)	36	58
All	All	628/688 (91%)	608 (97%)	17 (3%)	3 (0%)	24	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	376	LYS
1	A	268	PRO
1	B	268	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/304 (91%)	264 (96%)	12 (4%)	26	51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	276/304 (91%)	267 (97%)	9 (3%)	33	61
All	All	552/608 (91%)	531 (96%)	21 (4%)	29	56

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	ARG
1	A	369	MET
1	A	372	LEU
1	A	374	SER
1	A	378	ASN
1	A	452	GLN
1	A	474	LYS
1	A	480	LEU
1	A	484	ILE
1	A	489	LEU
1	A	508	GLU
1	A	545	ASP
1	B	234	PRO
1	B	339	MET
1	B	340	LEU
1	B	346	LEU
1	B	421	LYS
1	B	471	ARG
1	B	489	LEU
1	B	500	VAL
1	B	508	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	384	GLN
1	A	388	GLN
1	A	453	GLN
1	A	505	GLN
1	A	535	ASN
1	B	378	ASN
1	B	384	GLN
1	B	452	GLN
1	B	453	GLN
1	B	513	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	535	ASN
1	B	536	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/344 (91%)	0.40	13 (4%) 41 36	24, 41, 78, 114	0
1	B	316/344 (91%)	0.38	18 (5%) 29 24	23, 41, 78, 113	0
All	All	632/688 (91%)	0.39	31 (4%) 35 29	23, 41, 78, 114	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	418	SER	4.3
1	B	343	TYR	4.0
1	B	377	ALA	3.3
1	B	480	LEU	3.1
1	A	425	ALA	2.9
1	B	419	MET	2.8
1	A	480	LEU	2.8
1	A	380	VAL	2.8
1	B	513	HIS	2.7
1	A	510	TRP	2.7
1	B	510	TRP	2.7
1	A	420	HIS	2.6
1	B	230	THR	2.6
1	B	427	GLY	2.5
1	B	469	MET	2.5
1	A	427	GLY	2.5
1	A	377	ALA	2.4
1	A	230	THR	2.4
1	B	430	VAL	2.3
1	B	425	ALA	2.3
1	B	420	HIS	2.3
1	A	421	LYS	2.2
1	A	469	MET	2.2
1	A	379	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	498	VAL	2.2
1	A	498	VAL	2.1
1	B	422	VAL	2.1
1	B	379	PRO	2.1
1	A	375	ILE	2.0
1	B	481	GLY	2.0
1	B	362	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.