



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 02:51 PM EDT

PDB ID : 8I58 / pdb_00008i58
Title : Uroporphyrin I (UPI)-bound CfbA
Authors : Ogawa, S.; Fujishiro, T.
Deposited on : 2023-01-24
Resolution : 3.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	143	 64% 17% 17%
1	B	143	 65% 17% 17%

2 Entry composition ⓘ


There are 3 unique types of molecules in this entry. The entry contains 1968 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sirohydrochlorin cobaltochelata

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	119	Total	C	N	O	S	0	1	0
			950	617	164	167	2			
1	B	119	Total	C	N	O	S	0	1	0
			951	616	164	169	2			

- Molecule 2 is 3-[(1Z,9Z)-3,8,13,18-tetrakis(2-hydroxy-2-oxoethyl)-7,12,17-tris(3-hydroxy-3-oxopropyl)-21,23-dihydroporphyrin-2-yl]propanoic acid (CCD ID: UPI) (formula: C₄₀H₃₈N₄O₁₆) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			60	40	4	16		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	O 3	0	0
3	B	4	Total 4	O 4	0	0

- Molecule 1: Sirohydrochlorin cobaltochelatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	68.81Å 68.81Å 83.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	31.67 – 3.09 31.67 – 3.09	Depositor EDS
% Data completeness (in resolution range)	99.9 (31.67-3.09) 99.9 (31.67-3.09)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 3.07Å)	Xtriage
Refinement program	PHENIX 1.18.2	Depositor
R, R_{free}	0.157 , 0.204 0.157 , 0.206	Depositor DCC
R_{free} test set	361 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	77.6	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.479 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1968	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UPI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/969	1.02	1/1308 (0.1%)
1	B	0.55	0/970	1.08	1/1309 (0.1%)
All	All	0.55	0/1939	1.05	2/2617 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	7
All	All	0	11

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	114	ILE	N-CA-C	5.19	113.64	107.73
1	B	87	ILE	N-CA-CB	-5.11	103.05	111.44

There are no chirality outliers.

All (11) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	123	ARG	Mainchain
1	A	128	ALA	Mainchain
1	A	34	PRO	Mainchain
1	A	44	SER	Mainchain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	114	ILE	Mainchain
1	B	117	ASP	Mainchain
1	B	12	ARG	Mainchain
1	B	123	ARG	Mainchain
1	B	29	GLU	Mainchain
1	B	34	PRO	Mainchain
1	B	44	SER	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	950	0	1012	23	0
1	B	951	0	1007	21	0
2	A	60	0	0	1	0
3	A	3	0	0	0	0
3	B	4	0	0	0	0
All	All	1968	0	2019	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:75[B]:HIS:HD2	1:B:76:THR:HG23	1.36	0.91
1:B:75[B]:HIS:CD2	1:B:76:THR:H	2.09	0.70
1:B:20:LEU:HD12	1:B:40:LEU:HD21	1.76	0.66
1:A:128:ALA:HB2	1:B:19:LEU:HD12	1.80	0.62
1:B:17:LYS:HG3	1:B:40:LEU:HD11	1.82	0.62
1:A:20:LEU:HD12	1:A:40:LEU:HD21	1.83	0.60
1:A:64:ILE:HG12	1:A:121:ILE:HD11	1.83	0.59
1:A:19:LEU:HD12	1:B:128:ALA:HB2	1.85	0.59
1:A:35:ILE:CD1	1:A:58:GLN:HB2	2.36	0.56
1:B:75[B]:HIS:CD2	1:B:76:THR:HG23	2.27	0.56
1:B:8:GLY:HA3	1:B:20:LEU:HD11	1.88	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:ILE:HD13	1:B:58:GLN:HB2	1.90	0.54
1:A:40:LEU:HD13	1:A:44:SER:HB2	1.89	0.53
1:A:54:LYS:O	1:A:58:GLN:HG3	2.09	0.53
1:A:17:LYS:HG3	1:A:40:LEU:HD11	1.90	0.53
1:A:73:GLY:HA3	1:A:75[B]:HIS:CE1	2.45	0.52
1:A:27:VAL:HG22	1:B:133:VAL:HG13	1.92	0.51
1:A:71:ALA:HB2	1:B:19:LEU:HD22	1.93	0.51
1:A:133:VAL:HG13	1:B:27:VAL:HG22	1.92	0.50
1:A:8:GLY:HA3	1:A:20:LEU:HD11	1.93	0.50
1:B:20:LEU:HD12	1:B:40:LEU:CD2	2.40	0.50
1:A:35:ILE:HD13	1:A:58:GLN:HB2	1.95	0.49
1:A:20:LEU:HD12	1:A:40:LEU:CD2	2.43	0.48
1:B:41:MET:SD	1:B:75[A]:HIS:HB3	2.54	0.48
1:B:75[A]:HIS:HA	1:B:79:ASP:HB2	1.96	0.47
1:A:33:PHE:HA	1:A:34:PRO:HD3	1.86	0.45
1:A:19:LEU:HD22	1:B:71:ALA:HB2	1.98	0.45
2:A:201:UPI:CAV	1:B:75[B]:HIS:CE1	3.02	0.43
1:A:41:MET:SD	1:A:75[A]:HIS:HB3	2.59	0.43
1:A:75[B]:HIS:CD2	1:A:76:THR:OG1	2.72	0.43
1:B:61:LYS:HD3	1:B:118:VAL:HG12	1.99	0.43
1:A:25:GLU:O	1:A:29:GLU:HG3	2.19	0.42
1:A:67:PRO:HG3	1:A:122:TYR:CE1	2.54	0.42
1:B:75[B]:HIS:CG	1:B:76:THR:H	2.38	0.42
1:A:54:LYS:HE2	1:A:54:LYS:HB3	1.81	0.41
1:A:41:MET:HE2	1:A:41:MET:HB3	1.94	0.41
1:B:2:GLU:O	1:B:34:PRO:HD2	2.21	0.41
1:B:41:MET:HE2	1:B:41:MET:HB3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/143 (81%)	113 (97%)	3 (3%)	0	100	100
1	B	116/143 (81%)	111 (96%)	5 (4%)	0	100	100
All	All	232/286 (81%)	224 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/127 (82%)	104 (100%)	0	100	100
1	B	104/127 (82%)	104 (100%)	0	100	100
All	All	208/254 (82%)	208 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	B	50	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	UPI	A	201	-	56,64,64	3.90	20 (35%)	56,92,92	5.31	26 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	UPI	A	201	-	-	20/36/52/52	0/4/5/5

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	UPI	CCF-CBX	-16.34	1.21	1.39
2	A	201	UPI	CCG-CBY	-10.75	1.27	1.39
2	A	201	UPI	CCC-CBU	-8.24	1.16	1.45
2	A	201	UPI	CBF-CBR	7.39	1.66	1.51
2	A	201	UPI	CBU-CBS	-6.30	1.27	1.34
2	A	201	UPI	CBE-CBQ	6.21	1.64	1.51
2	A	201	UPI	CCB-CBT	-6.18	1.23	1.45
2	A	201	UPI	CAZ-CBT	6.12	1.60	1.51
2	A	201	UPI	CCA-CBS	-5.93	1.24	1.45
2	A	201	UPI	CCE-CBW	-5.36	1.33	1.39
2	A	201	UPI	CBD-CBV	3.64	1.58	1.50
2	A	201	UPI	CBV-CBT	-3.24	1.31	1.34
2	A	201	UPI	CBY-CBX	-3.12	1.32	1.39
2	A	201	UPI	CBF-CBZ	3.01	1.58	1.51
2	A	201	UPI	CBC-CBU	3.00	1.57	1.50
2	A	201	UPI	CCH-NBJ	-2.93	1.33	1.38
2	A	201	UPI	OAN-CBP	-2.21	1.23	1.30
2	A	201	UPI	CAX-CBN	2.18	1.55	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	UPI	CBZ-CBW	-2.13	1.34	1.39
2	A	201	UPI	CCF-NBI	2.08	1.42	1.38

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	UPI	CBX-CCF-NBI	-23.42	92.50	109.43
2	A	201	UPI	CBY-CCG-NBI	-17.14	97.04	109.43
2	A	201	UPI	CBC-CBU-CCC	-12.31	103.65	124.80
2	A	201	UPI	CCC-CBU-CBS	10.11	117.11	109.39
2	A	201	UPI	CBZ-CCH-NBJ	-9.36	102.67	109.43
2	A	201	UPI	CBZ-CBF-CBR	-9.19	103.57	114.20
2	A	201	UPI	CAZ-CBT-CCB	-7.86	104.34	124.70
2	A	201	UPI	CBY-CBE-CBQ	-6.40	106.80	114.20
2	A	201	UPI	CAW-CBA-CBW	5.09	121.09	112.54
2	A	201	UPI	CAY-CBS-CCA	-4.00	114.34	124.70
2	A	201	UPI	CAY-CAU-CBK	-3.67	103.94	113.67
2	A	201	UPI	OAK-CBM-CAW	-3.45	112.14	123.09
2	A	201	UPI	CCB-CBT-CBV	3.25	111.87	109.39
2	A	201	UPI	OAE-CBO-CBC	-3.23	112.91	122.11
2	A	201	UPI	OAA-CBK-CAU	-3.16	113.05	123.09
2	A	201	UPI	CAZ-CAV-CBL	-3.15	105.32	113.67
2	A	201	UPI	CBW-CCE-NBJ	2.70	111.38	109.43
2	A	201	UPI	CBB-CBX-CBY	-2.56	116.83	124.90
2	A	201	UPI	CBD-CBV-CCD	2.41	128.94	124.80
2	A	201	UPI	OAI-CBK-CAU	2.38	121.52	114.00
2	A	201	UPI	OAD-CBN-CAX	-2.31	115.75	123.09
2	A	201	UPI	CCG-NBI-CCF	2.20	112.03	108.82
2	A	201	UPI	OAC-CBM-CAW	2.20	120.96	114.00
2	A	201	UPI	OAG-CBQ-CBE	-2.14	116.50	122.94
2	A	201	UPI	CCD-CBV-CBT	-2.11	107.77	109.39
2	A	201	UPI	CBP-CBD-CBV	2.10	119.33	113.73

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	UPI	CAU-CAY-CBS-CCA
2	A	201	UPI	CAV-CAZ-CBT-CCB
2	A	201	UPI	CAX-CBB-CBX-CCF
2	A	201	UPI	CBO-CBC-CBU-CCC
2	A	201	UPI	CBN-CAX-CBB-CBX

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	201	UPI	CBZ-CBF-CBR-OAP
2	A	201	UPI	CBU-CBC-CBO-OAM
2	A	201	UPI	CBV-CBD-CBP-OAF
2	A	201	UPI	CBA-CAW-CBM-OAK
2	A	201	UPI	CAY-CAU-CBK-OAI
2	A	201	UPI	CBV-CBD-CBP-OAN
2	A	201	UPI	CAY-CAU-CBK-OAA
2	A	201	UPI	CAZ-CAV-CBL-OAB
2	A	201	UPI	CBA-CAW-CBM-OAC
2	A	201	UPI	CBY-CBE-CBQ-OAO
2	A	201	UPI	CAZ-CAV-CBL-OAJ
2	A	201	UPI	CBU-CBC-CBO-OAE
2	A	201	UPI	CBZ-CBF-CBR-OAH
2	A	201	UPI	CBY-CBE-CBQ-OAG
2	A	201	UPI	CBL-CAV-CAZ-CBT

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	UPI	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	119/143 (83%)	-1.29	0 100 100	40, 70, 109, 147	1 (0%)
1	B	119/143 (83%)	-1.26	0 100 100	42, 69, 114, 148	1 (0%)
All	All	238/286 (83%)	-1.27	0 100 100	40, 70, 110, 148	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	UPI	A	201	60/60	0.96	0.11	72,161,184,187	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around UPI A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.