



Full wwPDB X-ray Structure Validation Report ⓘ

May 6, 2026 – 10:12 am BST

PDB ID : 9I5B / pdb_00009i5b
Title : Crystal structure of perlecan region 3 mutant (P1019L) construct I876-V1272 including one laminin IV-like and four laminin EGF-like domains.
Authors : Sohail, A.A.; Koski, M.K.; Ruddock, L.W.
Deposited on : 2025-01-27
Resolution : 2.52 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.52 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	397	5% 76% 21% ..
1	B	397	4% 54% 15% 29% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 5331 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Basement membrane-specific heparan sulfate proteoglycan core protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	392	Total	C	N	O	S	0	2	0
			2941	1776	542	588	35			
1	B	282	Total	C	N	O	S	0	2	0
			2142	1312	387	425	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1019	LEU	PRO	engineered mutation	UNP Q05793
B	1019	LEU	PRO	engineered mutation	UNP Q05793

- Molecule 2 is SUCCINIC ACID (CCD ID: SIN) (formula: $C_4H_6O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 8	C 4	O 4	0	0
2	B	1	Total 8	C 4	O 4	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	142	Total 142	O 142	0	0
3	B	90	Total 90	O 90	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.65Å 74.97Å 81.37Å 75.23° 73.12° 80.11°	Depositor
Resolution (Å)	47.67 – 2.52 47.67 – 2.52	Depositor EDS
% Data completeness (in resolution range)	98.5 (47.67-2.52) 98.5 (47.67-2.52)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.88)	Depositor
R, R_{free}	0.188 , 0.220 0.188 , 0.220	Depositor DCC
R_{free} test set	1997 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.845	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 63.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5331	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	0/3022	1.27	18/4106 (0.4%)
1	B	0.68	1/2197 (0.0%)	1.24	9/2983 (0.3%)
All	All	0.67	1/5219 (0.0%)	1.26	27/7089 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	970	SER	C-O	-5.09	1.21	1.25

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	891	THR	CA-CB-OG1	-11.04	93.05	109.60
1	B	1002	VAL	N-CA-CB	8.32	120.87	110.47
1	A	891	THR	CA-CB-OG1	-7.96	97.65	109.60
1	A	1002	VAL	N-CA-CB	7.91	121.30	110.54
1	A	988	PRO	CB-CA-C	7.57	121.28	111.21
1	A	1151	LEU	N-CA-CB	7.18	122.62	110.49
1	A	1012	THR	CA-CB-OG1	-6.79	99.41	109.60
1	A	899	VAL	N-CA-CB	-6.75	101.26	112.06
1	B	988	PRO	CB-CA-C	6.72	120.15	111.21
1	A	1054	ASN	CB-CA-C	-6.40	99.41	109.84
1	A	1014	MET	CG-SD-CE	-6.24	87.18	100.90
1	B	1012	THR	CA-CB-OG1	-6.20	100.30	109.60
1	A	1097	PRO	CB-CA-C	6.02	118.87	110.98
1	A	1261	GLN	CB-CA-C	-5.99	99.64	109.46
1	B	1157	GLU	CB-CG-CD	5.82	122.50	112.60
1	B	881	GLU	CB-CG-CD	-5.65	103.00	112.60
1	A	1075	GLU	CB-CA-C	-5.64	101.79	110.81
1	A	975	GLU	CB-CG-CD	5.55	122.03	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1266	ASP	CA-CB-CG	5.52	118.12	112.60
1	B	1075	GLU	CB-CA-C	-5.44	101.75	110.79
1	A	1231	HIS	CA-CB-CG	-5.24	108.56	113.80
1	A	1227	ASP	CA-CB-CG	5.24	117.84	112.60
1	A	1200	ARG	N-CA-CB	5.18	117.52	110.01
1	A	964	THR	CA-CB-OG1	-5.13	101.91	109.60
1	B	1025	HIS	CA-CB-CG	-5.12	108.68	113.80
1	B	1062	GLN	N-CA-CB	5.11	118.70	110.42
1	A	1060	GLN	N-CA-CB	5.03	120.24	111.13

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2941	0	2689	46	0
1	B	2142	0	2028	35	0
2	A	8	0	4	0	0
2	B	8	0	4	0	0
3	A	142	0	0	9	0
3	B	90	0	0	5	0
All	All	5331	0	4725	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1021:SER:HA	3:A:1412:HOH:O	1.67	0.94
1:B:917:ASN:ND2	1:B:922:LEU:HD23	1.96	0.80
1:B:914:SER:HB2	1:B:916:GLN:NE2	1.99	0.77
1:A:1214:ALA:O	1:A:1238:SER:HB2	1.87	0.74
1:A:1256:ASN:C	1:A:1256:ASN:HD22	1.96	0.73

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:917:ASN:ND2	1:A:922:LEU:HD23	2.03	0.73
1:B:1060:GLN:OE1	1:B:1062:GLN:HG2	1.89	0.73
1:B:1026:ARG:HD3	3:B:1422:HOH:O	1.94	0.68
1:B:924:CYS:O	1:B:931:ARG:NH1	2.29	0.65
1:A:1193:GLY:HA2	1:A:1222:HIS:O	1.96	0.65
1:B:1092:SER:HB3	1:B:1097:PRO:HG3	1.79	0.64
1:A:1224:CYS:HB2	1:A:1233:THR:O	2.00	0.62
1:A:949:GLN:NE2	3:A:1406:HOH:O	2.32	0.61
1:A:1092:SER:HB3	1:A:1097:PRO:HG3	1.82	0.60
1:A:1115:THR:HB	1:B:1115:THR:HB	1.83	0.59
1:B:877:VAL:HG21	1:B:901:ARG:NH1	2.18	0.58
1:A:1227:ASP:C	1:A:1229:ASP:H	2.16	0.54
1:B:1026:ARG:CG	3:B:1422:HOH:O	2.56	0.53
1:B:917:ASN:HD22	1:B:922:LEU:HD23	1.73	0.53
1:B:1026:ARG:CD	3:B:1422:HOH:O	2.55	0.52
1:B:1003:THR:HB	1:B:1110:ALA:HB3	1.92	0.52
1:A:985:LEU:CD1	1:A:985:LEU:N	2.73	0.52
1:A:1003:THR:HB	1:A:1110:ALA:HB3	1.90	0.52
1:B:1008:GLU:HB2	1:B:1010:ARG:NH1	2.25	0.52
1:B:1128:CYS:HB3	1:B:1132:TYR:HB2	1.92	0.51
1:A:1008:GLU:HB2	1:A:1010:ARG:NH1	2.26	0.51
1:B:1066:ARG:NH1	1:B:1070:GLN:HG3	2.25	0.51
1:A:934:SER:HB2	3:A:1512:HOH:O	2.10	0.51
1:A:986:SER:HB2	3:A:1468:HOH:O	2.11	0.50
1:B:1027:GLN:HG3	3:B:1444:HOH:O	2.11	0.50
1:A:992:SER:OG	3:A:1401:HOH:O	2.20	0.50
1:A:1144:TYR:HA	1:A:1157:GLU:O	2.12	0.49
1:B:877:VAL:HG21	1:B:901:ARG:HH12	1.78	0.49
1:A:1128:CYS:HB3	1:A:1132:TYR:HB2	1.94	0.49
1:A:1215:PRO:HB2	1:A:1219:GLN:HB2	1.93	0.49
1:A:1238:SER:OG	1:A:1239:PRO:HD2	2.12	0.48
1:A:985:LEU:N	1:A:985:LEU:HD12	2.27	0.48
1:A:879:CYS:HB3	1:A:884:SER:OG	2.13	0.48
1:B:1149:SER:O	1:B:1151:LEU:N	2.47	0.48
1:B:1045:ARG:NH1	1:B:1053:SER:HB3	2.28	0.48
1:B:934:SER:HB2	3:B:1443:HOH:O	2.14	0.48
1:B:1092:SER:O	1:B:1097:PRO:HB3	2.14	0.48
1:A:1131:GLY:HA2	1:A:1144:TYR:CD1	2.48	0.47
1:B:916:GLN:H	1:B:916:GLN:CD	2.22	0.47
1:A:1024:LEU:HB3	1:A:1027:GLN:HG3	1.96	0.46
1:A:1237[B]:CYS:HA	1:A:1246:CYS:SG	2.56	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1013:VAL:O	1:B:1052:PRO:HA	2.16	0.46
1:B:1024:LEU:HB3	1:B:1027:GLN:OE1	2.16	0.46
1:A:1023:PRO:O	1:A:1095[B]:GLN:NE2	2.49	0.46
1:B:909:GLY:HA2	1:B:931:ARG:O	2.17	0.45
1:B:1023:PRO:O	1:B:1095[A]:GLN:NE2	2.50	0.44
1:A:1020:SER:N	3:A:1402:HOH:O	2.23	0.44
1:B:917:ASN:HD21	1:B:922:LEU:HD23	1.77	0.44
1:B:880:ASP:HB2	1:B:904:ASN:HB3	2.00	0.44
1:A:1066:ARG:NH1	1:A:1070:GLN:HG3	2.32	0.43
1:A:946:ALA:HB2	3:A:1474:HOH:O	2.17	0.43
1:A:998:ARG:HH11	1:A:998:ARG:HD3	1.64	0.43
1:A:1092:SER:O	1:A:1097:PRO:HB3	2.18	0.43
1:A:1256:ASN:C	1:A:1256:ASN:ND2	2.69	0.43
1:A:1227:ASP:O	1:A:1229:ASP:N	2.52	0.43
1:B:879:CYS:HB3	1:B:884:SER:OG	2.18	0.43
1:A:917:ASN:HD21	1:A:922:LEU:HD23	1.82	0.42
1:B:953:PHE:CG	1:B:976:LEU:HD11	2.53	0.42
1:B:1018:ARG:O	1:B:1019:LEU:C	2.61	0.42
1:A:953:PHE:CE2	1:A:994:PRO:HD3	2.55	0.42
1:A:1024:LEU:HB3	1:A:1027:GLN:CG	2.50	0.42
1:A:1149:SER:HB3	1:A:1150:GLY:H	1.65	0.42
1:A:1199:GLN:HE21	1:A:1199:GLN:HB3	1.62	0.41
1:B:953:PHE:CG	1:B:976:LEU:CD1	3.03	0.41
1:A:1143:GLY:HA2	1:A:1166:GLU:O	2.19	0.41
1:A:1227:ASP:C	1:A:1229:ASP:N	2.79	0.41
1:B:906:CYS:HB3	1:B:910:SER:HB2	2.02	0.41
1:A:1070:GLN:HG3	3:A:1413:HOH:O	2.19	0.41
1:A:939:SER:O	1:A:1110:ALA:HA	2.21	0.41
1:A:1249:CYS:SG	1:A:1255:GLY:O	2.79	0.41
1:B:1149:SER:HB2	1:B:1154:GLY:HA2	2.02	0.41
1:A:1125:GLN:NE2	3:A:1419:HOH:O	2.54	0.40
1:B:1131:GLY:HA2	1:B:1144:TYR:CD1	2.56	0.40
1:A:1223:THR:OG1	1:A:1236:SER:HB3	2.21	0.40
1:A:911:PHE:CD1	1:A:933:CYS:SG	3.14	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/397 (99%)	367 (94%)	20 (5%)	5 (1%)	9	17
1	B	282/397 (71%)	263 (93%)	17 (6%)	2 (1%)	18	32
All	All	674/794 (85%)	630 (94%)	37 (6%)	7 (1%)	12	23

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1151	LEU
1	A	1220	ALA
1	A	1228	THR
1	B	1021	SER
1	A	1021	SER
1	B	1150	GLY
1	A	1148	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	328/331 (99%)	311 (95%)	17 (5%)	21	40
1	B	242/331 (73%)	229 (95%)	13 (5%)	20	39
All	All	570/662 (86%)	540 (95%)	30 (5%)	20	39

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	876	ILE
1	A	877	VAL
1	A	881	GLU
1	A	887	THR
1	A	899	VAL
1	A	931	ARG
1	A	966	GLU
1	A	980	SER
1	A	982	HIS
1	A	986	SER
1	A	1051	GLN
1	A	1060	GLN
1	A	1186	SER
1	A	1202	THR
1	A	1254	TYR
1	A	1256	ASN
1	A	1259	GLN
1	B	887	THR
1	B	891	THR
1	B	916	GLN
1	B	966	GLU
1	B	980	SER
1	B	982	HIS
1	B	986	SER
1	B	1045	ARG
1	B	1048	SER
1	B	1062	GLN
1	B	1096	GLN
1	B	1145	THR
1	B	1151	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1051	GLN
1	A	1054	ASN
1	A	1096	GLN
1	A	1199	GLN
1	A	1245	HIS
1	A	1256	ASN
1	A	1259	GLN
1	B	1114	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SIN	B	1301	-	7,7,7	1.06	0	8,8,8	1.01	0
2	SIN	A	1301	-	7,7,7	1.16	0	8,8,8	1.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SIN	B	1301	-	-	2/5/5/5	-
2	SIN	A	1301	-	-	4/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1301	SIN	O1-C1-C2-C3
2	A	1301	SIN	O1-C1-C2-C3
2	A	1301	SIN	O2-C1-C2-C3
2	B	1301	SIN	O2-C1-C2-C3
2	A	1301	SIN	C2-C3-C4-O4
2	A	1301	SIN	C2-C3-C4-O3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	392/397 (98%)	0.15	19 (4%) 35 32	29, 52, 105, 136	2 (0%)
1	B	282/397 (71%)	0.29	15 (5%) 32 29	30, 51, 108, 137	2 (0%)
All	All	674/794 (84%)	0.21	34 (5%) 34 31	29, 52, 108, 137	4 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	876	ILE	5.7
1	B	1145	THR	4.2
1	B	1151	LEU	4.0
1	A	1022	ALA	3.7
1	A	1220	ALA	3.7
1	A	1214	ALA	3.7
1	A	1221	ALA	3.4
1	A	1267	GLY	3.4
1	B	1019	LEU	3.1
1	A	876	ILE	3.0
1	A	1254	TYR	2.8
1	B	1020	SER	2.8
1	A	1149	SER	2.7
1	B	950	PRO	2.5
1	B	1132	TYR	2.5
1	B	1146	ARG	2.5
1	A	1019	LEU	2.5
1	A	1240	GLY	2.4
1	A	1264	HIS	2.4
1	A	889	GLY	2.4
1	B	982	HIS	2.4
1	B	974	GLY	2.3
1	B	1026	ARG	2.3
1	A	1151	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1253	TYR	2.3
1	B	1147	VAL	2.2
1	B	1022	ALA	2.2
1	A	972	ALA	2.2
1	A	1024	LEU	2.2
1	A	1217	ALA	2.2
1	B	891	THR	2.2
1	A	1020	SER	2.1
1	B	933	CYS	2.0
1	A	1150	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SIN	B	1301	8/8	0.89	0.21	89,94,102,113	0
2	SIN	A	1301	8/8	0.90	0.18	80,93,106,107	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around SIN B 1301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.