



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:48 PM UTC

PDB ID : 1I6L / pdb_00001i6l
Title : 1.7 HIGH RESOLUTION EXPERIMENTAL PHASES FOR
TRYPTOPHANYL-TRNA SYNTHETASE COMPLEXED WITH
TRYPTOPHANYL-5'AMP
Authors : Retailleau, P.; Carter, C.W.
Deposited on : 2001-03-02
Resolution : 1.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	364	-	-	X	-
2	SO4	A	366	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	370	-	X	-	-
5	GOL	A	371	-	X	-	-
5	GOL	A	372	-	X	-	-
5	GOL	A	373	-	X	-	-
5	GOL	A	374	-	X	-	-
5	GOL	A	375	-	X	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3064 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRYPTOPHANYL-TRNA SYNTHETASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	Se	0	5	0
			2604	1649	449	493	3	10			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	modified residue	UNP P00953
A	92	MSE	MET	modified residue	UNP P00953
A	105	MSE	MET	modified residue	UNP P00953
A	129	MSE	MET	modified residue	UNP P00953
A	184	MSE	MET	modified residue	UNP P00953
A	193	MSE	MET	modified residue	UNP P00953
A	291	MSE	MET	modified residue	UNP P00953
A	314	MSE	MET	modified residue	UNP P00953
A	318	MSE	MET	modified residue	UNP P00953
A	322	MSE	MET	modified residue	UNP P00953

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is AMMONIUM ION (CCD ID: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	N	0	0
			1	1		

- Molecule 4 is TRYPTOPHANYL-5'AMP (CCD ID: TYM) (formula: C₂₁H₂₄N₇O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			37	21	7	8	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	351	Total	O	0	0
			351	351		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRYPTOPHANYL-TRNA SYNTHETASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	59.77Å 59.77Å 232.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.72 20.00 – 1.72	Depositor EDS
% Data completeness (in resolution range)	96.8 (20.00-1.72) 96.7 (20.00-1.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.71Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER	Depositor
R, R_{free}	0.209 , 0.241 0.194 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtriage
Anisotropy	0.492	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3064	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH4, SO4, GOL, TYM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.30	7/2670 (0.3%)	1.11	11/3592 (0.3%)

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	79	ILE	CA-CB	-7.83	1.45	1.54
1	A	88	GLN	N-CA	7.15	1.54	1.46
1	A	171	ILE	CA-C	6.86	1.58	1.53
1	A	115	LYS	CA-C	-5.70	1.45	1.53
1	A	17	GLY	CA-C	5.55	1.58	1.52
1	A	180	GLY	CA-C	5.33	1.58	1.51
1	A	22	ALA	CA-CB	5.31	1.60	1.53

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	117	ALA	N-CA-C	-9.04	91.75	107.49
1	A	34	ASN	N-CA-C	-7.68	97.54	109.76
1	A	126	PRO	N-CA-C	6.40	118.51	110.70
1	A	138	THR	N-CA-C	6.04	119.09	110.10
1	A	17	GLY	N-CA-C	-5.75	105.40	112.77
1	A	120	ALA	N-CA-C	5.68	118.24	111.71
1	A	94[A]	GLN	N-CA-C	-5.65	105.21	111.71
1	A	94[B]	GLN	N-CA-C	-5.65	105.21	111.71
1	A	30	GLN	N-CA-C	5.33	118.57	111.75
1	A	190	THR	N-CA-C	-5.10	106.84	113.16
1	A	164	ARG	N-CA-C	5.04	116.85	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2604	0	2621	44	0
2	A	35	0	0	4	0
3	A	1	0	0	0	0
4	A	37	0	23	1	0
5	A	36	0	24	2	0
6	A	351	0	0	16	0
All	All	3064	0	2668	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:195[A]:LYS:HG2	5:A:374:GOL:H12	1.66	0.77
1:A:201:LYS:HG3	6:A:578:HOH:O	1.86	0.74
1:A:14:ILE:HD13	1:A:195[B]:LYS:HE2	1.73	0.69
1:A:107:GLN:HG2	6:A:467:HOH:O	1.95	0.66
1:A:228:ILE:HD12	1:A:260:TYR:HB3	1.77	0.65
1:A:285:GLU:HG2	6:A:655:HOH:O	1.98	0.64
1:A:107:GLN:HG3	1:A:150:HIS:CE1	2.33	0.63
1:A:86:HIS:HD2	1:A:132:ASP:OD2	1.82	0.62
1:A:214:GLU:O	1:A:218:LYS:HD3	2.00	0.62
1:A:112:SER:O	1:A:115:LYS:HB2	2.00	0.61
1:A:111:LYS:HD3	6:A:675:HOH:O	2.00	0.61
1:A:225:GLU:OE2	1:A:227:THR:HB	2.00	0.61
1:A:288:HIS:HE1	6:A:562:HOH:O	1.87	0.57
1:A:288:HIS:HD2	6:A:549:HOH:O	1.88	0.57
1:A:125:TYR:HB3	1:A:126:PRO:HD3	1.88	0.56
1:A:294:GLU:HG3	6:A:714:HOH:O	2.07	0.53
1:A:229:ARG:HB2	6:A:741:HOH:O	2.10	0.52
1:A:125:TYR:OH	4:A:350:TYM:NH3	2.44	0.51
1:A:86:HIS:HE1	1:A:136:TYR:OH	1.93	0.51
1:A:182:ARG:HD2	6:A:680:HOH:O	2.13	0.49
1:A:60:ARG:NH1	6:A:687:HOH:O	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ARG:NH2	1:A:291:MSE:HE2	2.28	0.48
1:A:124:THR:O	1:A:127:PRO:HD2	2.13	0.48
1:A:182:ARG:HG3	6:A:680:HOH:O	2.12	0.48
1:A:4:ILE:HD12	1:A:33:TYR:CG	2.50	0.47
1:A:60:ARG:NH2	2:A:366:SO4:S	2.89	0.45
1:A:211:LYS:NZ	6:A:588:HOH:O	2.50	0.45
1:A:259:GLN:O	1:A:263:LYS:HE3	2.17	0.44
5:A:373:GOL:H11	6:A:551:HOH:O	2.17	0.44
1:A:119[A]:SER:HB2	6:A:436:HOH:O	2.17	0.44
1:A:232:LYS:HE3	2:A:364:SO4:O1	2.18	0.44
1:A:252:SER:OG	1:A:255:GLU:HG3	2.18	0.44
1:A:218:LYS:HA	1:A:269:LYS:HD3	2.00	0.43
1:A:211:LYS:HG3	6:A:727:HOH:O	2.18	0.43
1:A:201:LYS:NZ	1:A:219:SER:OG	2.46	0.43
1:A:211:LYS:HB3	1:A:211:LYS:HE3	1.81	0.43
1:A:164:ARG:HD2	6:A:559:HOH:O	2.19	0.42
1:A:159:GLU:HG3	1:A:171:ILE:HD12	2.02	0.42
1:A:11:SER:OG	1:A:195[B]:LYS:HD3	2.20	0.42
1:A:287:TYR:OH	2:A:366:SO4:O3	2.29	0.41
1:A:6:SER:HB3	1:A:26:PHE:CZ	2.55	0.41
1:A:285:GLU:OE2	1:A:285:GLU:HA	2.20	0.41
1:A:232:LYS:NZ	2:A:364:SO4:O4	2.53	0.41
1:A:274:GLN:HG3	1:A:278:GLU:OE2	2.21	0.40
1:A:122:LEU:HD23	1:A:122:LEU:HA	1.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/328 (100%)	318 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	283/270 (105%)	269 (95%)	14 (5%)	22 4

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MSE
1	A	4	ILE
1	A	32	GLU
1	A	109	LYS
1	A	115	LYS
1	A	116	GLU
1	A	164	ARG
1	A	201	LYS
1	A	205	THR
1	A	218	LYS
1	A	225	GLU
1	A	254	GLU
1	A	285	GLU
1	A	295	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	56	GLN
1	A	86	HIS
1	A	274	GLN
1	A	288	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 1 is modelled with single atom - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	372	-	5,5,5	4.75	5 (100%)	5,5,5	6.13	3 (60%)
2	SO4	A	366	-	4,4,4	1.61	1 (25%)	6,6,6	0.69	0
5	GOL	A	370	-	5,5,5	4.59	5 (100%)	5,5,5	5.83	3 (60%)
2	SO4	A	365	-	4,4,4	1.52	1 (25%)	6,6,6	0.70	0
2	SO4	A	362	-	4,4,4	1.56	1 (25%)	6,6,6	0.68	0
2	SO4	A	363	-	4,4,4	1.54	1 (25%)	6,6,6	0.67	0
5	GOL	A	371	-	5,5,5	4.78	4 (80%)	5,5,5	5.78	3 (60%)
5	GOL	A	373	-	5,5,5	4.76	5 (100%)	5,5,5	6.16	3 (60%)
5	GOL	A	374	-	5,5,5	4.98	5 (100%)	5,5,5	6.16	3 (60%)
5	GOL	A	375	-	5,5,5	4.70	5 (100%)	5,5,5	6.16	3 (60%)
2	SO4	A	361	-	4,4,4	1.56	1 (25%)	6,6,6	0.66	0
2	SO4	A	364	-	4,4,4	1.56	1 (25%)	6,6,6	0.66	0
4	TYM	A	350	-	40,41,41	1.84	7 (17%)	58,61,61	1.67	12 (20%)
2	SO4	A	360	-	4,4,4	1.51	1 (25%)	6,6,6	0.61	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	370	-	-	2/4/4/4	-
5	GOL	A	371	-	-	2/4/4/4	-
5	GOL	A	373	-	-	2/4/4/4	-
5	GOL	A	374	-	-	3/4/4/4	-
5	GOL	A	375	-	-	3/4/4/4	-
4	TYM	A	350	-	-	0/21/39/39	0/5/5/5
5	GOL	A	372	-	-	2/4/4/4	-

All (43) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	371	GOL	C3-C2	-8.71	1.18	1.51
5	A	374	GOL	C3-C2	-8.31	1.20	1.51
4	A	350	TYM	P-OPP	8.20	1.75	1.60
5	A	372	GOL	C3-C2	-7.93	1.21	1.51
5	A	373	GOL	C3-C2	-7.93	1.21	1.51
5	A	370	GOL	C3-C2	-7.84	1.21	1.51
5	A	375	GOL	C3-C2	-7.75	1.22	1.51
5	A	374	GOL	O1-C1	4.84	1.62	1.42
5	A	375	GOL	O1-C1	4.60	1.61	1.42
5	A	372	GOL	O1-C1	4.53	1.61	1.42
5	A	373	GOL	O1-C1	4.38	1.60	1.42
5	A	371	GOL	O1-C1	4.32	1.60	1.42
5	A	370	GOL	O1-C1	4.21	1.60	1.42
5	A	374	GOL	C1-C2	-3.76	1.37	1.51
5	A	375	GOL	O3-C3	3.56	1.57	1.42
5	A	373	GOL	O3-C3	3.41	1.56	1.42
5	A	372	GOL	O3-C3	3.33	1.56	1.42
4	A	350	TYM	P-O5'	3.30	1.72	1.59
5	A	373	GOL	C1-C2	-3.29	1.39	1.51
5	A	374	GOL	O3-C3	3.29	1.56	1.42
5	A	371	GOL	C1-C2	-3.26	1.39	1.51
5	A	370	GOL	C1-C2	-3.19	1.39	1.51
5	A	372	GOL	C1-C2	-3.19	1.39	1.51
5	A	375	GOL	O2-C2	-3.00	1.34	1.43
5	A	373	GOL	O2-C2	-3.00	1.34	1.43
5	A	370	GOL	O2-C2	-2.94	1.34	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	372	GOL	O2-C2	-2.89	1.35	1.43
4	A	350	TYM	CD2-CG	-2.83	1.39	1.44
5	A	370	GOL	O3-C3	2.74	1.53	1.42
5	A	375	GOL	C1-C2	-2.73	1.41	1.51
2	A	366	SO4	O1-S	2.61	1.60	1.44
5	A	374	GOL	O2-C2	-2.54	1.36	1.43
2	A	365	SO4	O1-S	2.54	1.59	1.44
2	A	362	SO4	O1-S	2.53	1.59	1.44
2	A	363	SO4	O1-S	2.51	1.59	1.44
2	A	364	SO4	O1-S	2.51	1.59	1.44
2	A	361	SO4	O1-S	2.50	1.59	1.44
5	A	371	GOL	O2-C2	-2.50	1.36	1.43
2	A	360	SO4	O1-S	2.39	1.59	1.44
4	A	350	TYM	C3'-C4'	2.35	1.59	1.53
4	A	350	TYM	CB-CG	2.15	1.54	1.50
4	A	350	TYM	C2'-C1'	2.14	1.60	1.53
4	A	350	TYM	CZ2-CE2	2.09	1.43	1.39

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	373	GOL	O3-C3-C2	11.23	160.93	110.38
5	A	375	GOL	O3-C3-C2	11.19	160.74	110.38
5	A	374	GOL	O3-C3-C2	11.17	160.67	110.38
5	A	372	GOL	O3-C3-C2	11.12	160.46	110.38
5	A	371	GOL	O3-C3-C2	10.47	157.52	110.38
5	A	370	GOL	O3-C3-C2	9.88	154.84	110.38
5	A	370	GOL	O2-C2-C3	7.69	141.03	109.18
5	A	374	GOL	O2-C2-C3	7.31	139.44	109.18
5	A	372	GOL	O2-C2-C3	7.28	139.31	109.18
5	A	373	GOL	O2-C2-C3	7.20	138.97	109.18
5	A	375	GOL	O2-C2-C3	7.09	138.52	109.18
5	A	371	GOL	O2-C2-C3	6.82	137.42	109.18
4	A	350	TYM	CD2-CG-CD1	5.94	112.24	106.16
4	A	350	TYM	CG-CD1-NE1	-4.87	104.96	110.31
4	A	350	TYM	CE2-CD2-CG	-4.57	102.55	107.17
5	A	375	GOL	O1-C1-C2	3.71	127.07	110.38
4	A	350	TYM	OPP-P-O5'	-3.47	92.59	103.00
5	A	373	GOL	O1-C1-C2	3.41	125.73	110.38
5	A	374	GOL	O1-C1-C2	3.34	125.42	110.38
5	A	370	GOL	O1-C1-C2	3.29	125.19	110.38
5	A	372	GOL	O1-C1-C2	3.28	125.13	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	371	GOL	O1-C1-C2	3.17	124.64	110.38
4	A	350	TYM	CB-CG-CD1	-2.96	121.20	126.97
4	A	350	TYM	CZ2-CE2-CD2	-2.71	119.56	122.19
4	A	350	TYM	CB-CA-C	-2.55	102.97	110.46
4	A	350	TYM	CE3-CD2-CG	2.44	137.50	133.85
4	A	350	TYM	O1P-P-O2P	2.34	123.32	112.44
4	A	350	TYM	O4'-C1'-C2'	-2.33	101.62	106.62
4	A	350	TYM	CD2-CE2-NE1	2.14	109.31	107.52
4	A	350	TYM	CE2-NE1-CD1	2.09	110.93	109.08

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	370	GOL	C1-C2-C3-O3
5	A	372	GOL	C1-C2-C3-O3
5	A	373	GOL	O1-C1-C2-C3
5	A	373	GOL	C1-C2-C3-O3
5	A	374	GOL	O1-C1-C2-C3
5	A	374	GOL	C1-C2-C3-O3
5	A	375	GOL	C1-C2-C3-O3
5	A	371	GOL	O2-C2-C3-O3
5	A	372	GOL	O1-C1-C2-O2
5	A	375	GOL	O1-C1-C2-C3
5	A	371	GOL	O1-C1-C2-O2
5	A	370	GOL	O1-C1-C2-O2
5	A	374	GOL	O1-C1-C2-O2
5	A	375	GOL	O1-C1-C2-O2


There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	366	SO4	2	0
5	A	373	GOL	1	0
5	A	374	GOL	1	0
2	A	364	SO4	2	0
4	A	350	TYM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	316/328 (96%)	0.23	24 (7%)	20 25	10, 19, 51, 79	5 (1%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	114	GLY	4.9
1	A	113	ALA	4.8
1	A	115	LYS	4.6
1	A	224	SER	3.8
1	A	285	GLU	3.8
1	A	226	GLY	3.7
1	A	117	ALA	3.4
1	A	111	LYS	3.4
1	A	116	GLU	3.1
1	A	31[A]	HIS	3.0
1	A	104	ARG	2.9
1	A	12	GLY	2.7
1	A	118	VAL	2.7
1	A	227	THR	2.6
1	A	262	GLY	2.6
1	A	225	GLU	2.5
1	A	228	ILE	2.4
1	A	263	LYS	2.3
1	A	178	LYS	2.1
1	A	218	LYS	2.1
1	A	32	GLU	2.1
1	A	258	ARG	2.0
1	A	211	LYS	2.0
1	A	107	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	372	6/6	0.48	0.40	32,37,41,46	6
2	SO4	A	363	5/5	0.56	0.28	47,51,52,52	5
5	GOL	A	375	6/6	0.62	0.19	47,52,57,62	0
2	SO4	A	365	5/5	0.65	0.28	40,45,45,45	5
5	GOL	A	373	6/6	0.70	0.20	46,51,56,61	0
2	SO4	A	364	5/5	0.72	0.30	41,46,46,46	5
2	SO4	A	366	5/5	0.74	0.22	44,49,50,50	5
5	GOL	A	371	6/6	0.76	0.17	23,27,29,32	0
5	GOL	A	374	6/6	0.77	0.20	33,39,44,49	0
2	SO4	A	361	5/5	0.82	0.17	40,44,45,45	5
2	SO4	A	362	5/5	0.85	0.15	34,38,39,39	5
2	SO4	A	360	5/5	0.85	0.23	29,34,34,34	5
5	GOL	A	370	6/6	0.88	0.11	18,23,26,29	0
3	NH4	A	400	1/1	0.97	0.06	20,20,20,20	0
4	TYM	A	350	37/37	0.97	0.06	9,14,20,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.