



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:42 AM UTC

PDB ID : 3I68 / pdb_00003i68
Title : Plasmodium falciparum dihydroorotate dehydrogenase bound with triazologyrimidine-based inhibitor DSM2
Authors : Deng, X.; Phillips, M.A.
Deposited on : 2009-07-06
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3129 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydroorotate dehydrogenase homolog, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2999	1916	502	566	15			

There are 63 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	125	MET	-	expression tag	UNP Q08210
A	126	GLY	-	expression tag	UNP Q08210
A	127	SER	-	expression tag	UNP Q08210
A	128	SER	-	expression tag	UNP Q08210
A	129	HIS	-	expression tag	UNP Q08210
A	130	HIS	-	expression tag	UNP Q08210
A	131	HIS	-	expression tag	UNP Q08210
A	132	HIS	-	expression tag	UNP Q08210
A	133	HIS	-	expression tag	UNP Q08210
A	134	HIS	-	expression tag	UNP Q08210
A	135	SER	-	expression tag	UNP Q08210
A	136	SER	-	expression tag	UNP Q08210
A	137	GLY	-	expression tag	UNP Q08210
A	138	LEU	-	expression tag	UNP Q08210
A	139	VAL	-	expression tag	UNP Q08210
A	140	PRO	-	expression tag	UNP Q08210
A	141	ARG	-	expression tag	UNP Q08210
A	142	GLY	-	expression tag	UNP Q08210
A	143	SER	-	expression tag	UNP Q08210
A	144	HIS	-	expression tag	UNP Q08210
A	145	MET	-	expression tag	UNP Q08210
A	146	ALA	-	expression tag	UNP Q08210
A	147	SER	-	expression tag	UNP Q08210
A	148	MET	-	expression tag	UNP Q08210
A	149	THR	-	expression tag	UNP Q08210
A	150	GLY	-	expression tag	UNP Q08210
A	151	GLY	-	expression tag	UNP Q08210

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	152	GLN	-	expression tag	UNP Q08210
A	153	GLN	-	expression tag	UNP Q08210
A	154	GLY	-	expression tag	UNP Q08210
A	155	ARG	-	expression tag	UNP Q08210
A	156	ASP	-	expression tag	UNP Q08210
A	157	PRO	-	expression tag	UNP Q08210
A	?	-	SER	deletion	UNP Q08210
A	?	-	THR	deletion	UNP Q08210
A	?	-	TYR	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	GLU	deletion	UNP Q08210
A	?	-	ASP	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	LYS	deletion	UNP Q08210
A	?	-	ILE	deletion	UNP Q08210
A	?	-	VAL	deletion	UNP Q08210
A	?	-	GLU	deletion	UNP Q08210
A	?	-	LYS	deletion	UNP Q08210
A	?	-	LYS	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	PHE	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	LYS	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210
A	?	-	SER	deletion	UNP Q08210
A	?	-	HIS	deletion	UNP Q08210
A	?	-	MET	deletion	UNP Q08210
A	?	-	MET	deletion	UNP Q08210
A	?	-	LYS	deletion	UNP Q08210
A	?	-	ASP	deletion	UNP Q08210
A	?	-	ALA	deletion	UNP Q08210
A	?	-	LYS	deletion	UNP Q08210
A	?	-	ASP	deletion	UNP Q08210
A	?	-	ASN	deletion	UNP Q08210


- Molecule 2 is N-anthracen-2-yl-5-methyl[1,2,4]triazolo[1,5-a]pyrimidin-7-amine (CCD ID: J4Z) (formula: C₂₀H₁₅N₅).



-
- The chemical structure of FMN (Flavin Mononucleotide) is shown. It consists of an isoalloxazine ring system (a fused bicyclic system with two nitrogen atoms, N1 and N5) attached to a ribitol chain. The ribitol chain is a five-carbon chain (C1' to C5') with hydroxyl groups at C1', C2', C3', and C4'. The C5' carbon is attached to a phosphate group (O1P, O2P, O3P, O4P). The ribitol chain is shown in a specific conformation with stereochemistry indicated by wedges and dashes. The isoalloxazine ring system is shown with atoms labeled C1, C2, C3, C4, C5, C6, C7, C8, C9, C10, N1, N5, and N10. The ribitol chain is shown with atoms labeled C1', C2', C3', C4', and C5'. The phosphate group is shown with atoms labeled O1P, O2P, O3P, and O4P.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 31	C 17	N 4	O 9	P 1	0	0

- Molecule 4 is OROTIC ACID (CCD ID: ORO) (formula: $\text{C}_5\text{H}_4\text{N}_2\text{O}_4$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	5	2	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	63	Total	O	0	0
			63	63		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase homolog, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	85.92Å 85.92Å 138.70Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.00 – 2.40 50.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (50.00-2.40) 99.8 (50.00-2.40)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.214 , 0.267 0.232 , 0.267	Depositor DCC
R_{free} test set	1477 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	55.6	Xtriage
Anisotropy	0.368	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.053 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3129	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ORO, J4Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/3050	1.01	10/4107 (0.2%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	353	ASP	N-CA-C	-12.17	93.80	110.35
1	A	414	PHE	N-CA-C	10.21	122.48	111.36
1	A	346	PRO	N-CA-C	-9.25	97.52	111.57
1	A	319	VAL	CB-CA-C	-8.21	101.28	111.87
1	A	234	ILE	N-CA-C	7.84	117.95	110.42
1	A	344	SER	N-CA-C	7.67	122.84	113.17
1	A	319	VAL	N-CA-C	6.49	116.52	110.42
1	A	163	PRO	N-CA-C	-6.32	104.93	113.40
1	A	319	VAL	N-CA-CB	-6.16	103.69	110.65
1	A	354	ASN	N-CA-C	5.29	117.85	109.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2999	0	3037	77	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	25	0	15	1	0
3	A	31	0	19	1	0
4	A	11	0	3	2	0
5	A	63	0	0	2	0
All	All	3129	0	3074	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:ARG:HE	1:A:354:ASN:HA	0.99	1.11
1:A:351:LEU:HD12	1:A:352:ARG:N	1.66	1.09
1:A:352:ARG:HG2	1:A:353:ASP:O	1.53	1.08
1:A:319:VAL:HG22	1:A:320:ASN:N	1.68	1.06
1:A:253:ARG:HD3	1:A:254:GLY:H	1.18	1.06
1:A:351:LEU:HD12	1:A:351:LEU:C	1.94	0.93
1:A:352:ARG:NE	1:A:354:ASN:HA	1.84	0.91
1:A:160:SER:OG	1:A:161:TYR:HA	1.72	0.89
1:A:346:PRO:O	1:A:347:ASN:HB2	1.73	0.87
1:A:230:ASN:HB3	1:A:281:MET:HE2	1.58	0.85
1:A:159:GLU:N	1:A:201:THR:H	1.77	0.82
1:A:319:VAL:HG13	1:A:320:ASN:H	1.46	0.80
1:A:537:LYS:HE2	1:A:541:GLN:HE22	1.47	0.80
1:A:320:ASN:HD22	1:A:323:ASP:CG	1.90	0.80
1:A:253:ARG:HD3	1:A:254:GLY:N	1.97	0.78
1:A:346:PRO:O	1:A:347:ASN:CB	2.30	0.78
1:A:320:ASN:HB3	1:A:323:ASP:OD2	1.85	0.76
1:A:352:ARG:HH21	1:A:354:ASN:HB3	1.51	0.75
1:A:191:LEU:HD22	1:A:196:ILE:HD11	1.69	0.74
1:A:162:ASN:HB2	1:A:165:PHE:HD1	1.54	0.73
1:A:356:GLU:C	1:A:356:GLU:OE1	2.33	0.71
1:A:482:LYS:HE2	5:A:39:HOH:O	1.90	0.70
1:A:537:LYS:HE2	1:A:541:GLN:NE2	2.09	0.68
1:A:351:LEU:C	1:A:351:LEU:CD1	2.68	0.66
1:A:356:GLU:OE1	1:A:358:GLY:N	2.30	0.65
1:A:320:ASN:ND2	1:A:323:ASP:OD1	2.30	0.63
1:A:351:LEU:HD12	1:A:352:ARG:CA	2.29	0.62
1:A:354:ASN:N	1:A:354:ASN:OD1	2.30	0.62
1:A:160:SER:OG	1:A:161:TYR:CA	2.48	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:346:PRO:HA	1:A:351:LEU:HD13	1.83	0.61
1:A:356:GLU:OE1	1:A:359:LYS:N	2.30	0.61
1:A:329:ILE:HD11	1:A:367:VAL:HG13	1.83	0.61
1:A:175:CYS:HB3	1:A:184:CYS:SG	2.41	0.60
1:A:159:GLU:HB2	1:A:199:TYR:O	2.03	0.59
1:A:351:LEU:CD1	1:A:352:ARG:N	2.55	0.57
1:A:320:ASN:ND2	1:A:323:ASP:CG	2.61	0.56
1:A:159:GLU:N	1:A:201:THR:N	2.51	0.55
1:A:463:ILE:HD13	1:A:480:LYS:HB3	1.89	0.55
1:A:352:ARG:O	1:A:352:ARG:CD	2.55	0.55
1:A:235:ASP:OD2	1:A:296:ARG:NH1	2.41	0.53
1:A:319:VAL:HG22	1:A:320:ASN:CB	2.38	0.53
1:A:162:ASN:ND2	5:A:62:HOH:O	2.36	0.52
1:A:457:SER:N	1:A:505:SER:O	2.33	0.51
1:A:321:ILE:HG23	1:A:322:VAL:N	2.25	0.51
1:A:414:PHE:CD1	1:A:414:PHE:N	2.79	0.51
1:A:454:MET:HE2	1:A:500:ILE:HB	1.94	0.50
1:A:244:PHE:HB3	1:A:307:ILE:HB	1.93	0.50
1:A:252:PRO:HG2	1:A:324:ASP:OD1	2.11	0.50
1:A:226:GLY:HA3	3:A:1002:FMN:N5	2.26	0.50
1:A:346:PRO:O	1:A:346:PRO:CD	2.56	0.50
1:A:240:LEU:HD11	2:A:1001:J4Z:H25	1.93	0.50
1:A:356:GLU:OE1	1:A:356:GLU:O	2.30	0.48
1:A:343:VAL:HA	1:A:353:ASP:HB2	1.96	0.48
1:A:319:VAL:HG22	1:A:320:ASN:CA	2.43	0.48
1:A:160:SER:HA	1:A:161:TYR:C	2.39	0.47
1:A:319:VAL:HG13	1:A:320:ASN:N	2.23	0.47
1:A:345:SER:HB2	4:A:1003:ORO:C4	2.44	0.47
1:A:348:THR:O	1:A:351:LEU:O	2.32	0.47
1:A:359:LYS:HA	1:A:359:LYS:HD3	1.46	0.47
1:A:159:GLU:O	1:A:160:SER:HB3	2.15	0.47
1:A:345:SER:OG	1:A:346:PRO:O	2.31	0.46
1:A:382:ASP:O	1:A:383:GLU:C	2.56	0.46
1:A:345:SER:HB2	4:A:1003:ORO:O4	2.16	0.45
1:A:352:ARG:HG2	1:A:353:ASP:C	2.35	0.45
1:A:159:GLU:HG3	1:A:199:TYR:O	2.17	0.44
1:A:274:ASN:O	1:A:474:GLY:HA3	2.17	0.44
1:A:319:VAL:H	1:A:319:VAL:HG12	1.34	0.43
1:A:528:TYR:CD2	1:A:528:TYR:C	2.96	0.43
1:A:169:ASP:O	1:A:173:LYS:HG3	2.18	0.43
1:A:179:ILE:HB	1:A:184:CYS:SG	2.60	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:329:ILE:CD1	1:A:367:VAL:HG13	2.47	0.42
1:A:352:ARG:HG3	1:A:355:GLN:HE21	1.83	0.42
1:A:326:LYS:HG2	1:A:370:GLU:HG3	2.02	0.42
1:A:222:GLY:HA3	1:A:244:PHE:CE2	2.55	0.41
1:A:162:ASN:HA	1:A:163:PRO:HD2	1.81	0.41
1:A:255:GLN:OE1	1:A:315:ASN:HA	2.21	0.41
1:A:192:GLY:HA3	1:A:233:CYS:SG	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	376/415 (91%)	361 (96%)	13 (4%)	2 (0%)	24 37

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	SER
1	A	347	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/365 (92%)	321 (96%)	14 (4%)	26	45

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	162	ASN
1	A	167	LEU
1	A	235	ASP
1	A	253	ARG
1	A	316	LYS
1	A	318	THR
1	A	319	VAL
1	A	351	LEU
1	A	352	ARG
1	A	354	ASN
1	A	356	GLU
1	A	383	GLU
1	A	487	LYS
1	A	524	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	289	ASN
1	A	320	ASN
1	A	330	ASN
1	A	355	GLN
1	A	450	ASN
1	A	541	GLN
1	A	548	HIS
1	A	566	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FMN	A	1002	-	33,33,33	1.07	2 (6%)	48,50,50	1.30	6 (12%)
4	ORO	A	1003	-	11,11,11	1.52	2 (18%)	14,15,15	2.39	6 (42%)
2	J4Z	A	1001	-	29,29,29	1.74	7 (24%)	37,42,42	2.14	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMN	A	1002	-	-	1/18/18/18	0/3/3/3
4	ORO	A	1003	-	-	4/4/4/4	0/1/1/1
2	J4Z	A	1001	-	-	0/4/4/4	0/5/5/5

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	J4Z	C15-N2	-3.95	1.32	1.38
3	A	1002	FMN	C4A-N5	3.75	1.38	1.30
2	A	1001	J4Z	C9-C4	3.38	1.49	1.42
2	A	1001	J4Z	C7-C6	3.27	1.49	1.42
3	A	1002	FMN	C10-N1	3.19	1.39	1.33
2	A	1001	J4Z	C11-N2	-2.82	1.34	1.37
2	A	1001	J4Z	C16-N3	2.42	1.36	1.32
2	A	1001	J4Z	C2-N1	-2.35	1.36	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	J4Z	C13-N5	2.35	1.36	1.33
4	A	1003	ORO	C4-N3	-2.20	1.34	1.38
4	A	1003	ORO	O72-C7	-2.15	1.24	1.30

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	J4Z	C16-N4-C15	8.83	108.26	102.36
4	A	1003	ORO	C4-N3-C2	-4.81	120.88	125.55
2	A	1001	J4Z	N3-C16-N4	-4.53	109.59	116.78
4	A	1003	ORO	O4-C4-C5	-4.13	119.76	125.46
4	A	1003	ORO	N3-C2-N1	3.29	120.92	115.74
4	A	1003	ORO	C5-C4-N3	3.26	119.29	115.24
2	A	1001	J4Z	C16-N3-N2	3.21	103.48	100.86
2	A	1001	J4Z	C14-C13-N5	3.10	120.29	116.84
2	A	1001	J4Z	C15-N2-N3	3.01	112.14	110.11
3	A	1002	FMN	C4A-C10-N10	3.00	120.78	116.48
2	A	1001	J4Z	N2-C15-N4	-3.00	106.51	109.19
3	A	1002	FMN	C9A-C5A-N5	-2.95	119.32	122.45
2	A	1001	J4Z	C12-C13-N5	-2.83	119.92	122.71
3	A	1002	FMN	C10-C4A-N5	-2.68	119.33	124.81
3	A	1002	FMN	C4-C4A-N5	2.39	121.50	118.21
3	A	1002	FMN	C5A-C9A-N10	2.34	120.08	117.97
4	A	1003	ORO	C7-C6-N1	2.27	119.62	115.47
4	A	1003	ORO	C6-N1-C2	-2.08	120.42	122.65
3	A	1002	FMN	C4A-C4-N3	2.03	118.41	113.25

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1003	ORO	N1-C6-C7-O71
4	A	1003	ORO	N1-C6-C7-O72
3	A	1002	FMN	C4'-C5'-O5'-P
4	A	1003	ORO	C5-C6-C7-O71
4	A	1003	ORO	C5-C6-C7-O72


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1002	FMN	1	0
4	A	1003	ORO	2	0
2	A	1001	J4Z	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand FMN A 1002



Ligand J4Z A 1001



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/415 (91%)	0.56	23 (6%) 27 23	42, 61, 83, 96	1 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	351	LEU	6.4
1	A	161	TYR	6.2
1	A	353	ASP	5.2
1	A	348	THR	4.1
1	A	356	GLU	3.7
1	A	160	SER	3.6
1	A	194	TYR	3.5
1	A	319	VAL	3.4
1	A	350	GLY	3.4
1	A	159	GLU	3.1
1	A	497	ASN	3.0
1	A	343	VAL	2.8
1	A	471	ASN	2.8
1	A	554	GLY	2.7
1	A	354	ASN	2.5
1	A	349	PRO	2.4
1	A	566	HIS	2.4
1	A	541	GLN	2.2
1	A	562	ILE	2.2
1	A	352	ARG	2.1
1	A	162	ASN	2.0
1	A	357	ALA	2.0
1	A	551	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ORO	A	1003	11/11	0.94	0.07	53,54,55,55	0
2	J4Z	A	1001	25/25	0.95	0.08	49,50,52,53	0
3	FMN	A	1002	31/31	0.96	0.07	42,48,51,52	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around J4Z A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FMN A 1002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.