



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 04:26 PM UTC

PDB ID : 1I7A / pdb_00001i7a
Title : EVH1 DOMAIN FROM MURINE HOMER 2B/VESL 2
Authors : Barzik, M.; Carl, U.D.; Schubert, W.-D.; Wehland, J.; Heinz, D.W.
Deposited on : 2001-03-08
Resolution : 2.24 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	111	71% 18% • 7%
1	B	111	67% 23% • 7%
1	C	111	60% 30% • 7%
1	D	111	66% 27% 7%
2	E	3	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	B	302	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3513 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HOMER 2B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	103	Total	C	N	O	S	0	0	0
			830	531	143	155	1			
1	B	103	Total	C	N	O	S	0	0	0
			830	531	143	155	1			
1	C	103	Total	C	N	O	S	17	0	0
			830	531	143	155	1			
1	D	103	Total	C	N	O	S	0	0	0
			830	531	143	155	1			

- Molecule 2 is a protein called PHE-ALA-PHE.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	3	Total	C	N	O	27	0	0
			27	21	3	3			

- Molecule 3 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is PHENYLALANINE (CCD ID: PHE) (formula: $C_9H_{11}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			11	9	1	1		
4	B	1	Total	C	N	O	0	0
			11	9	1	1		
4	D	1	Total	C	N	O	0	0
			12	9	1	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	25	Total	O	0	0
			25	25		
5	B	26	Total	O	0	0
			26	26		
5	C	29	Total	O	0	0
			29	29		
5	D	25	Total	O	0	0
			25	25		
5	E	1	Total	O	0	0
			1	1		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: HOMER 2B

Chain A: 




• Molecule 1: HOMER 2B

Chain B: 



• Molecule 1: HOMER 2B

Chain C: 



• Molecule 1: HOMER 2B

Chain D: 



• Molecule 2: PHE-ALA-PHE

Chain E: 

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.72Å 50.01Å 73.10Å 90.00° 101.19° 90.00°	Depositor
Resolution (Å)	15.00 – 2.24	Depositor
% Data completeness (in resolution range)	85.7 (15.00-2.24)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS, REFMAC	Depositor
R, R_{free}	0.235 , 0.296	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	3513	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/848	0.89	2/1146 (0.2%)
1	B	0.46	0/848	0.91	4/1146 (0.3%)
1	C	0.42	0/848	0.88	1/1146 (0.1%)
1	D	0.42	0/848	0.85	2/1146 (0.2%)
2	E	0.35	0/28	0.58	0/36
All	All	0.45	0/3420	0.88	9/4620 (0.2%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	62	THR	CA-C-N	7.03	126.85	119.05
1	B	62	THR	C-N-CA	7.03	126.85	119.05
1	B	54	LYS	N-CA-C	6.11	119.15	109.50
1	A	59	SER	N-CA-C	5.88	118.11	108.52
1	C	59	SER	N-CA-C	5.64	118.15	109.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	830	0	817	23	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	830	0	817	17	3
1	C	830	0	817	34	0
1	D	830	0	817	29	2
2	E	27	0	22	0	0
3	A	13	0	5	0	0
3	B	13	0	5	0	0
4	A	11	0	8	0	0
4	B	11	0	8	0	0
4	D	12	0	8	4	0
5	A	25	0	0	2	0
5	B	26	0	0	1	0
5	C	29	0	0	2	0
5	D	25	0	0	3	0
5	E	1	0	0	0	0
All	All	3513	0	3324	100	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 100 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:GLN:NE2	1:A:107:ARG:HH21	1.50	1.10
1:A:104:GLN:HE22	1:A:107:ARG:NH2	1.56	1.03
1:C:46:ARG:HD3	1:C:58:ASN:ND2	1.84	0.93
1:A:46:ARG:HD3	1:A:58:ASN:ND2	1.89	0.88
1:A:104:GLN:HE22	1:A:107:ARG:HH21	0.82	0.79

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:98:LYS:NZ	1:B:111:ARG:O[2_546]	2.09	0.11
1:B:58:ASN:ND2	1:D:58:ASN:OD1[2_556]	2.16	0.04
1:B:58:ASN:OD1	1:D:58:ASN:ND2[2_556]	2.16	0.04

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/111 (89%)	93 (94%)	6 (6%)	0	100	100
1	B	99/111 (89%)	95 (96%)	4 (4%)	0	100	100
1	C	99/111 (89%)	91 (92%)	6 (6%)	2 (2%)	6	2
1	D	99/111 (89%)	93 (94%)	6 (6%)	0	100	100
2	E	1/3 (33%)	0	1 (100%)	0	100	100
All	All	397/447 (89%)	372 (94%)	23 (6%)	2 (0%)	24	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	24	TRP
1	C	108	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/97 (93%)	85 (94%)	5 (6%)	19	18
1	B	90/97 (93%)	86 (96%)	4 (4%)	25	27
1	C	90/97 (93%)	87 (97%)	3 (3%)	33	39
1	D	90/97 (93%)	90 (100%)	0	100	100
2	E	2/2 (100%)	2 (100%)	0	100	100
All	All	362/390 (93%)	350 (97%)	12 (3%)	33	39

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	72	GLN
1	B	94	LEU
1	C	69	LYS
1	C	25	VAL
1	A	81	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	30	GLN
1	D	72	GLN
1	C	12	HIS
1	C	15	GLN
1	C	23	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PHE	D	401	-	11,12,12	0.71	0	11,15,15	0.26	0
3	FLC	A	301	-	12,12,12	4.23	7 (58%)	17,17,17	1.43	4 (23%)
4	PHE	B	402	-	10,11,12	0.57	0	8,13,15	0.21	0
3	FLC	B	302	-	12,12,12	4.16	7 (58%)	17,17,17	1.44	5 (29%)
4	PHE	A	403	-	10,11,12	0.72	0	8,13,15	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PHE	D	401	-	-	5/8/8/8	0/1/1/1
3	FLC	A	301	-	-	5/16/16/16	-
4	PHE	B	402	-	-	2/5/6/8	0/1/1/1
3	FLC	B	302	-	-	6/16/16/16	-
4	PHE	A	403	-	-	2/5/6/8	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	FLC	CG-CB	9.20	1.65	1.54
3	A	301	FLC	CG-CB	8.68	1.64	1.54
3	A	301	FLC	CA-CB	8.22	1.64	1.54
3	B	302	FLC	CA-CB	7.76	1.63	1.54
3	A	301	FLC	CB-CBC	6.52	1.60	1.53

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	FLC	CB-CG-CGC	2.65	121.18	113.92
3	A	301	FLC	OHB-CB-CG	-2.46	103.78	109.38
3	A	301	FLC	CB-CA-CAC	2.38	120.42	113.92
3	A	301	FLC	CG-CB-CBC	2.33	115.18	110.03
3	A	301	FLC	OB2-CBC-CB	2.25	117.46	113.14

There are no chirality outliers.

5 of 20 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	FLC	CG-CB-CBC-OB1
3	A	301	FLC	CG-CB-CBC-OB2
3	A	301	FLC	OHB-CB-CBC-OB1
3	A	301	FLC	OHB-CB-CBC-OB2
4	D	401	PHE	O-C-CA-N

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	401	PHE	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.