



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 04:08 AM UTC

PDB ID : 1I7R / pdb_00001i7r
Title : CRYSTAL STRUCTURE OF CLASS I MHC A2 IN COMPLEX WITH PEP-
TIDE P1058
Authors : Busslep, J.; Zhao, R.; Loftus, D.; Appella, E.; Collins, E.J.
Deposited on : 2001-03-10
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	5% 80% 16% .
1	D	275	7% 79% 16% 5%
2	B	100	2% 77% 18% . .
2	E	100	4% 75% 20% . .
3	C	9	22% 78% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	9	22% 56% 44%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			
1	D	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			

- Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	E	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	cloning artifact	UNP P01884
E	0	MET	-	cloning artifact	UNP P01884

- Molecule 3 is a protein called 9 RESIDUE PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O	0	0	0
			77	57	9	11			
3	F	9	Total	C	N	O	0	0	0
			76	57	9	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	34	Total 34	O 34	0	0
4	B	22	Total 22	O 22	0	0
4	C	1	Total 1	O 1	0	0
4	D	32	Total 32	O 32	0	0
4	E	17	Total 17	O 17	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN



- Molecule 2: BETA-2-MICROGLOBULIN



- Molecule 2: BETA-2-MICROGLOBULIN




● Molecule 3: 9 RESIDUE PEPTIDE

Chain C: 



● Molecule 3: 9 RESIDUE PEPTIDE

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.86Å 63.82Å 75.87Å 82.09° 76.20° 78.13°	Depositor
Resolution (Å)	30.00 – 2.20 30.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.7 (30.00-2.20) 96.7 (30.00-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.20Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.264 , 0.307 0.242 , 0.282	Depositor DCC
R_{free} test set	2252 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6427	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.58	0/2312	1.50	32/3137 (1.0%)
1	D	0.59	0/2312	1.51	29/3137 (0.9%)
2	B	0.60	0/860	1.51	8/1162 (0.7%)
2	E	0.58	0/860	1.47	5/1162 (0.4%)
3	C	0.71	0/82	1.65	1/110 (0.9%)
3	F	0.68	0/81	1.55	3/110 (2.7%)
All	All	0.59	0/6507	1.50	78/8818 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	E	0	1
All	All	0	2

There are no bond length outliers.

All (78) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	223	ASP	CA-CB-CG	11.81	124.41	112.60
1	A	223	ASP	CA-CB-CG	11.58	124.18	112.60
2	B	21	ASN	OD1-CG-ND2	-9.43	113.17	122.60
1	A	44	ARG	NE-CZ-NH1	-8.55	112.94	121.50
1	D	25	VAL	CA-C-N	8.29	127.29	121.65
1	D	25	VAL	C-N-CA	8.29	127.29	121.65
1	A	273	ARG	NE-CZ-NH1	-8.03	113.47	121.50
1	D	44	ARG	NE-CZ-NH2	7.56	126.00	119.20
2	B	97	ARG	NE-CZ-NH2	7.41	125.87	119.20
1	A	197	HIS	CA-CB-CG	-7.35	106.45	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	83	ASN	CA-CB-CG	-7.33	105.27	112.60
1	D	179	LEU	N-CA-C	7.23	120.21	111.82
1	A	25	VAL	CA-C-N	7.12	126.49	121.65
1	A	25	VAL	C-N-CA	7.12	126.49	121.65
3	C	1	PHE	CA-CB-CG	7.10	120.90	113.80
1	A	97	ARG	CD-NE-CZ	7.07	134.30	124.40
1	A	211	ALA	N-CA-C	7.05	118.96	111.28
2	E	83	ASN	CA-CB-CG	-6.97	105.63	112.60
1	A	9	PHE	CA-CB-CG	6.79	120.59	113.80
1	A	111	ARG	CA-C-N	6.75	126.81	121.61
1	A	111	ARG	C-N-CA	6.75	126.81	121.61
1	D	197	HIS	CA-CB-CG	-6.57	107.23	113.80
1	D	116	TYR	CA-CB-CG	6.54	125.68	113.90
1	D	44	ARG	NE-CZ-NH1	-6.44	115.06	121.50
1	D	97	ARG	CD-NE-CZ	6.44	133.41	124.40
1	D	55	GLU	N-CA-C	-6.30	101.62	110.50
1	D	141	GLN	OE1-CD-NE2	-6.20	116.40	122.60
1	A	273	ARG	NE-CZ-NH2	6.07	124.67	119.20
1	D	26	GLY	O-C-N	-6.04	118.66	123.49
1	D	9	PHE	CA-CB-CG	6.03	119.83	113.80
1	A	77	ASP	CA-C-O	-5.99	114.16	120.63
1	D	44	ARG	CD-NE-CZ	-5.97	116.04	124.40
1	D	102	ASP	CA-CB-CG	5.96	118.56	112.60
1	A	35	ARG	CD-NE-CZ	5.94	132.71	124.40
1	A	116	TYR	CA-CB-CG	5.89	124.51	113.90
1	A	234	ARG	NE-CZ-NH1	-5.73	115.77	121.50
1	D	211	ALA	N-CA-C	5.71	117.59	111.36
1	D	35	ARG	CD-NE-CZ	5.69	132.36	124.40
2	E	44	GLU	CB-CG-CD	5.66	122.22	112.60
2	B	56	PHE	CA-CB-CG	5.64	119.44	113.80
3	F	1	PHE	CA-CB-CG	5.60	119.40	113.80
1	A	32	GLN	CA-C-O	-5.59	114.67	121.05
1	D	219	ARG	CD-NE-CZ	5.59	132.23	124.40
1	A	84	TYR	CA-CB-CG	5.57	123.93	113.90
1	D	84	TYR	CA-CB-CG	5.55	123.89	113.90
1	A	234	ARG	O-C-N	5.53	125.25	121.71
1	D	52	ILE	CA-C-O	5.48	125.01	119.20
1	D	32	GLN	CA-C-O	-5.47	114.81	121.05
1	D	28	VAL	N-CA-CB	5.47	118.79	111.90
1	A	44	ARG	NE-CZ-NH2	5.47	124.12	119.20
1	A	149	ALA	CA-C-O	5.47	126.26	119.97
2	B	34	ASP	N-CA-CB	5.46	118.05	109.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4	THR	CA-C-O	5.46	124.62	119.76
1	D	256	ARG	CD-NE-CZ	5.37	131.91	124.40
2	E	33	SER	N-CA-C	5.36	119.54	112.89
2	B	44	GLU	CB-CG-CD	5.36	121.72	112.60
2	B	5	PRO	CB-CA-C	-5.36	104.98	111.46
1	D	218	GLN	OE1-CD-NE2	-5.35	117.25	122.60
1	A	52	ILE	CA-C-O	5.34	124.86	119.20
1	D	129	ASP	CA-CB-CG	5.34	117.94	112.60
1	A	244	TRP	CA-CB-CG	5.33	123.72	113.60
1	D	244	TRP	CA-CB-CG	5.33	123.72	113.60
1	A	219	ARG	CD-NE-CZ	5.25	131.76	124.40
1	A	112	GLY	O-C-N	-5.25	118.75	123.37
1	A	234	ARG	NE-CZ-NH2	5.24	123.92	119.20
1	D	77	ASP	CA-C-O	-5.22	115.01	120.55
2	B	2	GLN	OE1-CD-NE2	-5.21	117.39	122.60
2	E	34	ASP	N-CA-CB	5.18	117.65	109.83
1	A	204	TRP	N-CA-C	5.18	117.92	109.59
1	A	102	ASP	CA-CB-CG	5.17	117.77	112.60
1	A	26	GLY	O-C-N	-5.13	119.39	123.49
1	A	93	HIS	CA-CB-CG	-5.13	108.67	113.80
1	A	45	MET	CA-C-O	-5.11	115.44	121.16
3	F	2	ALA	CA-C-N	5.08	124.98	119.85
3	F	2	ALA	C-N-CA	5.08	124.98	119.85
1	A	28	VAL	N-CA-CB	5.05	118.27	111.90
1	D	111	ARG	CA-C-N	5.03	125.48	121.61
1	D	111	ARG	C-N-CA	5.03	125.48	121.61

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	0	MET	Mainchain
2	E	0	MET	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2096	26	0
1	D	2247	0	2096	25	0
2	B	837	0	803	11	0
2	E	837	0	803	12	0
3	C	77	0	71	1	0
3	F	76	0	71	1	0
4	A	34	0	0	1	0
4	B	22	0	0	2	0
4	C	1	0	0	0	0
4	D	32	0	0	1	0
4	E	17	0	0	1	0
All	All	6427	0	5940	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:0:MET:HG2	2:B:0:MET:O	1.67	0.92
2:B:73:THR:HG22	2:B:76:ASP:H	1.48	0.79
2:E:73:THR:HG22	2:E:76:ASP:H	1.49	0.77
1:D:19:GLU:HB3	1:D:20:PRO:HD2	1.71	0.73
1:A:19:GLU:HB3	1:A:20:PRO:HD2	1.70	0.73
2:B:85:VAL:HG13	4:B:100:HOH:O	1.97	0.65
2:B:12:ARG:HB3	2:B:21:ASN:HD21	1.62	0.65
1:D:99:TYR:OH	3:F:3:PRO:HD2	1.98	0.62
1:D:98:MET:HE3	1:D:98:MET:C	2.26	0.60
1:A:99:TYR:OH	3:C:3:PRO:HD2	2.02	0.59
1:A:253:GLN:NE2	1:A:256:ARG:HH11	2.00	0.59
1:A:98:MET:HE3	1:A:98:MET:C	2.28	0.58
1:A:263:HIS:HD2	1:A:265:GLY:H	1.51	0.58
2:B:13:HIS:HD2	2:B:21:ASN:ND2	2.02	0.58
1:D:263:HIS:HD2	1:D:265:GLY:H	1.52	0.57
1:A:98:MET:HE1	1:A:113:TYR:HB3	1.86	0.56
2:E:12:ARG:HB3	2:E:21:ASN:HD21	1.70	0.56
1:A:44:ARG:NH1	4:A:278:HOH:O	2.39	0.56
2:B:21:ASN:CG	2:B:22:PHE:H	2.14	0.55
1:D:98:MET:HE1	1:D:113:TYR:HB3	1.88	0.55
1:D:263:HIS:CD2	1:D:265:GLY:H	2.24	0.55
2:E:21:ASN:CG	2:E:22:PHE:H	2.16	0.54
1:A:98:MET:HE3	1:A:98:MET:O	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:0:MET:O	2:B:0:MET:CG	2.48	0.54
1:D:274:TRP:O	1:D:275:GLU:HB3	2.07	0.54
1:D:98:MET:HE3	1:D:98:MET:O	2.08	0.54
2:E:13:HIS:HD2	2:E:21:ASN:ND2	2.07	0.53
1:A:263:HIS:CD2	1:A:265:GLY:H	2.26	0.53
2:B:22:PHE:CZ	2:B:69:GLU:HG2	2.44	0.52
1:D:9:PHE:HE2	1:D:99:TYR:CE2	2.27	0.52
1:D:234:ARG:HD2	1:D:242:GLN:HB2	1.92	0.51
1:D:253:GLN:NE2	1:D:256:ARG:HH11	2.09	0.50
1:D:28:VAL:HG11	1:D:179:LEU:HD13	1.92	0.50
1:D:111:ARG:HD3	1:D:113:TYR:CZ	2.47	0.50
1:A:111:ARG:HD3	1:A:113:TYR:CZ	2.49	0.48
1:A:9:PHE:HE2	1:A:99:TYR:CE2	2.31	0.48
1:A:28:VAL:HG11	1:A:179:LEU:HD13	1.96	0.48
2:E:85:VAL:HG13	4:E:101:HOH:O	2.14	0.47
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.49	0.47
1:D:98:MET:HE2	1:D:115:GLN:HG2	1.97	0.47
2:E:21:ASN:ND2	2:E:22:PHE:H	2.14	0.46
1:D:44:ARG:HH11	1:D:44:ARG:HD3	1.44	0.46
1:D:127:LYS:HD2	1:D:132:SER:HB2	1.98	0.45
1:A:182:THR:CG2	1:A:265:GLY:HA2	2.47	0.45
2:B:21:ASN:ND2	2:B:22:PHE:H	2.15	0.45
1:A:273:ARG:HH11	1:A:273:ARG:HD2	1.48	0.45
1:A:44:ARG:HH11	1:A:44:ARG:HD3	1.47	0.44
1:D:15:PRO:HG2	1:D:91:GLY:O	2.18	0.44
2:E:83:ASN:OD1	2:E:90:PRO:HG3	2.17	0.44
1:D:16:GLY:O	1:D:17:ARG:NE	2.50	0.44
2:E:22:PHE:CZ	2:E:69:GLU:HG2	2.53	0.44
1:A:234:ARG:HD2	1:A:242:GLN:HB2	2.00	0.44
1:A:111:ARG:HD3	1:A:113:TYR:OH	2.18	0.43
1:D:253:GLN:HE21	1:D:256:ARG:HD3	1.83	0.43
2:E:51:HIS:HA	2:E:65:LEU:O	2.18	0.43
1:D:44:ARG:NH1	4:D:279:HOH:O	2.52	0.42
1:A:97:ARG:HD2	1:A:116:TYR:CZ	2.54	0.42
1:A:108:ARG:O	1:A:109:PHE:C	2.63	0.42
1:A:127:LYS:HD2	1:A:132:SER:HB2	2.02	0.42
1:A:197:HIS:ND1	1:A:197:HIS:N	2.68	0.42
1:D:55:GLU:OE2	1:D:170:ARG:NH2	2.52	0.42
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.54	0.42
1:D:184:ALA:HB2	1:D:265:GLY:O	2.21	0.41
2:E:73:THR:CG2	2:E:75:LYS:HG2	2.50	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:21:ASN:CG	4:B:120:HOH:O	2.64	0.41
1:D:97:ARG:HD2	1:D:116:TYR:CZ	2.56	0.41
1:D:111:ARG:HD3	1:D:113:TYR:OH	2.20	0.41
2:E:25:CYS:HB2	2:E:39:LEU:HD21	2.03	0.41
1:A:182:THR:HG23	1:A:265:GLY:HA2	2.02	0.40
1:A:257:TYR:O	1:A:273:ARG:HG3	2.22	0.40
1:A:98:MET:HE2	1:A:115:GLN:HG2	2.02	0.40
1:A:15:PRO:HG2	1:A:91:GLY:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	266 (97%)	7 (3%)	0	100	100
1	D	273/275 (99%)	267 (98%)	6 (2%)	0	100	100
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	E	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	756/768 (98%)	738 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	217 (94%)	14 (6%)	17	20
1	D	231/231 (100%)	217 (94%)	14 (6%)	17	20
2	B	95/95 (100%)	85 (90%)	10 (10%)	6	6
2	E	95/95 (100%)	86 (90%)	9 (10%)	8	8
3	C	7/7 (100%)	7 (100%)	0	100	100
3	F	7/7 (100%)	6 (86%)	1 (14%)	3	3
All	All	666/666 (100%)	618 (93%)	48 (7%)	13	15

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	SER
1	A	98	MET
1	A	105	SER
1	A	132	SER
1	A	176	LYS
1	A	182	THR
1	A	196	ASP
1	A	201	LEU
1	A	207	SER
1	A	223	ASP
1	A	225	THR
1	A	230	LEU
1	A	234	ARG
1	A	268	LYS
2	B	0	MET
2	B	1	ILE
2	B	34	ASP
2	B	48	LYS
2	B	58	LYS
2	B	73	THR
2	B	75	LYS
2	B	83	ASN
2	B	88	SER
2	B	94	LYS
1	D	11	SER
1	D	98	MET
1	D	105	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	108	ARG
1	D	132	SER
1	D	176	LYS
1	D	182	THR
1	D	196	ASP
1	D	201	LEU
1	D	207	SER
1	D	223	ASP
1	D	225	THR
1	D	230	LEU
1	D	234	ARG
2	E	0	MET
2	E	34	ASP
2	E	48	LYS
2	E	58	LYS
2	E	73	THR
2	E	75	LYS
2	E	83	ASN
2	E	88	SER
2	E	91	LYS
3	F	6	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	70	HIS
1	A	87	GLN
1	A	174	ASN
1	A	192	HIS
1	A	253	GLN
1	A	262	GLN
1	A	263	HIS
2	B	8	GLN
2	B	13	HIS
1	D	87	GLN
1	D	141	GLN
1	D	174	ASN
1	D	192	HIS
1	D	242	GLN
1	D	253	GLN
1	D	262	GLN
1	D	263	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	2	GLN
2	E	13	HIS
2	E	21	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.26	13 (4%) 36 33	8, 18, 31, 40	0
1	D	275/275 (100%)	0.44	19 (6%) 23 20	9, 20, 33, 41	0
2	B	100/100 (100%)	0.06	2 (2%) 65 62	8, 15, 29, 33	0
2	E	100/100 (100%)	0.17	4 (4%) 42 39	8, 16, 31, 35	0
3	C	9/9 (100%)	1.39	2 (22%) 2 1	8, 22, 29, 29	1 (11%)
3	F	9/9 (100%)	1.53	2 (22%) 2 1	11, 22, 28, 31	1 (11%)
All	All	768/768 (100%)	0.32	42 (5%) 30 27	8, 18, 32, 41	2 (0%)

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	99	TYR	5.5
3	F	5	PHE	4.8
3	C	5	PHE	4.7
1	A	18	GLY	4.7
1	A	86	ASN	4.3
1	D	86	ASN	3.6
2	B	0	MET	3.4
1	A	99	TYR	3.2
2	E	0	MET	3.2
1	D	226	GLN	3.0
2	B	97	ARG	3.0
2	E	44	GLU	3.0
1	A	1	GLY	3.0
3	F	6	PHE	2.9
1	D	192	HIS	2.9
1	D	105	SER	2.9
1	D	17	ARG	2.8
1	A	136	ALA	2.7
1	A	54	GLN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	C	6	PHE	2.6
1	A	196	ASP	2.5
1	D	223	ASP	2.5
1	D	195	SER	2.4
1	D	138	MET	2.4
1	D	196	ASP	2.4
1	A	268	LYS	2.4
1	D	141	GLN	2.3
1	A	223	ASP	2.3
1	D	250	PRO	2.3
1	A	192	HIS	2.3
1	D	193	ALA	2.2
1	A	267	PRO	2.2
1	D	89	GLU	2.2
2	E	97	ARG	2.2
1	D	150	ALA	2.2
1	D	19	GLU	2.2
1	A	17	ARG	2.1
1	D	1	GLY	2.1
1	D	222	GLU	2.1
1	D	54	GLN	2.1
1	A	89	GLU	2.0
2	E	98	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.