



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:16 AM UTC

PDB ID : 4I84 / pdb_00004i84
Title : The crystal structure of the Haemophilus influenzae HxuA secretion domain involved in the two-partner secretion pathway
Authors : Baelen, S.; Dewitte, F.; Clantin, B.; Villeret, V.
Deposited on : 2012-12-03
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	5% 80% 14% • •
1	B	307	4% 77% 11% • •

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4879 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heme/hemopexin-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	15	0
			2292	1422	397	470	3			
1	B	274	Total	C	N	O	S	0	14	0
			2158	1337	380	439	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP P44602
A	-4	HIS	-	expression tag	UNP P44602
A	-3	HIS	-	expression tag	UNP P44602
A	-2	HIS	-	expression tag	UNP P44602
A	-1	HIS	-	expression tag	UNP P44602
A	0	HIS	-	expression tag	UNP P44602
B	-5	HIS	-	expression tag	UNP P44602
B	-4	HIS	-	expression tag	UNP P44602
B	-3	HIS	-	expression tag	UNP P44602
B	-2	HIS	-	expression tag	UNP P44602
B	-1	HIS	-	expression tag	UNP P44602
B	0	HIS	-	expression tag	UNP P44602


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	258	Total	O	0	0
			258	258		
2	B	171	Total	O	0	0
			171	171		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Heme/hemopexin-binding protein



• Molecule 1: Heme/hemopexin-binding protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	38.99Å 70.84Å 104.98Å 90.00° 98.10° 90.00°	Depositor
Resolution (Å)	41.90 – 1.50 41.90 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (41.90-1.50) 99.2 (41.90-1.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.159 , 0.210 (Not available) , 0.202	Depositor DCC
R_{free} test set	4489 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4879	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.53	13/2348 (0.6%)	1.29	7/3166 (0.2%)
1	B	1.42	6/2177 (0.3%)	1.18	14/2928 (0.5%)
All	All	1.48	19/4525 (0.4%)	1.24	21/6094 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	277	ILE	CG1-CD1	-7.33	1.23	1.51
1	B	136	GLN	N-CA	7.15	1.54	1.45
1	B	270	GLU	CA-C	6.88	1.61	1.53
1	B	283	SER	CB-OG	6.10	1.54	1.42
1	B	135	GLY	N-CA	6.09	1.55	1.45
1	A	237	VAL	C-O	6.09	1.30	1.24
1	A	110	ARG	CB-CG	-6.03	1.34	1.52
1	A	9	VAL	CA-CB	-6.00	1.47	1.53
1	B	111	ILE	CA-CB	5.93	1.60	1.54
1	A	124	ARG	CD-NE	-5.72	1.38	1.46
1	A	273	ASN	N-CA	-5.62	1.39	1.46
1	A	139	ASN	N-CA	5.56	1.52	1.46
1	A	281	THR	C-O	5.46	1.30	1.23
1	B	88	VAL	CA-CB	5.36	1.61	1.54
1	A	141	GLY	N-CA	5.31	1.51	1.45
1	A	244	VAL	CA-CB	5.30	1.60	1.54
1	A	76	ALA	C-O	5.11	1.30	1.23
1	A	299	ILE	CA-CB	5.10	1.60	1.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	216	LYS	CA-C	-5.04	1.46	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	124	ARG	NE-CZ-NH2	-8.66	111.41	119.20
1	B	299	ILE	N-CA-C	7.53	119.92	108.71
1	A	124	ARG	NE-CZ-NH1	7.08	128.58	121.50
1	B	300	ASN	N-CA-C	-7.08	95.72	110.80
1	A	269	ASN	N-CA-C	-7.02	99.14	108.34
1	B	299	ILE	CB-CA-C	-6.46	101.88	110.98
1	B	269	ASN	CA-C-N	-6.15	111.02	120.86
1	B	269	ASN	C-N-CA	-6.15	111.02	120.86
1	B	167	ASN	CA-C-N	-5.92	116.30	122.69
1	B	167	ASN	C-N-CA	-5.92	116.30	122.69
1	B	269	ASN	N-CA-C	-5.78	100.09	108.07
1	A	298	ASN	CA-CB-CG	5.63	118.23	112.60
1	A	92	GLN	CA-C-N	-5.53	114.39	122.46
1	A	92	GLN	C-N-CA	-5.53	114.39	122.46
1	B	270	GLU	CA-C-N	-5.50	114.01	119.56
1	B	270	GLU	C-N-CA	-5.50	114.01	119.56
1	B	295	LYS	N-CA-C	-5.49	104.69	111.33
1	B	241	ASN	CB-CA-C	-5.43	108.72	115.89
1	B	232	LEU	N-CA-C	-5.23	99.76	108.34
1	B	300	ASN	N-CA-CB	-5.21	101.69	110.49
1	A	282	GLY	N-CA-C	-5.13	108.09	113.58

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	300	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2292	0	2345	46	0
1	B	2158	0	2181	25	0
2	A	258	0	0	12	0
2	B	171	0	0	5	0
All	All	4879	0	4526	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ARG:HH11	1:A:218:ARG:HG2	1.09	1.05
1:A:291:ASN:HD22	1:B:291[A]:ASN:HD22	1.04	0.98
1:A:291:ASN:ND2	1:B:291[A]:ASN:HD22	1.70	0.89
1:A:218:ARG:HG2	1:A:218:ARG:NH1	1.81	0.87
1:B:252:GLN:HG2	2:B:562:HOH:O	1.72	0.87
1:A:218:ARG:HH11	1:A:218:ARG:CG	1.89	0.85
1:A:291:ASN:HD22	1:B:291[A]:ASN:ND2	1.74	0.83
1:A:194:ASN:HA	1:A:218:ARG:HE	1.46	0.80
1:A:301:GLY:OXT	1:B:301:GLY:O	2.01	0.79
1:A:138[A]:ILE:HD12	1:A:157:ILE:HD12	1.69	0.74
1:A:291:ASN:ND2	1:B:291[A]:ASN:ND2	2.36	0.72
1:A:124:ARG:HD3	2:A:640:HOH:O	1.89	0.72
1:A:252[B]:GLN:HE21	1:A:254:ASN:HD21	1.37	0.71
2:A:658:HOH:O	1:B:301:GLY:HA3	1.90	0.71
1:B:26:ASP:OD2	2:B:570:HOH:O	2.09	0.70
1:B:270:GLU:HB2	1:B:273:ASN:ND2	2.08	0.69
1:A:260:LYS:HE3	2:A:615:HOH:O	1.93	0.68
1:A:89[B]:ILE:HD11	2:A:650:HOH:O	1.93	0.67
1:A:260:LYS:HD2	2:A:561:HOH:O	1.94	0.67
1:B:155:LYS:HE2	1:B:157:ILE:HD11	1.78	0.66
1:A:287:SER:HB3	1:A:290[A]:ILE:CD1	2.28	0.63
1:A:260:LYS:HE2	2:A:633:HOH:O	1.99	0.63
1:A:287:SER:HB3	1:A:290[A]:ILE:HD11	1.81	0.62
1:A:270:GLU:HB2	1:A:273:ASN:ND2	2.15	0.62
1:A:270:GLU:HA	1:A:270:GLU:OE1	1.99	0.62
1:A:138[A]:ILE:HG23	1:A:157:ILE:HB	1.83	0.61
1:A:260:LYS:HG3	2:A:551:HOH:O	2.01	0.60
1:B:299:ILE:HG22	1:B:300:ASN:N	2.19	0.58
1:A:199[A]:ILE:HD13	1:A:225:LEU:HB3	1.88	0.55
1:A:298:ASN:HB3	2:B:501:HOH:O	2.07	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:VAL:HG22	2:A:519:HOH:O	2.05	0.54
1:A:298:ASN:ND2	2:A:622:HOH:O	2.19	0.54
1:A:194:ASN:CA	1:A:218:ARG:HE	2.20	0.52
1:A:155:LYS:HE3	1:A:199[A]:ILE:HG21	1.91	0.52
1:A:7:SER:HB3	1:A:36:TRP:CZ3	2.46	0.51
1:B:6:GLY:O	1:B:37[B]:HIS:HD2	1.94	0.50
1:A:218:ARG:O	2:A:571:HOH:O	2.19	0.50
1:A:217:GLY:C	1:A:218:ARG:NH1	2.70	0.48
1:A:270:GLU:HG2	1:A:273:ASN:HD22	1.78	0.48
1:B:250:ASP:OD1	1:B:276:LYS:HD3	2.13	0.48
1:A:284:LYS:HG2	1:B:300:ASN:OD1	2.14	0.47
1:A:138[A]:ILE:CD1	1:A:157:ILE:HD12	2.40	0.47
1:A:243:LYS:HE2	1:A:245:VAL:CG2	2.44	0.47
1:B:252:GLN:CG	2:B:562:HOH:O	2.45	0.47
1:B:299:ILE:CG2	1:B:300:ASN:N	2.77	0.47
1:B:126:LEU:HD23	1:B:126:LEU:HA	1.66	0.46
1:A:260:LYS:CE	2:A:615:HOH:O	2.60	0.46
1:B:98:VAL:O	1:B:145:ALA:HA	2.14	0.46
1:A:218:ARG:HB2	1:A:220:GLN:HG3	1.98	0.45
1:B:7:SER:HB3	1:B:36:TRP:CZ3	2.52	0.45
1:A:194:ASN:HA	1:A:218:ARG:NE	2.24	0.43
1:A:104:THR:HA	1:A:152:ASN:O	2.19	0.43
1:A:237:VAL:O	1:A:237:VAL:HG23	2.17	0.43
1:A:287:SER:HB3	1:A:290[A]:ILE:HD13	2.00	0.42
1:A:2:ASP:HB3	1:A:45:LYS:HE3	2.01	0.42
1:A:260:LYS:HD2	2:A:551:HOH:O	2.20	0.42
1:A:284:LYS:HE2	1:A:284:LYS:HB2	1.70	0.42
1:A:98:VAL:O	1:A:145:ALA:HA	2.21	0.41
1:B:11:VAL:O	1:B:32[A]:THR:HG23	2.20	0.41
1:A:73:LYS:HE3	1:A:73:LYS:HB3	1.78	0.41
1:B:260:LYS:HG3	2:B:538:HOH:O	2.20	0.41
1:B:269:ASN:O	1:B:270:GLU:C	2.63	0.40
1:B:247:SER:HA	1:B:268:THR:O	2.21	0.40
1:B:255:ASN:HD22	1:B:255:ASN:H	1.69	0.40
1:B:278:THR:HA	1:B:300:ASN:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/307 (100%)	301 (98%)	6 (2%)	0	100	100
1	B	275/307 (90%)	267 (97%)	8 (3%)	0	100	100
All	All	582/614 (95%)	568 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	264/259 (102%)	255 (97%)	9 (3%)	32	7
1	B	245/259 (95%)	240 (98%)	5 (2%)	48	20
All	All	509/518 (98%)	495 (97%)	14 (3%)	39	11

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138[A]	ILE
1	A	138[B]	ILE
1	A	185	SER
1	A	218	ARG
1	A	222	LEU
1	A	239	ASN
1	A	272	LYS
1	A	277	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	284	LYS
1	B	1	ARG
1	B	142	LYS
1	B	196	VAL
1	B	255	ASN
1	B	270	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	66	ASN
1	A	69	GLN
1	A	71	GLN
1	A	152	ASN
1	A	167	ASN
1	A	194	ASN
1	A	239	ASN
1	A	241	ASN
1	A	273	ASN
1	A	291	ASN
1	B	15	ASN
1	B	86	ASN
1	B	239	ASN
1	B	255	ASN
1	B	273	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	296/307 (96%)	-0.27	14 (4%)	36 40	6, 15, 35, 55	15 (5%)
1	B	274/307 (89%)	-0.06	12 (4%)	39 43	7, 20, 35, 47	14 (5%)
All	All	570/614 (92%)	-0.17	26 (4%)	37 41	6, 17, 35, 55	29 (5%)

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	112[A]	SER	4.7
1	A	131	VAL	4.4
1	A	132	VAL	4.0
1	B	301	GLY	3.8
1	A	301	GLY	3.8
1	A	219	ASN	3.7
1	B	126	LEU	3.7
1	B	191	LEU	3.6
1	B	97[A]	ASN	3.5
1	A	182	LEU	3.5
1	B	55[A]	ASN	3.4
1	B	271	PRO	3.2
1	A	221	ALA	3.2
1	B	135	GLY	3.2
1	B	196	VAL	2.9
1	A	217	GLY	2.5
1	B	270	GLU	2.4
1	B	222	LEU	2.4
1	A	181	THR	2.3
1	A	194	ASN	2.3
1	A	115	GLY	2.3
1	A	117	GLY	2.2
1	B	300	ASN	2.2
1	A	270	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	114	ASN	2.1
1	A	183	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.