



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:05 PM UTC

PDB ID : 4I8N / pdb_00004i8n
Title : CRYSTAL STRUCTURE of PROTEIN TYROSINE PHOSPHATASE 1B IN COMPLEX WITH AN INHIBITOR [(4-{(2S)-2-(1,3-BENZOXAZOL-2-YL)-2-[(4-FLUOROPHENYL)SULFAMOYL]ETHYL}PHENYL)AMINO](OXO) ACETIC ACID
Authors : Reddy, S.M.V.V.V.; Rao, K.N.; Subramanya, H.
Deposited on : 2012-12-03
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	354	60% 21% • • 16%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2526 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	298	Total	C	N	O	S	0	0	0
			2434	1541	419	458	16			

There are 34 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-33	MET	-	expression tag	UNP P18031
A	-32	GLY	-	expression tag	UNP P18031
A	-31	SER	-	expression tag	UNP P18031
A	-30	SER	-	expression tag	UNP P18031
A	-29	HIS	-	expression tag	UNP P18031
A	-28	HIS	-	expression tag	UNP P18031
A	-27	HIS	-	expression tag	UNP P18031
A	-26	HIS	-	expression tag	UNP P18031
A	-25	HIS	-	expression tag	UNP P18031
A	-24	HIS	-	expression tag	UNP P18031
A	-23	SER	-	expression tag	UNP P18031
A	-22	SER	-	expression tag	UNP P18031
A	-21	GLY	-	expression tag	UNP P18031
A	-20	LEU	-	expression tag	UNP P18031
A	-19	VAL	-	expression tag	UNP P18031
A	-18	PRO	-	expression tag	UNP P18031
A	-17	ARG	-	expression tag	UNP P18031
A	-16	GLY	-	expression tag	UNP P18031
A	-15	SER	-	expression tag	UNP P18031
A	-14	HIS	-	expression tag	UNP P18031
A	-13	MET	-	expression tag	UNP P18031
A	-12	ALA	-	expression tag	UNP P18031
A	-11	SER	-	expression tag	UNP P18031
A	-10	MET	-	expression tag	UNP P18031
A	-9	THR	-	expression tag	UNP P18031
A	-8	GLY	-	expression tag	UNP P18031
A	-7	GLY	-	expression tag	UNP P18031

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	GLN	-	expression tag	UNP P18031
A	-5	GLN	-	expression tag	UNP P18031
A	-4	MET	-	expression tag	UNP P18031
A	-3	GLY	-	expression tag	UNP P18031
A	-2	ARG	-	expression tag	UNP P18031
A	-1	GLY	-	expression tag	UNP P18031
A	0	SER	-	expression tag	UNP P18031

- Molecule 2 is [(4-{(2S)-2-(1,3-benzoxazol-2-yl)-2-[(4-fluorophenyl)sulfamoyl]ethyl}phenyl)amino](oxo)acetic acid (CCD ID: 1CG) (formula: C₂₃H₁₈FN₃O₆S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	
			34	23	1	3	6	1	


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	58	Total O 58 58	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.89Å 87.89Å 103.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	27.74 – 2.50 27.74 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.8 (27.74-2.50) 93.8 (27.74-2.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.29 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0001	Depositor
R, R_{free}	0.184 , 0.248 0.193 , 0.253	Depositor DCC
R_{free} test set	831 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	40.3	Xtriage
Anisotropy	0.015	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 30.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2526	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 1CG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.63	28/2489 (1.1%)	1.34	9/3355 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	147	GLU	C-O	-6.90	1.15	1.23
1	A	23	ILE	CA-CB	6.68	1.62	1.54
1	A	32	CYS	N-CA	-6.63	1.37	1.46
1	A	234	LEU	C-O	-6.59	1.16	1.24
1	A	286	SER	CA-C	6.31	1.61	1.52
1	A	214	HIS	C-O	6.28	1.31	1.23
1	A	63	ASP	CA-C	6.09	1.60	1.52
1	A	180	PRO	N-CA	6.06	1.54	1.47
1	A	157	GLN	N-CA	-5.91	1.39	1.46
1	A	72	ILE	C-O	-5.83	1.17	1.23
1	A	29	ASP	C-O	-5.75	1.16	1.23
1	A	126	PRO	CA-C	-5.71	1.46	1.52
1	A	217	ALA	C-O	-5.71	1.16	1.24
1	A	44	ASN	C-O	-5.68	1.17	1.24
1	A	31	PRO	N-CA	-5.60	1.40	1.47
1	A	212	VAL	C-O	-5.45	1.17	1.24
1	A	69	ALA	C-O	-5.42	1.17	1.23
1	A	114	MET	CA-C	-5.42	1.46	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	90	ASN	N-CA	5.38	1.53	1.46
1	A	178	THR	N-CA	5.38	1.52	1.46
1	A	212	VAL	CA-CB	-5.27	1.47	1.54
1	A	141	LYS	CA-C	-5.27	1.46	1.52
1	A	214	HIS	CA-C	-5.25	1.46	1.52
1	A	179	TRP	N-CA	-5.15	1.41	1.46
1	A	229	ASP	C-O	5.12	1.30	1.24
1	A	175	HIS	C-O	-5.11	1.18	1.24
1	A	87	PRO	CA-C	-5.06	1.46	1.52
1	A	49	VAL	C-O	5.01	1.29	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	37	LEU	CA-C-N	-8.32	110.37	119.19
1	A	37	LEU	C-N-CA	-8.32	110.37	119.19
1	A	166	GLN	N-CA-C	6.24	120.21	112.47
1	A	92	CYS	N-CA-C	5.67	118.23	111.71
1	A	62	GLU	N-CA-C	-5.64	105.90	112.89
1	A	203	SER	N-CA-C	5.49	118.78	111.75
1	A	15	SER	N-CA-C	5.29	118.86	111.30
1	A	19	ILE	CB-CA-C	-5.06	105.41	112.04
1	A	66	TYR	N-CA-C	5.02	117.75	109.76

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	2	GLU	Peptide
1	A	298	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2434	0	2392	40	0
2	A	34	0	17	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	58	0	0	3	0
All	All	2526	0	2409	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:MET:CE	1:A:246:ILE:HD11	1.81	1.08
1:A:3:MET:HE1	1:A:246:ILE:HD11	1.43	0.98
1:A:143:THR:HG23	3:A:524:HOH:O	1.73	0.87
1:A:6:GLU:OE2	1:A:247:LYS:NZ	2.22	0.72
1:A:48:ASP:O	2:A:401:1CG:H18	1.90	0.71
1:A:3:MET:CE	1:A:246:ILE:CD1	2.67	0.70
1:A:2:GLU:HB3	1:A:4:GLU:HB2	1.73	0.70
1:A:156:ARG:HH11	1:A:173:HIS:HD2	1.45	0.64
1:A:105:ARG:HG2	1:A:105:ARG:HH11	1.64	0.62
1:A:143:THR:CG2	3:A:524:HOH:O	2.37	0.62
1:A:124:TYR:OH	1:A:214:HIS:HE1	1.83	0.61
1:A:140:LEU:HD23	1:A:162:ASN:HA	1.83	0.60
2:A:401:1CG:O11	2:A:401:1CG:H1	2.01	0.60
1:A:3:MET:HE3	1:A:246:ILE:HD11	1.80	0.59
1:A:61:GLN:HE21	1:A:63:ASP:HB3	1.69	0.58
1:A:105:ARG:HH12	1:A:170:GLU:HG2	1.70	0.57
1:A:84:THR:O	1:A:214:HIS:HB2	2.05	0.56
1:A:61:GLN:NE2	1:A:63:ASP:HB3	2.20	0.56
1:A:214:HIS:HD2	1:A:215:CYS:O	1.88	0.56
1:A:97:GLU:OE2	1:A:101:GLU:OE2	2.24	0.54
2:A:401:1CG:H3	3:A:542:HOH:O	2.09	0.53
1:A:20:TYR:CE2	1:A:24:ARG:HD2	2.45	0.52
1:A:42:ASN:C	1:A:42:ASN:OD1	2.52	0.50
1:A:21:GLN:HE22	1:A:24:ARG:HH11	1.59	0.50
1:A:196:PHE:O	1:A:200:GLU:HG3	2.11	0.49
1:A:156:ARG:HH11	1:A:173:HIS:CD2	2.28	0.49
1:A:50:SER:O	1:A:258:MET:HE3	2.12	0.49
1:A:3:MET:HE3	1:A:246:ILE:CD1	2.39	0.47
1:A:111:ASN:HD21	1:A:121:CYS:HB2	1.79	0.46
1:A:124:TYR:OH	1:A:214:HIS:CE1	2.67	0.44
1:A:238:ARG:NH1	1:A:243:SER:O	2.51	0.43
1:A:97:GLU:HG2	1:A:101:GLU:CD	2.43	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:114:MET:HE2	1:A:114:MET:HB2	1.98	0.43
1:A:157:GLN:NE2	1:A:170:GLU:OE1	2.52	0.42
1:A:124:TYR:HH	1:A:214:HIS:HE1	1.67	0.42
1:A:257:ARG:HB3	1:A:260:LEU:HD21	2.01	0.42
1:A:165:THR:OG1	1:A:167:GLU:HG3	2.20	0.42
1:A:2:GLU:C	1:A:4:GLU:N	2.79	0.41
1:A:225:PHE:CD2	1:A:225:PHE:C	2.98	0.41
1:A:191:PHE:HZ	1:A:225:PHE:HA	1.86	0.41
1:A:2:GLU:HB3	1:A:4:GLU:CB	2.49	0.40
1:A:192:LEU:O	1:A:193:ASN:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	296/354 (84%)	279 (94%)	14 (5%)	3 (1%)	12 24

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	MET
1	A	63	ASP
1	A	261	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	271/319 (85%)	255 (94%)	16 (6%)	18	37

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	8	GLU
1	A	21	GLN
1	A	23	ILE
1	A	72	ILE
1	A	114	MET
1	A	118	SER
1	A	127	GLN
1	A	134	ILE
1	A	143	THR
1	A	150	LYS
1	A	190	SER
1	A	207	GLU
1	A	225	PHE
1	A	287	VAL
1	A	299	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	GLN
1	A	61	GLN
1	A	111	ASN
1	A	173	HIS
1	A	193	ASN
1	A	214	HIS
1	A	262	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1CG	A	401	-	34,37,37	2.26	10 (29%)	42,53,53	3.04	13 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1CG	A	401	-	-	14/20/27/27	0/4/4/4

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	1CG	C22-C23	-8.11	1.44	1.54
2	A	401	1CG	C5-N8	-4.79	1.35	1.43
2	A	401	1CG	O26-C23	-3.63	1.20	1.30
2	A	401	1CG	C13-N27	3.46	1.35	1.29
2	A	401	1CG	O25-C23	3.24	1.30	1.22
2	A	401	1CG	C18-N21	-3.17	1.35	1.41
2	A	401	1CG	C29-C28	-2.90	1.35	1.38
2	A	401	1CG	C28-N27	-2.41	1.35	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	1CG	O10-S9	2.16	1.46	1.43
2	A	401	1CG	O24-C22	-2.15	1.19	1.23

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	1CG	O11-S9-O10	-10.11	104.12	118.88
2	A	401	1CG	O30-C13-N27	-8.40	109.64	115.62
2	A	401	1CG	C28-N27-C13	7.63	110.18	104.27
2	A	401	1CG	O30-C29-C28	5.06	111.23	107.80
2	A	401	1CG	O10-S9-C12	4.39	113.35	107.64
2	A	401	1CG	O24-C22-C23	-3.93	116.19	121.36
2	A	401	1CG	C29-C28-N27	-3.86	105.02	108.60
2	A	401	1CG	C1-C2-C3	-3.82	117.79	122.80
2	A	401	1CG	C4-C3-C2	2.97	121.43	118.38
2	A	401	1CG	C14-C15-C20	2.54	125.63	120.90
2	A	401	1CG	C6-C1-C2	2.43	120.87	118.38
2	A	401	1CG	C31-C28-N27	2.27	134.66	128.21
2	A	401	1CG	O26-C23-O25	-2.11	118.88	123.90

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	1CG	O24-C22-C23-O26
2	A	401	1CG	O24-C22-C23-O25
2	A	401	1CG	N21-C22-C23-O25
2	A	401	1CG	C13-C12-S9-O10
2	A	401	1CG	C13-C12-S9-O11
2	A	401	1CG	C14-C12-S9-O10
2	A	401	1CG	C14-C12-S9-O11
2	A	401	1CG	C5-N8-S9-O10
2	A	401	1CG	C5-N8-S9-O11
2	A	401	1CG	C6-C5-N8-S9
2	A	401	1CG	C4-C5-N8-S9
2	A	401	1CG	C5-N8-S9-C12
2	A	401	1CG	C12-C14-C15-C16
2	A	401	1CG	C12-C14-C15-C20

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	1CG	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	298/354 (84%)	-0.29	2 (0%) 84 81	30, 40, 61, 73	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	299	LEU	5.7
1	A	63	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	1CG	A	401	34/34	0.78	0.20	53,91,101,101	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.