



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:27 PM UTC

PDB ID : 1IAR / pdb_00001iar
Title : INTERLEUKIN-4 / RECEPTOR ALPHA CHAIN COMPLEX
Authors : Hage, T.; Sebald, W.; Reinemer, P.
Deposited on : 1999-02-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	129	
2	B	207	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (INTERLEUKIN-4).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			1048	653	192	196	7			

- Molecule 2 is a protein called PROTEIN (INTERLEUKIN-4 RECEPTOR ALPHA CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	188	Total	C	N	O	S	0	0	0
			1517	972	254	282	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	182	ALA	CYS	engineered mutation	UNP P24394
B	1	PHE	MET	engineered mutation	UNP P24394

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total	O	0	0
			63	63		
3	B	167	Total	O	0	0
			167	167		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: PROTEIN (INTERLEUKIN-4)

Chain A: 



• Molecule 2: PROTEIN (INTERLEUKIN-4 RECEPTOR ALPHA CHAIN)

Chain B: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.72Å 67.98Å 108.46Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 2.30	Depositor
% Data completeness (in resolution range)	98.2 (25.00-2.30)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.218 , 0.289	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2795	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/1063	1.03	5/1426 (0.4%)
2	B	0.69	1/1565 (0.1%)	1.11	9/2144 (0.4%)
All	All	0.70	1/2628 (0.0%)	1.08	14/3570 (0.4%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	50	ILE	CA-CB	6.10	1.60	1.53

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	55	GLY	N-CA-C	-8.70	103.29	112.08
2	B	120	ASN	N-CA-C	-7.63	98.28	109.50
2	B	189	GLU	N-CA-C	-6.47	102.00	110.53
1	A	38	ASN	N-CA-C	5.98	118.47	108.73
2	B	155	LEU	N-CA-C	5.93	121.10	111.37
2	B	98	PRO	N-CA-C	5.87	120.21	111.41
2	B	30	SER	N-CA-C	5.70	119.34	112.38
1	A	59	HIS	N-CA-C	5.59	120.79	113.30
2	B	188	SER	N-CA-C	-5.32	103.67	110.53
1	A	24	CYS	N-CA-C	5.29	119.59	113.20
1	A	108	THR	N-CA-C	-5.21	102.77	110.48
2	B	192	PRO	N-CA-C	-5.18	102.98	111.21
2	B	97	LYS	N-CA-C	-5.05	97.83	108.73
1	A	66	LEU	N-CA-C	5.03	119.41	113.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1048	0	1063	32	0
2	B	1517	0	1437	37	0
3	A	63	0	0	6	0
3	B	167	0	0	4	0
All	All	2795	0	2500	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:LYS:HB3	1:A:127:CYS:HB3	1.42	1.00
2:B:3:VAL:HG22	2:B:23:MET:HE3	1.57	0.85
2:B:140:SER:HB2	2:B:170:ILE:HG21	1.59	0.83
1:A:45:PHE:CE1	1:A:120:MET:HE1	2.14	0.81
1:A:15:ASN:O	1:A:19:GLU:HG2	1.88	0.74
1:A:108:THR:HG22	1:A:111:ASN:H	1.53	0.73
1:A:8:GLN:O	1:A:12:LYS:HG3	1.89	0.71
2:B:1:PHE:CD1	2:B:1:PHE:N	2.62	0.68
2:B:144:PRO:C	2:B:146:ASP:H	2.02	0.68
2:B:23:MET:HE1	2:B:85:LEU:HD13	1.80	0.64
1:A:12:LYS:HE2	2:B:131:HIS:HE1	1.64	0.63
2:B:23:MET:SD	2:B:33:LEU:HD11	2.38	0.62
2:B:6:GLU:HG3	3:B:328:HOH:O	2.00	0.60
1:A:50:THR:O	1:A:54:GLN:HG3	2.05	0.57
1:A:94:ALA:HB3	3:A:174:HOH:O	2.05	0.57
2:B:54:ASN:CG	2:B:54:ASN:O	2.48	0.56
2:B:139:TRP:HZ3	2:B:175:ARG:HG3	1.71	0.56
1:A:22:THR:HG22	1:A:24:CYS:H	1.71	0.56
1:A:123:LYS:O	1:A:126:LYS:HB2	2.07	0.55
2:B:4:LEU:N	2:B:22:LYS:O	2.39	0.54
1:A:52:LEU:HD22	1:A:83:LEU:HD22	1.90	0.54
2:B:122:TYR:CZ	2:B:128:LEU:HD12	2.43	0.54
2:B:82:GLN:HG2	3:B:341:HOH:O	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:ARG:NH2	3:A:142:HOH:O	2.41	0.54
2:B:80:ALA:HB2	2:B:85:LEU:HD11	1.90	0.54
2:B:141:GLU:HG3	2:B:173:ARG:HE	1.72	0.54
2:B:144:PRO:O	2:B:146:ASP:N	2.41	0.53
2:B:137:ASN:O	2:B:174:ALA:HA	2.09	0.53
2:B:142:ASN:HD22	2:B:170:ILE:HG23	1.77	0.50
2:B:170:ILE:HG22	2:B:171:SER:N	2.26	0.50
2:B:140:SER:HB2	2:B:170:ILE:CG2	2.37	0.48
2:B:170:ILE:HG22	2:B:171:SER:H	1.79	0.48
1:A:37:LYS:HA	1:A:37:LYS:HE2	1.95	0.48
1:A:101:VAL:HA	3:A:180:HOH:O	2.14	0.48
1:A:31:ASP:HB2	1:A:107:SER:HB2	1.96	0.48
2:B:144:PRO:C	2:B:146:ASP:N	2.72	0.48
2:B:142:ASN:ND2	2:B:170:ILE:HG23	2.29	0.47
2:B:96:VAL:HB	2:B:185:THR:HG22	1.96	0.47
1:A:23:LEU:H	1:A:23:LEU:HD23	1.79	0.47
2:B:54:ASN:HB3	2:B:60:VAL:HG23	1.96	0.47
1:A:97:ASN:ND2	3:A:152:HOH:O	2.46	0.47
1:A:22:THR:CG2	1:A:24:CYS:H	2.27	0.47
1:A:22:THR:HG22	1:A:24:CYS:N	2.29	0.47
1:A:117:LYS:O	1:A:121:ARG:HB2	2.15	0.47
2:B:3:VAL:HG23	3:B:266:HOH:O	2.15	0.46
1:A:13:THR:CG2	1:A:82:PHE:HB3	2.45	0.46
2:B:54:ASN:O	2:B:54:ASN:ND2	2.49	0.46
1:A:41:GLU:OE2	1:A:123:LYS:HD3	2.16	0.45
2:B:77:ASP:OD1	2:B:87:LYS:HG3	2.17	0.45
2:B:177:ARG:HG3	2:B:187:TRP:CE3	2.52	0.45
2:B:115:LEU:HD13	2:B:160:ARG:HG2	1.99	0.44
2:B:14:MET:HG3	3:B:218:HOH:O	2.17	0.44
1:A:33:PHE:C	1:A:35:ALA:N	2.75	0.43
1:A:17:LEU:HD11	1:A:86:LEU:HD22	2.00	0.43
1:A:64:ARG:HE	1:A:64:ARG:HB2	1.64	0.43
2:B:21:TRP:CE2	2:B:59:CYS:HB2	2.54	0.43
1:A:42:LYS:HD3	1:A:94:ALA:O	2.18	0.43
1:A:20:GLN:HG3	1:A:20:GLN:O	2.18	0.43
1:A:71:GLN:HB2	3:A:190:HOH:O	2.18	0.42
1:A:97:ASN:ND2	3:A:176:HOH:O	2.50	0.42
1:A:93:LEU:HD12	1:A:93:LEU:HA	1.87	0.42
2:B:139:TRP:CZ3	2:B:175:ARG:HG3	2.53	0.42
2:B:154:TYR:CG	2:B:155:LEU:N	2.88	0.42
1:A:81:ARG:NH1	2:B:125:ASP:OD2	2.53	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:17:SER:OG	2:B:90:PHE:HE2	2.03	0.42
2:B:2:LYS:HE3	2:B:2:LYS:HB2	1.96	0.40
1:A:43:GLU:O	1:A:47:ARG:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/129 (98%)	119 (94%)	6 (5%)	2 (2%)	7	7
2	B	182/207 (88%)	170 (93%)	9 (5%)	3 (2%)	7	7
All	All	309/336 (92%)	289 (94%)	15 (5%)	5 (2%)	7	7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	144	PRO
1	A	36	SER
2	B	143	ASP
2	B	145	ALA
1	A	128	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/118 (100%)	109 (92%)	9 (8%)	12	16
2	B	167/186 (90%)	151 (90%)	16 (10%)	8	10
All	All	285/304 (94%)	260 (91%)	25 (9%)	9	12

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	THR
1	A	23	LEU
1	A	37	LYS
1	A	38	ASN
1	A	39	THR
1	A	105	ASN
1	A	108	THR
1	A	126	LYS
1	A	127	CYS
2	B	1	PHE
2	B	5	GLN
2	B	13	TYR
2	B	17	SER
2	B	20	GLU
2	B	50	ILE
2	B	54	ASN
2	B	61	CYS
2	B	65	MET
2	B	76	LEU
2	B	84	LEU
2	B	106	VAL
2	B	112	ASP
2	B	142	ASN
2	B	175	ARG
2	B	186	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	59	HIS
1	A	71	GLN
1	A	74	HIS
1	A	78	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	97	ASN
2	B	47	HIS
2	B	54	ASN
2	B	131	HIS
2	B	142	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.